



AMPHIPOD NEWSLETTER

17



AMPHIPOD NEWSLETTER

17



Bibliography by Wim Vader, Tromsø Museum
Produced by Kathleen Conlan,
Canadian Museum of Nature

THE AMPHIPOD NEWSLETTER: WORTH SAVING?

The Amphipod Newsletter has appeared in 16 issues between 1972 and 1986, with first Wim Vader and later Les Walling as editors. Unfortunately, Les has gotten increasingly occupied with other commitments in later years, and AN16 in 1986 has been the last issue to come out. In 1988 editorship was transferred to another colleague, without visible results.

In 1989 Wim Vader took an initiative to revive his brainchild, and got positive reactions from most regional editors, from the Maine conference and from British amphipodologists, polled by Mike Thurston. It was therefore decided to try to bring out AN17 and 18 as quickly as possible, with Kathleen Conlan, Jim Lowry, and Wim Vader as editorial committee. AN17, produced in Ottawa, contains mainly the usual annotated bibliography of amphipod literature, collated by Wim, as well as a questionnaire asking our subscribers once more what exactly they expect to get from AN, what they themselves will be able to contribute to it, and how much they are willing to pay for it. We also ask for correct addresses, and for names and addresses of colleagues that may be interested in receiving AN.

AN18 will be produced in Sydney in November 1990 and will mainly consist of the Index to AN11-17, prepared by Wim Vader and George Crawford. On receipt of the questionnaire we will be able to announce more definite plans for the future of the Amphipod Newsletter, and where and by whom it will be edited and produced.

The present transition has unfortunately not gone completely smoothly. This has resulted in some gaps in the bibliography, especially for 1986 and 1987, and the use of a number of obsolete addresses. We hope to be able to rectify these weaknesses in AN18.

We shall probably be able to produce and send out AN17 and 18 with existing funds, but we shall need more money in 1991.

**Wim Vader
**Ottawa, July 1990

QUESTIONNAIRE

The questionnaire that you received with this issue is meant to provide the editors with four types of information:

1. Do you still want to receive AN, and is your address correct?
2. Do you know of colleagues who may be interested in AN, but do not presently receive it (see list of subscribers in AN16)? Please send us their names and addresses, and we will send them AN17, with this questionnaire, free.
3. What can AN do for you? Is the mix of subjects the right one, does the bibliography satisfy your needs, is your particular field of research suitably covered, etc. etc.?
4. What can you do for AN? Take the subscription, of course, but there is more to it than that. Do you send us your reprints for inclusion in the bibliography, do you contribute to 'News from colleagues', do you help to fill the obvious gaps in the bibliography compiler's access to the literature (speleology, genetics, French and South American journals, Russian literature)?

Please take the time to fill in this questionnaire. It will be of enormous help in charting the waters ahead and finding a crew that can keep AN afloat.


**Wim Vader

ABOUT THE COVER

The cover to AN17 was illustrated by Susan Laurie-Bourque, who has produced many of the amphipod illustrations for Dr. Bousfield and myself over the last 10 years. Susan is a freelance scientific illustrator who works with invertebrates, plants, fishes, mammals, and various ecological themes. The cover illustrates some mating behaviors exhibited by amphipods. On the front cover, left to right, are free swimming male and female *Rhepoxynius* (males have the longer antennae), who have left the sediment to mate-search in the water column. On the sediment a small male *Crangonyx* is copulating with a recently molted female. Within the sediment, the two sexes of *Eohaustorius* are meeting. To the far right a male *Rhinoecetes* is ensuring his parentage by glueing his mates by their shells to his own. In the algae, a "major form" male *Jassa*, who is attending a female in her tube, is confronted by a "minor form" male, who may be acting as a sneak or satellite. Not so easily visible on the front cover, but magnified below, is a female *Dulichia rhabdoplastis* on her rod, which she has accreted to the tip of a spine of *Strongylocentrotus franciscanus*. She is being attended (and defended) by a male until she molts, at which time her cuticle will be sufficiently flexible for her to ovulate and her eggs be fertilized. The rod is still occupied by the offspring of her previous mating.

On the back cover, at the left, is a large male *Orchestia* who has grasped a recently molted female and has dragged her under cover to mate with her. Further to the right is a large male *Gammarus* guarding his mate by carrying her until she molts. The male *Ampithoe* in the alga is also waiting for his mate to molt, and is guarding her in her tube. At the far right a male *Paramoera* is copulating with a female. Unlike the males exhibited to the left, there is little appendage enlargement in males of *Paramoera*, suggesting that mate-guarding and defence is limited.

**Kathleen Conlan



Vllth INTERNATIONAL COLLOQUIUM ON AMPHIPODA

The Vllth International Colloquium on Amphipoda was hosted by Les Watling on September 14-16, 1989. There were 42 registered participants and 33 papers were given. These papers will be published in the journal *Hydrobiologia*. The setting for the meeting was the beautiful, spacious grounds of the Darling Marine Center at Walpole, Maine. Participants were treated to an all-you-can eat lobster and clam bake, tours of local villages, and a post-conference day trip to Monhegan Island for birding, seal watching, and (of course) hopper collecting. Thank you, Les and staff for all your efforts to collect and deliver weary colleagues at odd hours and provide us with a stimulating and most pleasant meeting.

Papers presented

- Marsden, I. A comparison of water loss and gill areas in two supralittoral amphipods from New Zealand.
 Vassilenko, S. Ecologo-physiological characteristics of common caprellid species of the Japan Sea.
 Takeuchi, I. & R. Hirano. Clinging behavior of the Caprellidea (Crustacea, Amphipoda) inhabiting the *Sargassum* zone.
 Thomas, J. Ecology and phylogeny of commensal amphipods - Anamixidae.
 Richardson, A. & R. Swain. Zonation of terrestrial amphipods in maritime western Tasmania.
 Haley, C. & A. Buikema. The role of the amphipod, *Gammarus minus*, in the food webs of two Virginia streams.
 Meijering, M.P.D. Low pH and lack of oxygen as limiting factors for *Gammarus* in hessian brooks and streams.
 Brunel, P. & J.C. Dauvin. Gammaridean recovery in a disturbed suprabenthic sublittoral community from the Lower St. Lawrence estuary.
 Chevrier, A. & P. Brunel. Seasonal and daily densities of suprabenthic Gammaridea over a deep soft bottom in the Bay of Fundy.
 Jazdzewski, K., A. Konopacka & S. Rakusa-Suszczewski. Notes on the biology of some Antarctic peracarids (Amphipoda and Isopoda).
 Jazdzewski, K. & W. Teodorczyk. Amphipod crustaceans as an important component of zoobenthos of the shallow Antarctic sublittoral.
 Quigley, M. & H.A. Vanderploeg. Feeding ecology of the Great Lakes amphipod, *Pontoporeia hoyi*.
 Jones, A. Patterns of abundance of intertidal exoedicerotid amphipods near Sydney, Australia.
 Stock, J. Distribution of anchialine amphipods.
 Krapp-Schickel, T. Comparative ecology of marine Mediterranean and Indonesian amphipods.
 Bhat, U.G. & K. Vamsee. Toxicity of mercury on a gammarid amphipod *Corophium* sp. from the Karwar region, central west coast of India.
 Conlan, K. Sexual dimorphism and mating behaviour of amphipods.
 Aoki, M. Reproductive characteristics of *Sargassum* bed caprellids in Amakusa, Kyushu, Japan.
 Gonzalez, E. Actual state of the Amphipoda taxonomy in Chile.
 Wakabara, Y., F.P. Leite, A.S. Tararam, M.T. Valerio-Berardo & W. Duleba. Gammaridean and caprellidean fauna from Brazil.
 Lowry, J.K. & H.E. Stoddart. Phylogenetic relationships within the Lysianassidae, *sensu stricto*.
 Chapman, J. The possible contribution of human introductions to the tropical Pacific dispersals of gammaridean amphipods.
 Holsinger, John R. What can vicariance biogeography models tell us about the distributional history of subterranean amphipods?
 Vonk, R. Some zoogeographic remarks on Ingolfiellidea from the Canary Islands.
 Takeuchi, I. & Shin-ichi Ishimaru. First record of *Caprogammarus* (Crustacea, Amphipoda) from Hokkaido, Japan.
 Wildish, D.J. & B. Frost. Volumetric growth in gammaridean Amphipoda.

Bousfield, E.L. Convergent morphologies in sand-burrowing members of phyletically unrelated gammaridean superfamilies.
 Oshel, P.E. SEM studies on Macrohectopis branickii from Lake Baikal.
 Boudrias, M. Turning and stopping in swimming amphipods.
 Steele, V.J. The structure and distribution of the type II microtrichs in selected gammaridean amphipods.
 Fong, D. Optic structures of Gammarus minus: comparison between spring and cave populations.
 Coleman, O. Comparative fore-gut morphology of Antarctic amphipods adapted to different food sources.
 Steele, D.H. Is oostegite structure related to ecology or phylogeny?

**Kathleen Conlan

PROFILE OF THE CRUSTACEAN SECTION OF THE CANADIAN MUSEUM OF NATURE

Just as the Canadian Museum of Nature has changed its name - from the National Museum of Natural Sciences, National Museums of Canada - so has the Crustacean Section changed its composition since we last reported on our activities. Ed Bousfield has left us for the more salubrious climate of the Pacific Coast, though he continues to drop in fairly regularly in his ongoing production of revisions to the Pacific coast amphipods.

Chang-tai (Mark) Shih is working on three major hyperiid projects. In collaboration with Dr. H.-E. Gruner, the hyperiid volume of Crustaceorum Catalogus is underway; this has been delayed due to the appointment of HEG to the directorship of the Humboldt University Museum. Significant changes have been made to the format of this volume which will reduce costs and increase accessibility. With Professor Chen Qing-chao, Mark is working on the Hyperiidea of the South China Sea, ultimately to produce a volume in the Fauna Sinica series. Lastly, Mark is reviewing and revising the family Phronimidae, and has already come up with two new species. In his spare time, Mark works on Copepoda: current and future projects include taxonomic reviews of the families of marine Calanoida of Canada, and a survey of the freshwater copepods west of the Rockies.

Diana Laubitz is the Head of the section and tries to protect the others from excessive bureaucratic interference. In between whiles, she is hoping to be able to complete a review of all caprellid genera, and go on to do a revision based on newly discovered or overlooked characters. As a result, she hopes that identification of caprellids will be simplified, and the current proliferation of monotypic genera will be reduced. Future plans include a review of Cyamidae in Canadian waters, with Leo Margolis.

Kathy Conlan is our newest staff member, and is still in the enviable position of establishing her research programs and deciding which of the many fascinating aspects of amphipods she will investigate. Current projects include reproductive biology, particularly mating behaviour in local freshwater gammarids; behaviour of rod-building Podoceridae; effects of iceberg scour, both on behaviour of local scavenging and predatory amphipods and on benthos energetics. Other projects have been or will be: deepwater surveys on the Pacific Coast; the Exxon Valdez Spring (1990) Shoreline Assessment; Antarctic field work; and, of course, this volume of AN.

As a change from amphipods, we have Fahmida Rafi to look after our isopod problems. She is currently describing a new species of the hyperparasitic genus Liriopsis from the Pacific coast, and is starting a revision of the genus Edotia. A major paper revising the Idoteidae of the Canadian Pacific is in press. Fahmida also works on Tanaidacea and Cumacea.

You are reminded that we have an excellent amphipod collection, as well as extensive material of Canadian crustaceans. We welcome research on our collections, either in house or through loans.

**Diana Laubitz

VISITING FELLOWSHIPS AT THE CANADIAN MUSEUM OF NATURE

Visiting Fellowships

The Canadian Museum of Nature offers visiting fellowships to both Canadians and non-Canadians. Applicants should hold a doctorate not more than five years prior to the date of application. Applicants who hold a master's degree obtained within the past eight years and who have at least three years of scientific experience beyond this degree conducting independent research may also be eligible. Applications are also accepted for doctoral graduates who withdrew from active research for the purpose of child bearing and rearing. The fellowships have an annual value of \$32,239, and are subject to Canadian income tax. Fellows will be provided with an allowance towards the cost of travel between the place of residence at the time the award is made and the Canadian Museum of Nature. Spouses and children are eligible to receive additional indemnity. Similar allowances will be provided for the return journey upon termination of the fellowship. The travel allowance is also considered a taxable benefit. Fellows are provided with office space, microscopes, a PC, secretarial service, and some research assistance. Appointments are for one year and renewable for a second year.

The Canadian Museum of Nature has a staff of 200 comprising Collections and Research, Public Programming, and administrative sections. There are 36 research scientists and 49 support staff working in the fields of zoology, botany, paleobiology, and mineral sciences. The Canadian Museum of Nature is situated in Ottawa, the capital of Canada. Metropolitan Ottawa has a population of 500,000. It is located at the junction of the Ottawa, Rideau, and Gatineau Rivers, within a day's drive of Montreal, Toronto, Quebec City, and the northeastern U.S. Ottawa has two universities and numerous government labs.

For more information and applications, please write to:

Visiting Fellowships Office
Natural Sciences and Engineering Research Council
200 Kent Street
Ottawa, Canada
K1A 1H5

and also to:

Assistant Director, Collections and Research
Canadian Museum of Nature
P.O. Box 3443, Stn. D
Ottawa, Canada
K1P 6P4

ACKNOWLEDGEMENTS

Part of the budget that comes from fees for receipt of the Amphipod Newsletter was used for typing and printing of AN17 and production of the mailing labels. I would like to acknowledge the support of the Canadian Museum of Nature for providing envelopes, paying mailing costs, and providing the services of Elemae Lashley who inputted nearly 700 references. As well, the Museum allowed me to set aside my own research program to produce this newsletter, which was a considerably greater time investment than I had anticipated.

**Kathleen Conlan

BIBLIOGRAPHY

This bibliography is set up along the usual AN lines, but because of its long gestation period and my pre-technological background, it is split up in five different parts. There will probably be a gap in the coverage of 1985-87 papers, as I have not yet been able to retrieve what I have sent to previous AN editors, and we had a major computer breakdown in Tromsø in 1988. I hope to be able to supply the missing parts by AN18 or 19.

I am most grateful to all colleagues who sent me reprints of their work. Special thanks, as always, to Jan Stock, who has continued to supply me with lists of references, even during the 'drought' of 1986-1990.

**Wim Vader

OBTAINING THE BIBLIOGRAPHY ON DISKETTE

If you would like a copy of this bibliography for word searches or to add to your reference file, I will copy it for you on Wordperfect 5.1 or in ASCII format, provided that you send me two 5.25 inch double density or one 5.25 inch high density or one 3.50 inch diskette. All diskettes will be formatted in MS-DOS; ASCII files will not have underlines. Please send to:

Dr. Kathleen Conlan
Zoology Division
Canadian Museum of Nature
P.O. Box 3443, stn. D
Ottawa, Ontario K1P 6P4
Canada
(tel. (613) 954-7677)
(fax (613) 954-6439)

BIBLIOGRAPHY

May 1988

- AHSANULLAH, M., M.C. MOBLEY & P. RANKIN, 1988. Individual and combined effects of zinc, cadmium and copper on the marine amphipod Allochrestes compressa. ____ Austr. J. mar. Freshw. Res. **39**, 33-38.
- AINLEY, D.G., W.R. FRASER, C.W. SULLIVAN, J.J. TORRES, T.L. HOPKINS & W.O. SMITH, 1986. Antarctic mesopelagic micronekton: Evidence from seabirds that pack ice affects community structure. ____ Science (N.Y.) **232**, 847-849.
- ALONSO, G., 1989. Gondogeneia thurstoni new species (Amphipoda, Eusiridae) from the south-west Atlantic, Argentina. ____ Crustaceana **56**, 1-7.
- ANDRES, H.G., 1988. Hyaella paramoensis sp. n. aus dem Paramo de Chisaca (Anden). (Crustacea: Amphipoda: Talitroidea). ____ Mitt. hamb. zool. Mus. Inst. **85**, 103-109. (The type locality is in the Colombian Andes).
- ANDRES, H.G., 1988. Zwei neue Acanthonotozomatidae aus der Bransfield Strasse, Antarktis (Crustacea: Amphipoda). ____ Mitt. hamb. zool. Mus. Inst. **85**, 111-120. (Iphimediella paracuticoxa n. sp. and Parapanaploea recessa n. sp.).
- BARLOCHER, F. & S.L. HOWATT, 1986. Digestion of carbohydrates and protein by Gammarus mucronatus Say (Amphipoda). ____ J. exp. mar. Biol. Ecol. **104**, 229-237.
- BAUMER H-P. & G-P. ZAUKE, 1987. Modelling the dynamic relationship between Cd- concentration in Gammarus tigrinus and water temperature. ____ Environm. Technol. Lett. **8**, 529-544.
- BEUGNON, G., M. LAMBIN & A. UGOLINI, 1987. Visual and binocular field size in Talitrus saltator Montagu (Crustacea Amphipoda Talitridae). ____ Monit. zool. ital. **21**, 151-155.
- BEUKEMA, J.J., 1988. An evaluation of the ABC-method (abundance/biomass comparison) as applied to macrozoobenthic communities living on tidal flats in the Dutch Wadden Sea. ____ Mar. Biol. **99**, 425-434.
- BORTKEVICH, L.V., 1988. (Corophiidae (Crustacea, Amphipoda) of the Dniester liman (Ukrainian SSR, USSR)). ____ Vestnik Zool. 1988 (1), 65-68. (In Russian, not seen. Deals with 7 Corophium spp.).
- BOTOSANEANU, L. & J. NOTENBOOM, 1988. Un Amphipode stygobie de la Thaïlande. ____ Annls. Limnol. **24**, 51-59. (Not seen. Bogidiella (B.) thai n. sp.).
- BOTOSANEANU, L. & J.H. STOCK, 1989. A remarkable new genus of cavernicolous Bogidiellidae (Crustacea, Amphipoda) from Thailand. ____ Pp. 171-184 in 'Studies in honour of Dr. Pieter Wagenaar Hummelinck', Found. scient. Res. Surinam & Neth. Antilles **123**, Amsterdam. (Aequigidiella aquilifera n. gen., n. sp. This species is intermediate between Bogidiellidae and Artesiidae, two families to be united).
- BOUSFIELD, E.L., 1988. Crustaceans of tropical seas - The amphipods. ____ Sea Wind **2**, 15-19.
- BOUTIN, C. & M. MESSOULI, 1988. (Metacrangonyx gineti, new species of the genus Metacrangonyx Chevreux, 1909, from a spring in the High-Atlas, Morocco, and the family Metacrangonyctidae new family (Crustacea, Amphipoda). ____ Vie Milieu **38**, 67-84. (In French, not seen. The Metacrangonyctidae n. fam. contains M. and Longipodocrangonyx).
- BOXSHALL, G.A. & K. HARRISON, 1988. New nicothoid copepods (Copepoda: Siphonostomatoida) from an amphipod and from deep-sea isopods. ____ Bull. Br. Mus. nat. Hist. (Zool.) **54**, 285-299. (i.a. Sphaeronella australis n. sp. from Amaryllis sp., Tasmania).
- BROWN, A.F., 1989. Seasonal dynamics of the acanthocephalan Pomphorhynchus laevis (Muller, 1776) in its intermediate and preferred definitive hosts. ____ J. Fish Biol. **34**, 183-194.
- BUCHANAN, J.A., B.A. STEWART & B.R. DAVIES, 1988. Thermal acclimation and tolerance to lethal high temperature in the mountain stream amphipod Paramelita nigroculus (Barnard). ____ Comp. Biochem. Physiol. A **89**, 425-431.
- BUDNIKOVA, L.L., 1989. (Redescription of the species Cerapus erae (Amphipoda, Corophioidea) from the Japan Sea and its position in the system of corophioid amphipods). ____ Zool. Zh. **68**, 48-57. (In Russian. C. erae Bulycheva, 1952 is revived as a valid sp. The genus Paracerapus n. gen. is erected for Cerapus polutovi (type) and C. comparativus. Baracuma alquira is transferred to Cerapus, making Baracuma a junior synonym of Cerapus. The author prefers (contra Barnard) to keep Cerapus and Runanga in the Corophiidae).
- CHARMASSON, S.S. & D.P. CALMET, 1987. Distribution of scavenging Lysianassidae amphipods Eurythenes gryllus in the northeast Atlantic: comparison with studies held in the Pacific. ____ Deep-Sea Res. A. **34**, 1509-1523.
- COLEMAN, C.O., 1989. On the nutrition of two Antarctic Acanthonotozomatidae (Crustacea: Amphipoda). Gut contents and functional morphology of mouthparts. ____ Polar Biol. **9**, 287-294. (Echiniphimedia hodgsoni feeds on sponges, Maxilliphimedia longipes ingests Cnidaria (sea anemones)).
- COLEMAN, O., 1988. Verbreitung und Biologie benthischer Amphipoden. ____ Pp. 51-52 in D.K. Fütterer (ed.). Die Expedition Antarktis VI mit Fls 'Polarstern' in 1987/1988. ____ Ber. Polarforsch. **58**.

- COLEMAN, O. & H. G. ANDRES, 1988. Neue Echiniphimedia - Arten aus der Antarktis (Crustacea: Amphipoda: Acanthonotozomatidae). ____ Mitt. hamb. zool. Mus. Inst. 85, 121-140. (Echiniphimedia gabrielae n. sp., E. waegelei n. sp. and E. barnardi n. sp., all from Antarctic waters. With a key to all Echiniphimedia species).
- COSTELLO, M.J., 1987. Alien amphipod species in Ireland. ____ Porcupine Newsl. 4, 3-7. (Not seen).
- COSTELLO, M.J. & A.A. MYERS, 1989. Observations on the parasitism of Aora gracilis (Bate) (Amphipoda) by Sphaeronella leuckartii Salensky (Copepoda), with a review of amphipod - Sphaeronella associations. ____ J. nat. Hist. 23, 81-91.
- COYLE, K.D. & R.C. HIGHSMITH, 1989. Arctic ampeliscid amphipods: three new species. ____ J. crust. Biol. 9, 157-175. (Deals with Ampelisca erythrorhabdota n. sp. (northern Bering Sea), Byblis robustus n. sp. (northern Bering Sea), and B. frigidus n. sp. (southern Chukchi Sea). A key to the Ampelisca spp. of the area is provided).
- CROXALL, J.P., H.J. HILL, R. LIDSONE-SCOTT, M.J. O'CONNELL & P.A. PRINCE, 1988. Food and feeding habits of Wilson's storm petrel Oceanites oceanicus at South Georgia (Antarctic). ____ J. Zool. (Lond.) 216, 83-102. (Crustacea dominant food, Themisto gaudichaudii dominant crustacean).
- DAUVIN, J.-C., 1988. Bilan des additions aux Inventaires de la faune marine de Roscoff à partir des observations effectuées de 1977 à 1987 en baie de Morlaix avec la signalisation de deux nouvelles espèces d'Amphipodes pour la faune: Ampelisca spooneri Dauvin et Bellan-Santini et Scopelocheirus hopei Costa. ____ Cah. Biol. mar. 29, 419-426.
- DAUVIN, J.-C., 1988. Role du macrobenthos dans l'alimentation des poissons démersaux vivant sur les fond de sédiments fins de la Manche occidentale. ____ Cah. Biol. mar. 29, 445-467. (A most interesting paper, taking advantage of the 'natural experiment' of the 'Amoco Cadiz' oil spill).
- DE PATERNOSTER, I. Kreibohm, 1986. (Fauna associated with ____ Macrocystis pyrifera: quantitative aspects of the populations of Ampithoe femorata (Kroyer) (Amphipoda)). ____ Neotropica 31 (86), 1985, 119-130. (In Spanish).
- DERMOTT, R.M. & K. CORNING, 1988. Seasonal ingestion rates of Pontoporeia hoyi (Amphipoda) in Lake Ontario. ____ Can. J. Fish. aq. Sci. 45, 1886-1895.
- DESBROYERES, D., P. GEISTDORFER, C.L. INGRAM, A. KHRIPOUDOFF & J.P. LAGARDERE, 1985. Repartition des populations de l'épibenthos carnivore. ____ Pp. 233-251 in L. Laubier & Ch. Monniot (eds.). Peuplements profonds du golfe de Gascogne. IFREMER. Paris. (Lysianassid amphipods dominate below 1800 m).
- DEWITT, T.H., 1987. Microhabitat selection and colonization rates of a benthic amphipod. ____ Mar. Ecol. Progr. Ser. 36, 237-250. (Microdeutopus gryllotalpa).
- ELEFTHERIOU, A. & M.R. ROBERTSON, 1988. The intertidal fauna of sandy beaches - A survey of the East Scottish coasts. ____ Scott. Fish. Res. Rep. 38, 1-52.
- ESSINK, K., H.L. KLEEF & W. VISSER, 1989. On the pelagic occurrence and dispersal of the benthic amphipod Corophium volutator. ____ J. mar. biol. Ass. UK 69, 11-15.
- FAUCETT, M.S., 1988. Diet and prey selectivity of scyphomedusae from Port Philip Bay, Australia. ____ Mar. Biol. 98, 503-510.
- FRANCE, R.L. & P.M. STOKES, 1988. Isoetid-zoobenthos association in acid-sensitive lakes in Ontario, Canada. ____ Aquat. Bot. 32, 99-114. ('The microdistribution and abundance of amphipods was dependent on, and significantly correlated with, isoetid biomass').
- FRANZ, D.R., 1989. Population density and demography of a fouling community amphipod. ____ J. exp. mar. Biol. Ecol. 125, 117-136. (Jassa marmorata in New York).
- FRANZ, D.R. & W.H. HARRIS, 1988. Seasonal and spatial variability in macrobenthos communities in Jamaica Bay, New York: an urban estuary. ____ Estuaries 11, 15-28.
- GARCÉS-B., H.A. 1988. Desiccation tolerance of Platorchestia platensis (Kröyer, 1845) (Amphipoda, Talitridae). ____ Rev. Biol. trop. 36, 63-66. (Not seen).
- GOLIKOV, A.A., 1988. (Peculiarities of distribution and ecology of amphipods (Amphipoda, Gammaridea) depending on water dynamics in different parts of the Kandalaksha Bay off the Sonostrov). ____ Issledov. Fauni Morej SSSR 38(46), 110-141. (In Russian).
- GOOCH, J., 1989. Genetic differentiation in relation to stream distance in Gammarus minus (Crustacea, Amphipoda) in Appalachian watersheds. ____ Arch. Hydrobiol. 114, 505-519.
- GREBMEIER, J.M., H.M. FEDER & C.P. McROY, 1989. Pelagic-benthic coupling on the shelf of the northern Bering and Chukchi Seas. 2. Benthic community structure. ____ Mar. Ecol. Progr. Ser. 51, 253-268.
- HACKSTEIN, E., 1988. Die Veränderungen populations dynamischer Parameter bei Gammarus tigrinus Sexton (Crustacea, Amphipoda) als Ausdruck sublethaler Effekte durch die Wechselwirkung von Temperatur und Cadmium-kontaminiertem Futter. ____ Int. Rev. ges. Hydrobiol. 73, 213-228.
- HILL, C., 1988. Life cycle and spatial distribution of the amphipod Pallasea quadrispinosa in a lake in northern Sweden. ____ Holarct. Ecol. 11, 298-304.

- HIRAYAMA, A., 1987. Taxonomic studies on the shallow water gammaridean Amphipoda of West Kyushu, Japan. 7. Melitidae (Melita), Melphidippidae, Oedicerotidae, Philantidae (sic!) and Phoxocephalidae. ____ Publ. Seto mar. biol. Lab. **32**, 1-62. (Deals with Melitalongidactyla n. sp., M. unamoena n. sp., M. pilopropoda n. sp., Melphidippa lineata n. sp., Periculodes longirostratus n. sp., P. pinguis n. sp., Synchelidium rostriopiculum n. sp., Palinotus thomsoni japonicus n. ssp., Heterophilia (recte: Heterophilias) lepas n. sp., Harpiniopsis vadiculus n. sp., Paraphoxus tomiokaensis n. sp., Phoxocephalus prolixus n. sp. and Metaphoxus fulloni).
- HIRAYAMA, A., 1988. Taxonomic studies on the shallow water gammaridean Amphipoda of West Kyushu, Japan. 8. Pleustidae, Podoceridae, Priscomilitaridae, Stenothoidae, Synopiidae, and Urothoidae. ____ Publ. Seto mar. biol. Lab. **33**, 39-77. (Deals with Parapleustes filialis n. sp., Dactylopleustes obsolescens n. sp., Priscomilitaris tenuis n. gen., n. sp. (type of the new family Priscomilitaridae, related to the Isaeidae), Stenothoe valida, Synopia ultramarina, Tiron ovatibasis n. sp., T. galeatus n. sp., Urothoe grimaldii japonica n. ssp. and U. gelasina ambigua n. ssp.).
- HIRAYAMA, A., 1988. A ghost shrimp with four-articulate fifth pereopods (Crustacea: Caprellidea: Phtisicidae) from Northwest Australia. ____ Zool. Sci. Tokyo **5**, 1089-1093. (Quadrisegmentum trianqulum n. gen., n. spec., from Ashmore Reef, NW Australia, collected from the gorgonian Isis hippurus).
- HOBERG, P., 1986. Aspects of ecology and biogeography of Acanthocephala in Antarctic seabirds. ____ Ann. Parasitol. hum. comp. **61**, 199-214.
- HOPKINS, T.L., 1985. The zooplankton community of Croker Passage, Antarctic peninsula. ____ Polar Biol. **4**, 161-170. (Amph. p. 167).
- HOPKINS, T.L., 1987. Midwater food web in McMurdo Sound, Ross Sea, Antarctica. ____ Mar. Biol. **96**, 93-106. (Amph. p.101).
- HOPKINS, T.L. & J.J. TORRES, 1988. The zooplankton community in the vicinity of the ice edge, western Weddell Sea, March 1986. ____ Polar Biol. **9**, 79-87. (Amph. p. 83).
- IKEDA, T., 1988. Metabolism and chemical composition of crustaceans from the Antarctic mesopelagic zone. ____ Deep-Sea Res. **35** A, 1991-2002. (i.a. Cyphocaris sp.).
- IKEDA, T. & H.R. SKJOLDAL, 1989. Metabolism and elemental composition of zooplankton from the Barents Sea during early Arctic summer. ____ Mar. Biol. **100**, 173-183. (i.a. Parathemisto libellula).
- INGLIS, G., 1989. The colonisation and degradation of stranded Macrocyctis pyrifera (L.) C. Ag. by the macrofauna of a New Zealand sandy beach. ____ J. exp. mar. Biol. Ecol. **125**, 203-217. (i.a. Talorchestia quoyana).
- JO, Young Wan, 1988. Talitridae (Crustacea - Amphipoda) of the Korean coasts. ____ Beaufortia **38**, 153-179. (Extensive descriptions of Trinorchestia longiramus n. sp. (E. Korea), Platorchestia crassicornis, P. munmui n. sp. (SE Korea) and P. pachypus. A SEM study revealed that the microstructure of the retinacula may show diagnostic differences between species).
- JO, Young Wan, 1988. Taxonomic studies on Dogielinotidae (Crustacea - Amphipoda) from the Korean coasts. ____ Bijdr. Dierk. **58**, 25-46. (Deals with Haustorioides koreanus n. sp. (Pusan), H. latipalpus n. sp. (Pusan), H. nesogenes n. sp. (Cheonnam), and H. indivisus n. sp. (Kyengki). A key to all Haustorioides spp. is provided).
- KASIMOV, R. Y. & V.I. KRYUCHKOV, 1988. (Effect of nitro lignin on the vital activity of hydrobionts.) ____ Gidrobiol. Zh. **24**, 53-57. (In Russian, not seen. Deals i.a. with Gammarus pulex).
- KEVREKIDIS, T. & A. KOUKOURAS, 1988. Bionomy of the amphipods in the Evros delta (North Aegean Sea). ____ PSZNI MarEcol. **9**, 199-212. (17 spp).
- KEVREKIDIS, T. & M. LAZARIDOU - DIMITRIADOU, 1988. Relative growth and secondary production of the amphipod Gammarus aequicauda (Martynov, 1931) in the Evros delta (N. Aegean Sea) ____ Cah. Biol. mar. **29**, 483-495.
- KOCH, H., 1989. Desiccation resistance of the supralittoral amphipod Traskorchestia traskiana (Stimpson, 1857). ____ Crustaceana **56**, 162-175.
- KORCZYNSKI, R.E., M.J. LAWRENCE & G.E. HOPKY, 1989. Range extensions of some peracaridan crustaceans in the Beaufort Sea. ____ Crustaceana **56**, 215-219. (Haliragoides inermis, Oedicerus saginatus, Andaniella pectinata and Metopa longirama).
- LADLE, M., J.S. WELTON & M.C. BELL, 1987. Sinking rates and physical properties of faecal pellets of freshwater invertebrates of the genera Simulium and Gammarus. ____ Arch. Hydrobiol. **108**, 411-424.
- LALANO-RUEDA, R. & F. GOSSELCK, 1986. Investigations of the benthos of mangrove coastal lagoons in southern Cuba. ____ Int. Rev. ges. Hydrobiol. **71**, 779-794. (Not seen. Corophium insidiosum co-dominant).
- LÖNNE, O.J. & B. GULLIKSEN, 1983. Size, age and diet of Polar Cod, Boreogadus saida (Lepechin, 1773) in ice covered waters. ____ Polar Biol. **9**, 187-191.
- LUCZKOVICH, J.J., 1988. The role of prey detection in the selection of prey by pinfish Lagodon rhomboides (Linnaeus). ____ J. exp. mar. Biol. Ecol. **123**, 15-30.
- MACKIE, G.L., 1989. Tolerance of five benthic invertebrates to hydrogen ions and metals (cadmium, lead, and aluminum). ____ Arch. environm. Contam. Toxicol. **18**, 215-224. (Hyalella most sensitive. 'Populations of H. azteca from low-alkalinity waters can tolerate lower ph-levels than populations from high-alkalinity waters').

- MATTILA, J. & E. BONSDORFF, 1989. The impact of fish predation on shallow soft bottoms in brackish waters (SW Finland): an experimental study. ____ Neth. J. Sea Res. **23**, 69-81. (Using Corophium volutator as one of two prey species).
- MAYER, F.L. & M.R. ELLERSIECK 1988. Experiences with single-species tests for acute toxic effects on freshwater animals. ____ Ambio **17**, 367-375.
- McCAHON, C.P. & D. PASCOE, 1988. Increased sensitivity to cadmium of the freshwater amphipod Gammarus pulex (L.) during the reproductive period. ____ Aq. Toxicol. **13**, 183-194.
- MEADOWS, P.S. & J. TAIT, 1989. Modification of sediment permeability and shear strength by two burrowing invertebrates. ____ Mar. Biol. **101**, 75-82. (i.a. Corophium volutator).
- MEUSY, J.-J. & G.G. PAYEN, 1988. Female reproduction in malacostracan Crustacea. ____ Zool. Sci. Tokyo **5**, 217-265. (A review paper).
- MEYERING M.P.D., 1988. Emissionsbedingte Gewässerveränderungen und ihre Wirkung auf Wassertiere. ____ Pp. 129-134 in D. Manz (ed.). Tagung der Fachgruppe Fischkrankheiten, Giessen Febr. 1988, dtsh. vet-med. Ges., Frankfurt 1988.
- MOORE, P.G. & A.A. MYERS, 1988. An enigma from Australia: a new variation on the corophioid theme (Crustacea: Amphipoda). ____ J. nat. Hist. **22**, 1665-1675. (Aetiopedes gracilis n. gen., n. sp. (Isaeidae) from shallow waters in Bass Straits. It is closest to Amphideutopus, which acc. to the authors also is an isaeid and not a neomegamphopid. Pseudomegamphopus, on the other hand, belongs with the Neomegamphopidae).
- MOORE, P.G. & P.S. RAINBOW, 1989. Feeding biology of the mesopelagic gammaridean amphipod Parandania boeckii (Stebbing, 1888) (Crustacea: Amphipoda: Stegocephalidae) from the Atlantic Ocean. ____ Ophelia **30**, 1-19.
- MORRITT, D., 1987. Evaporative water loss under desiccation stress in semiterrestrial and terrestrial amphipods (Crustacea: Amphipoda: Talitridae). ____ J. exp. mar. Biol. Ecol. **111**, 145-157.
- MORRITT, D., 1988. Osmoregulation in littoral and terrestrial talitroidean amphipods (Crustacea) from Britain. ____ J. exp. mar. Biol. Ecol. **123**, 77-94.
- MYERS, A.A., 1988. The genera Archaeobembos n. gen., Bembos Shoemaker, Protolembos Myers and Globosolembos Myers (Amphipoda, Aoridae, Aorinae) from Australia. ____ Rec. austr. Mus. **40**, 265-322. (An important regional monograph, describing and illustrating the following taxa: Archaeobembos n. gen., monotypic for Autonoephilacantha; Bembos with the spp. B. mollis n. sp. (Lizard Island, Qld), B. ephippium n. sp. (Lizard Isl.), B. ephippium disjuncta n. ssp. (W. Austr.), B. australis, B. quadrimanus (the ssp. mozambicus is questionably different), B. salotiae, B. tridis n. sp. (NSW), B. tridentatus n. sp. (L.I.), B. triangulum n. sp. (Qld), B. bidens n. sp. (Qld), B. tris n. sp. (Vict.), B. dolichomanus n. sp. (Vict.), B. strigilis n. sp. (W. Austr.), B. arkoolus n. sp. (Vict.) and B. gilgi n. sp. (Vict.); Protolembos, with the spp. P. chiltoni, P. murrarum n. sp. (NSW), P. drummondiae n. sp. (Vict.), P. clematis, P. yaranus n. sp. (W.Austr.), P. arinyas n. sp. (Vict.), P. varrucularum, and Globosolembos, with G. ruffoi, G. excavatus and G. lunatus n. sp. (Vict.)).
- NOTENBOOM, J., 1988. Metahadzia uncispina, a new amphipod from phreatic groundwater of the Guadalquivir River basin of Southern Spain. ____ Bijdr. Dierk. **58**, 79-87. (With a discussion of generic limits in the Hadzia - complex).
- NOTENBOOM, J., 1988. Phylogenetic relationships and biogeography of the groundwater-dwelling amphipod genus Pseudoniphargus (Crustacea), with emphasis on the Iberian species. ____ Bijdr. Dierk. **58**, 159-204. (A methodologically important monographic treatment).
- ORTIZ, M. & J. NAZABAL, 1988. (A new amphipod species of the genus Lembos (Crustacea, Amphipoda) from Cuban waters). ____ Rev. Invest. mar. **9**, 29-35. (In Spanish. L. barnardi n. sp. from Ciudad de La Havana province).
- PLESHA, P.D., J.E. STEIN, M.H. SCHIEWE, B.B. MCCAIN & U. VARANASI, 1988. Toxicity of marine sediments supplemented with mixtures of selected chlorinated aromatic hydrocarbons to the infaunal amphipod Rhepoxynius abronius. ____ Mar. environm. Res. **25**, 85-98.
- POSTON, T.M., R.M. BEAN, D.R. KALKWARF, B.L. THOMAS, M.L. CLARK & B.W. KILLAND, 1988. Photooxidation products of smoke generator fuel (SGF) No. 2 fog oil and toxicity to Hyalella azteca. ____ Environm. Toxicol. Chem. **7**, 753-762.
- PRESING, M., 1989. Data to the toxic effect of K-Othrine on crustaceans. ____ Arch. Hydrobiol. **114**, 621-629. (i.a. Gammarus pulex).
- RAFFAELLI, D., A. CONACHER, H. McLACHLAN & C. EMES, 1989. The role of epibenthic crustacean predators in an estuarine food web. ____ Est. coast. Shelf Sci. **28**, 149-160. (i.a. Corophium volutator).
- RAGA, J.A. & C. SANPERA, 1987. Ectoparasites and epizoots of Balaenoptera physalus (L., 1758) in Atlantic Iberian waters. ____ Invest. Pesquera, Barcelona **50** (1986), 489-498. (i.a. Cyamus balaenopteræ, with data on life cycle).
- RAMIREZ, F.C. & M.D. VINAS, 1985. Hyperiid amphipods found in Argentine shelf water. ____ Physis A **43**, 25-37. (Not seen).
- RODGERS, K.A. & R. OLERÖD, 1988. A catalog of zoological specimens collected from Tuvalu (Ellice Islands) by Sixten Bock, 1917. ____ Pacif. Sci. **42**, 300-305.
- ROPER, D.S., S.F. THRUSH & D.G. SMITH, 1988. The influence of runoff on intertidal mudflat benthic communities. ____ Mar. environm. Res. **26**, 1-18. (A New Zealand study).
- ROUCH, R., 1988. Sur la répartition spatiale des Crustacés dans le sous-écoulement d'un ruisseau des Pyrénées. ____ Annls. Limnol. **24**, 213-234.

- SAINTE-MARIE, B. & B.T. HARGRAVE, 1987. Estimation of scavenger abundance and distance of attraction to bait. ____ Mar. Biol. 94, 431-443.
- SCHEEPMAKER, M., F.v.d. MEER & S. PINKSTER, 1988. Genetic differentiation of the Iberian amphipods Gammarus ibericus Margalef, 1951 and G. gauthieri S. Karaman, 1935, with reference to some related species in France. ____ Bijdr. Dierk. 58, 205-226.
- SCHNEPPENHEIM, R. & R. WEIGMANN-HAAS, 1986. Morphological and electrophoretic studies of the genus Themisto (Amphipoda: Hyperiidea) from the South and North Atlantic. ____ Polar Biol. 6, 215-225.
- SCIPIONE, M.B. & E. FRESI, 1984. Distribution of amphipod crustaceans in Posidonia oceanica (L.) Delile foliar stratum. ____ Pp. 319-329 in C.F. Boudouresque, A.J. de Grissac & J. Olivier (eds.). International workshop on Posidonia oceanica beds. G.S. Posidonie, Marseille. (Not seen).
- SKET, B., 1988. (Zoogeography of the freshwater and brackish Crustacea in the Kvarner-Velebit islands (Nordwest Adriatic, Yugoslavia). ____ Biol. Vestnik 36, 63-76. (In Slovenian, not seen).
- SKET, B., 1988. Fauna of the hypogean waters of the San Andres (St. Andrews) Islands, Colombia. ____ Biol. Vestnik 36, 77-82. (Not seen).
- STEPHENSON, M. & G.L. MACKIE, 1988. Multivariate analysis of correlations between environmental parameters and cadmium concentrations in Hyaella azteca (Crustacea: Amphipoda) from Central Ontario lakes. ____ Can. J. Fish. aq. Sci. 45, 1705-1710.
- STOCK, J.H., 1988. Stygofauna of the Canary Islands, 9. The amphipod genus Pseudoniphargus (Crustacea) in the Canary Islands. ____ Bijdr. Dierk. 58, 47-78. (Deals with P. porticola n. sp., P. longicauda n. sp., P. fontinalis n. sp. and P. unispinosus n. sp., all from Tenerife, P. cupicola n. sp. and P. multidentis n. sp. from La Palma, P. gomerae n. sp. from Gomera and P. salinus n. sp. from Hierro, all in the western Canary Islands).
- THOMAS, J.D. & J. L. BARNARD, 1989. Gammaropsis arawakia a new species of marine Amphipoda (Crustacea) from Jamaica. ____ Proc. biol. Soc. Wash. 102, 89-94.
- TIMMS, B.V., U.T. HAMMER & J.W. SHEARD, 1986. A study of benthic communities in some saline lakes in Saskatchewan and Alberta, Canada. ____ Int. Rev. ges. Hydrobiol. 71, 759-777.
- TUCKER, M.J., 1988. Temporal distribution and brooding behavior of selected benthic species from the shallow marine waters off the Vestfold Hills, Antarctica. ____ Hydrobiologia 165, 151-159. (With data on i.a. Heterophoxus videns, Orchomene franklini, Ampelisca sp. and A. barnardi).
- TUCKER, M.J. & H.R. BURTON, 1987. A survey of the marine fauna in shallow coastal waters off the Vestfold Hills and Rauer Islands, Antarctica. ____ Anare Res. Notes 55, 1-24. (Amph., - Lowry det. - pp. 12-13).
- ULIAN, G.B. & E.G. MENDES, 1988. Tolerances of a land amphipod, Talitrus pacificus Hurley, 1955, towards temperature and humidity variations and immersion in water. ____ Rev. brasil. Biol. 48, 179-188. (Not seen).
- UNDERWOOD, A.J. & P.H. VERSTEGEN, 1988. Experiments on the association between the intertidal amphipod Hyaella media Dana and the limpet Cellana tramoserica (Sowerby). ____ J. exp. mar. Biol. Ecol. 119, 83-98.
- VINOGRADOV, G.M., 1988. (Life forms of pelagic amphipods). ____ Zool. Zh. 67, 1765-1775. (In Russian).
- WESAWSKI, J.M., M. ZAJACZKOWSKI, S. KWASNIEWSKI, J. JEZIERSKI & W. MOSKAL, 1988. Seasonality in an Arctic fjord ecosystem: Hornsund, Spitsbergen. ____ Polar Res. 6, 185-189.
- WIEBE, P.H., N. COPLEY, C. VAN DOVER, A. TAMSE & F. MANRIQUE, 1988. Deep-water zooplankton of the Guaymas Basin hydrothermal vent field. ____ Deep-Sea Res. A 35, 985-1014.
- WILDISH, D.J., 1988. Ecology and natural history of aquatic Talitroidea. ____ Can. J. Zool. 66, 2340-2359.
- WILLIAMS, W.D. & J.L. BARNARD, 1988. The taxonomy of crangonyctoid Amphipoda (Crustacea) from Australian fresh waters: foundation studies. ____ Rec. austr. Mus., Suppl. 10, 1-180. (The long-awaited monographic review of this fascinating fauna, with an extensive discussion of the taxonomy of the group Crangonyctoidea. The following taxa are described: Paramelitidae, with Austrogammarus, incl. A. australis (type), A. smithi n. sp. (Tasmania = G. australis s. Smith 1909), A. haasei, A. saycei n. sp. (Victoria), A. spinatus n. sp. (Victoria) and A. multispinatus n. sp. (Victoria); Austrocrangonyx, with A. barringtonensis (type) and A. hynesi n. sp. (NSW); Antipodeus n. gen., with Gammarusantipodeus (type), Neoniphargus wellingtoni, N. niger, Gammarus ripensis, Niphargus mortoni and A. franklini n. sp. (Tasmania); Hurleya, with H. kalamundae; Uroctena (with U. affinis (revived, type), U. westralis, U. setosa and U. yellandi); Giniphargus, with G. pulchellus and Protocrangonyx, with P. fontinalis. The Neoniphargidae contain Tasniphargus n. gen., for T. tyleri n. sp. (Tasmania); Neoniphargus, with N. thomsoni, N. spenceri, N. fultoni, N. obrieni, N. alpinus, N. exiguus and N. tasmanicus; Wesniphargus n. gen., for Neoniphargus nichollsii and Yulia n. gen., for Neoniphargus yuli. The Perthiidae n. fam. are monotypic for Perthia, with Neoniphargus branchialis (type) and P. acutitelson).
- YAMATO, S., 1987. Four intertidal species of the genus Melita (Crustacea: Amphipoda) from Japanese waters, including descriptions of two new species. ____ Publ. Seto mar. biol. Lab. 32, 275-302. (M. rylovae, M. koreana, M. nagatai n. sp. and M. bingensis n. sp., all from the Seto Inland Sea of Japan).
- ZEIDLER, W., 1988. A redescription of Afrochiltonia capensis (K.H. Barnard, 1916) with a review of the genera of the family Ceinidae (Crustacea, Amphipoda). ____ Ann. S. Afr. Mus. 98, 105-119. (The genus Austrochiltonia is resurrected for the Australian species, and a key to all ceinid genera provided).

ZEIDLER, W., 1989. A new species of Melita (Crustacea: Amphipoda: Melitidae) from northern New South Wales, with a note on the genus Abludomelita Karaman, 1981. ____ Proc. linn. Soc. NSW 110 (1988), 327-338. (Melita plumulosa n. sp. from a coastal freshwater lake. The genus Abludomelita has been characterized by plesiomorphic characters and Zeidler is therefore reluctant to accept it).

BIBLIOGRAPHY December 1988

- AARSET, A.V. & T. AUNAAS, 1987. Physiological adaptations to low temperature and brine exposure in the circumpolar amphipod Gammarus wilkitzkii. ____ Polar Biol. 8 129-134.
- AARSET, A.V. & K-E. ZACHARIASSEN, 1988. Low temperature tolerance and osmotic regulation in the amphipod Gammarus oceanicus from Spitsbergen waters. ____ Polar Res. 6, 35-42.
- ABEL, T. & F. BÄRLOCHER, 1988. Uptake of cadmium by Gammarus fossarum (Amphipoda) from food and water. ____ J. appl. Ecol. 25, 223-231.
- ALLAN, J.D., G.N. HERBST, R.ORTAL & Y. REGEV, 1988. Invertebrate drift in the Dan river, Israel. ____ Hydrobiologia 160, 155-164.
- ALONSO, G.M., 1987. (New records of marine amphipods (Amphipoda, Gammaridea) for Argentina). ____ Physis (Buenos Aires) A 44 (1986), 59-66. (In Spanish. Deals with Ampithoe valida, Cymadusa filosa, Ischyrocerus anquipes, Melita palmata & Orchestia gammarellus).
- ALONSO, G.M., 1987. (Systematic studies of three Lysianassidae (Amphipoda, Gammaridae) from Argentina). Physis (Buenos Aires) A 45, 1-10. (In Spanish. Deals with Amaryllis macrophthalma, Tryphosites chevreuxi and Tmetonyx serratus).
- ALONSO, G.M., 1987. (On the occurrence of Parawaldeckia kidderi (Smith) (Amphipoda, Lysianassidae) in the Argentine Sea). Physis (Buenos Aires) A 45, 17-20. (In Spanish).
- ANDRES, H-G., 1987. Die Gammaridea der 76. Reise von FFS 'Walther Herwig' mit Beschreibung von Parachevreuxiella lobata gen. n. und sp. n. (Crustacea, Amphipoda). ____ Mitt. hamb. zool. Mus. Inst. 84, 95-103. (Deepwater pelagic trawls from the NE Atlantic. Of the 6 lysianassids and 1 stegocephalid caught P. lobata n. gen., n. sp. (Lysianassoidea) was found on the platytroctid fish Normichthys aperiatus at 53°03' N, 16° 36' W. Parachevreuxiella is close to Chevreuxiella. C. obensis is tentatively transferred to Danaella).
- ARRESTI, A., J.C. ITURRONDOBEITIA & A. ROLLO, 1987. (Ecological study of the amphipods of the bay of Bilbao). ____ Cuadern. Invest. biol. 10, 51-88. (In Spanish, not seen).
- AVDEEV, V.V. & N. V. AVDEEVA, 1986. (Occurrence of cestodes of the order Tetraphyllidea in marine invertebrates and fishes, and the probable schemes of their development). ____ Parazitologiya (Leningrad) 20, 448-454. (In Russian, not seen).
- BÄRLOCHER, F., J. GORDON & R.J. IRELAND, 1988. Organic composition of seafoam and its digestion by Corophium volutator (Pallas). ____ J. exp. mar. Biol. Ecol. 115, 179-186.
- BARNARD, J.L. & J.B.R. AGARD, 1986. A new species of Ampelisca (Crustacea, Amphipoda) from Trinidad. ____ Bull. mar. Sci. 39, 630-636. (A. paria n. sp.).
- BARNARD, J.L. & M.M. DRUMMOND, 1987. Rectification of Amphoedicerus willisi Fearn-Wannan (1968); genus and species removed to Paramoera Miers (1875). ____ Proc. R. Soc. Victoria 99, 13-18.
- BARNARD, J.L. & M.M. DRUMMOND, 1987. A new marine genus, Doowia, from eastern Australia (Amphipoda, Gammaridea). ____ Proc. R. Soc. Victoria 99, 117-126. (Doowia n. gen. 'forms a separate branch in the complex of families Oedicerotidae, Exoedicerotidae and Paracalliopiidae', but the authors refrain from erecting a new family. The type is Doowia cooma n. sp. from NS Wales beaches; further species D. dexterae n. sp. from a Queensland beach).
- BARNARD, J.L. & G.S. KARAMAN, 1987. Revision in classification of Gammaridean Amphipoda (Crustacea), part 3. ____ Proc. biol. Soc. Wash. 100, 856-873. (A new round of 'armchair revisions'. The Ochlesinae are reduced to subfamily rank sub Acanthonotozomatidae, and a key to all genera provided. Meraldia n. gen. (A. Ochlesinae) is erected for Ochlesis meraldj. The family Cardenioidae, close to the synopiids, comprises only the type genus Cardenio. Also the Clarenciidae n. fam. are monotypic, for Clarencia. New genera in the Eusiridae are Abdia n. gen. (for Atylopsis latipalpus), Manerogeneia n. gen. (for Pontogeneiella maneroo), Membrilopus n. gen. (for Metaleptamphopus membrisetata), Nasageneia n. gen. (for Pontogeneia nasa (type) and P. guinsan) and Whangarusa n. gen. (for Panoploea translucens). h te baab, Photis geniculata is transferred to Cheiriphotis. A key to liljeborgiid genera is provided and Isipingus n. gen. erected for Liljeborgia epistomata. New lysianassid genera are Bruunosa n. gen. (for Tryphosa bruuni), Cedrosella n. gen. (for ? Ambasiopsis fomes), Cicadosa n. gen. (for Anonyx cicadoides), Galathella n. gen. (for Schisturella galathea), Lepiduristes n. gen. (for ? Uristes lepidus), and Rimakoroga n. gen. (for Pseudokoroga rima). In the as yet (for nomenclatorial reasons) not formally named nuuanuid family-group the genera Pherusa, Pherusana, Nuuanu, Cottesloe and Valettiella (transferred from Lysianassidae) are all submerged in Gammarella. The monotypic new families Pseudamphilochidae Schellenberg, 1931 (revived) (for Pseudamphilochus) and Bolttsiidae (for Bolttsia) are erected. Stegosoladius n. gen. (Stegocephalidae) is monotypic, type Andaniotes simplex. New genera in the Stenothoidae are Aurometopa n. gen. (for Metopoides aurorae), Knysmetopa n. gen. (for Parametopa grandimana), Torometopa n. gen. (for Metopa crenatipalmata (type), M. aequalis, M. antarctica, M. carinata, M. compacta, M. crassicornis, M. dentimana, M. palmata, M. parallelocheir, M. perlata, M. porcellana and M. stephensi), Vonimetopa n. gen.

- (for Metopella dubia (type), M. brazhnikovii, M. schellenbergi, M. shoemakeri and M. zernovi), and Zaikometopa n. gen. (for Metopelloides erythrophthalmus). Finally, in the Temnophliantidae, Hystriphlias n. gen. is erected for Temnophlias hystrix.
- BARNARD, J.L. & J. D. THOMAS, 1988. Vadosiapus copacabanus, a new genus and species of Exoedicerotidae from Brazil (Crustacea, Amphipoda). ____ Proc. biol. Soc. Wash. **103**, 366-374. (With keys to Exoedicerotid genera, and to Bathyporeiapus and Vadosiapus species. V. copacabanus was indeed collected from the famous beach in Rio de Janeiro).
- BARNARD, J.L. & J.D. THOMAS, 1988. Ipanemidae, new family, Ipanema talpa, new genus and species, from the surf zone of Brazil (Crustacea: Amphipoda; Haustoriidea). ____ Proc. biol. Soc. Wash. **101**, 614-621. (from Rio de Janeiro).
- BARNARD, J.L., J.D. THOMAS & K.B. SANDVED, 1988. Behavior of gammaridean Amphipoda: Corophium, Grandidierella, Podocerus and Gibberosus (American Megaluropus) in Florida. ____ Crustaceana, Suppl. **13**, 234-244.
- BARNES, R.S.K., 1987. Coastal lagoons of East Anglia, U.K. ____ J. coast. Res. **3**, 417-427.
- BAUDET, J., 1986. Dynamique d'une population d' Orchestia cavimana Heller, 1865 (amphipodes Talitridae) dans l'ouest de la France: croissance et cycle reproducteur. ____ Bull. Ecol. **17**, 97-108.
- BAYLISS, D. & R.R. HARRIS, 1988. Chloride ion regulation in the freshwater amphipod Corophium curvispinum and acclimatory effects of external climate. ____ J. comp. Physiol. B **158**, 81-90.
- BELL, J. D. & M. WESTABY, 1986. Abundance of macrofauna in dense seagrass is due to habitat preference, not predation. ____ Oecologia **68**, 205-209.
- BELLAN-SANTINI, D. & J.C. DAUVIN, 1986. Morphologie et microstructure d'une formation cuticulaire énigmatique chez la nouvelle espèce du genre Ampelisca (sic!), A. remorai. ____ Cah. Biol. mar. **27**, 328. (Abstract only).
- BELLAN-SANTINI, D. & J. C. DAUVIN, 1988. Actualisation des données sur l'écologie, la biogéographie et la phylogénie des Ampeliscidae (Crustacés-Amphipodes) atlantiques après la révision des collections d' E. Chevreux. ____ Aspects récents de la Biologie des Crustacés. IFREMER Actes Coll. **8**, 207-216.
- BELLAN-SANTINI, D. & J.C. DAUVIN, 1988. Eléments de synthèse sur les Ampeliscidae du nord-est Atlantique. ____ Crustaceana, Suppl. **13**, 20-60. (A most useful review).
- BELLAN-SANTINI, D., A. WILLISIE & A. ARNOUX, 1986. Distribution comparée des crustacés amphipodes de la matée d'herbier de posidonies mort et vivant. ____ Rapp. P.-v. Reun. Comm. int. Explor. scient. Mer Méditerran. **30**, 8.
- BEUGNON, G., M. LAMBIN & A. UGOLINI, 1987. Visual and binocular field size in Talitrus saltator Montagu (Crustacea Amphipoda, Talitridae). ____ Monit. zool. ital. **21**, 151-155.
- BIRSHTJEV, Y.A., 1985. (Genesis of freshwater, cave and deep-sea fauna). ____ Nauka Moskva, 247 pp. (In Russian, not seen. Can anybody furnish an abstract of this book?).
- BLANCHET-TOURNIER, M-F., 1987. Mise en évidence d'une activité neurohormonale inhibitrice de la vitellogénèse chez l'amphipode Orchestia gammarella. ____ Can. J. Zool. **65**, 1917-1922.
- BLANCHET-TOURNIER, M-F & M. CAZES, 1985. Effets de l'implantation d'une glande androgène de Decapode (Carcinus maenas) sur le fonctionnement de l'ovaire d'une Amphipode pubère (Orchestia gammarella) ____ CR Acad. Sci. Paris D **15**.
- BLINN, D.W., R.W. DAVIES & B. DEHDASTI, 1987. Specialized pelagic feeding by Erpobdella montezuma (Hirundinea). ____ Holarct. Ecol. **10**, 235-240. (Diet is 90% Hyalella montezuma).
- BLINN, D.W., N.E. GROSSNICKLE & B. DEHDASHTI, 1988. Diel vertical migration of a pelagic amphipod in the absence of fish predation. ____ Hydrobiologia **160**, 165-171. (Hyalella montezuma from Arizona, USA).
- BLINN, D.W., C. PINNEY & V.T. WAGNER, 1988. Intra-specific discrimination of amphipod prey by a freshwater leech through mechanoreception. ____ Can. J. Zool. **66**, 427-430. (Erpobdella montezuma distinguishes between adult and juvenile Hyalella montezuma).
- BOLT, S.R.I., 1986. Ecological, behavioural and physiological observations on under-ice populations of Arctic amphipods associated with salinity anomalies. ____ Progr. Underwat. Sci. **11**, 127-135.
- BOLT, S.R.I., 1986. Variation of water permeability in selected euryhaline amphipods. ____ Porcupine Newsl. **3** (7), 78-100. (Not seen).
- BOROWSKY, B., 1986. Laboratory observations on the pattern of reproduction of Elasmopus levis (Crustacea: Amphipoda). ____ Mar. Behav. Physiol. **12**, 245-256.
- BOROWSKY, B., C.E. AUGELLI & S.R. WILSON, 1987. Towards chemical characterization of waterborne pheromone of amphipod crustacean, Microdeutopus gryllotalpa. ____ J. Chem. Ecol. **13**, 1673-1680.
- BOROWSKY, B. & R. BOROWSKY, 1987. The reproductive behavior of the amphipod crustacean Gammarus palustris (Bousfield) and some insights into the nature of their stimuli. ____ J. exp. mar. Biol. Ecol. **107**, 131-144.

- BORTKEVICH, L.V., 1987. (Distribution and ecology of Corophium orientale (Amphipoda, Corophiidae) in the Black Sea estuaries). ____ Vestnik Zool. 1987 (3), 72-74. (In Russian, not seen).
- BORTKEVICH, L.V., 1987. (Ecology and production of Corophium curvispinum in the estuarine section of rivers of the northwestern Black Sea coast) ____ Gidrobiol. Zh. 23 (6), 91-93. (In Russian, not seen).
- BORTKEVICH, L.V., 1988. (Corophiidae (Crustacea, Amphipoda) of the Dnieper Bay). ____ Vestnik Zool. 1988 (1), 65-68. (In Russian, not seen. Seven Corophium spp. are dealt with).
- BOUDRIAS, M.A. & A.G. CAREY, 1988. Life history patterns of Pseudalibrotus litoralis (Crustacea: Amphipoda) on the inner continental shelf, SW Beaufort Sea. ____ Mar. Ecol. Progr. Ser. 49, 249-257.
- BOURNAUD, M., H. TACHAT, A-L. ROUX & Y. AUDA, 1987. The effects of seasonal and hydrological influences on the macroinvertebrates of the Rhone River, France. ____ Arch. Hydrobiol. 109, 287-304.
- BOUSFIELD, E.L., 1988. Ordered character states as a basis for phyletic classification within the Amphipoda. ____ Crustaceana, Suppl. 13, 279-280.
- BOUTIN, C. & COINEAU, 1988. Pseudoniphargus maroccanus n. sp. (Subterranean amphipod), the first representative of the genus in Morocco. Phylogenetic relationships and paleobiogeography. ____ Crustaceana, Suppl. 13, 1-19. (With a cladistic analysis of relationships within the genus, and a discussion of evolution and paleobiogeography).
- BOUTIN, C. & M. MESSOULI, 1988. Longipodacrangonyx maroccanus, n. gen., n. sp., nouveau représentant du groupe Metacrangonyx dans les eaux souterraines du Maroc. ____ Crustaceana, Suppl. 13, 256-271. (From the Marrakesh area).
- BRAZDA, J. & J. TEREK, 1987. (Makrozoobenthos in tributaries of the lake Izra (Czechoslovakia)). ____ Biologia (Bratislava) 42, 557-564. (In Slovak, not seen. Amph. dominant).
- BREY, T., H. RUMOHRE & S. ANKAR, 1988. Energy content of macrobenthic invertebrates. General conversion factors from weight to energy. ____ J. exp. mar. Biol. Ecol. 117, 271-278.
- BROWN, P.W. & L.H. FREDRICKSON, 1966. Food habits of breeding White-winged Scoters. ____ Can. J. Zool. 64, 1652-1654. (These ducks ate mainly Hyaella azteca in their Saskatchewan breeding area).
- BUCHANAN, J.A., B.A. STEWART & B.A. DAVIES, 1988. Thermal acclimation and tolerance to lethal high temperature in the mountain stream amphipod Paramelita nigroculus (Barnard). ____ Comp. Biochem. Physiol. A 89, 425-432.
- BUCKLIN, A.A., R.R. WILSON & K.L. SMITH, 1987. Genetic differentiation of seamount and basin populations of the deep-sea amphipod Eurythenes gryllus. ____ Deep-Sea Res. 34, 1795-1810.
- BUSHUEVA, I.V., 1988. (Two new species of antarctic scuds of the family Stenothoidae (Amphipoda, Gammaridea) ____ Zool. Zh. 67, 511-517. (In Russian. Describes Prometopa dorsoundata n. sp. and Proboloides bellansantiniae n. sp., both from the Davis Sea. The genus Prometopa is resurrected, and P. tuberculata and P. dorsoundata compared).
- CAREY, A.G. & M.A. BOUDRIAS, 1987. Feeding ecology of Pseudalibrotus (= Onisimus) litoralis Kröyer (Crustacea: Amphipoda) on the Beaufort Sea inner continental shelf. ____ Polar Biol. 8, 29-33.
- CEJAS, J.R., A. BRITO & G. LORENZO, 1983. (On some gammarideans (Crustacea Amphipoda) new for the marine fauna of the Canary islands). ____ Viera 12, 317-328. (In Spanish, not seen).
- CELLAT, B. & M. BOURNAUD, 1988. Dynamique spatio-temporelle des déplacements de macroinvertebrés dans une grande rivière. ____ Can. J. Zool. 66, 352-363. (river Rhone, i.a. Gammarus spp.).
- CHAMBERS, M.R., 1987. The status of the alien amphipod Gammarus tigrinus (Sexton, 1939) in Friesland twenty five years after its introduction into the Netherlands. ____ Bull. zool. Mus. A'dam 11, 65-68. (A strong decline in abundance, though not in area, compared to the early seventies).
- CHAPMAN, J.W., 1988. Invasions of the northeast Pacific by Asian and Atlantic gammaridean amphipod crustaceans, including a new species of Corophium, ____ J. crust. Biol. 8, 364-382. (Deals with Ampelisca abdita, Corophium alienense n. sp. (S. Francisco Bay), Melita nitida and Parapleustes derzhavini).
- CHARMASSON, S.S. & D.P. CALMET, 1987. Distribution of scavenging Lysianassidae amphipods Eurythenes gryllus in the northeast Atlantic: comparison with studies held in the Pacific. ____ Deep-Sea Res. 34 A, 1509-1524.
- CHEREPANOV, V.V., M.A. ARIPOVA, T.D. EVSTIGNEEVA, V.G. KRASNOV, I.N. NADELYAEV, A.L. NOVITSKIY & Z.V. SLUGINA, 1987. (Reactions of Baikal organisms to low concentrations of toxicants). ____ Izv. So An SSSR (Biol. Nauk) 3, 97-103. (In Russian, not seen. Deals with i.a. two Eulimnogammarus spp. The endemic Baikal animals are in general highly sensitive to toxicants).
- CHESUNOV, A.V. & M.V. PLETNIKOVA, 1986. (New data on commensal nematodes from the genus Gammarinema Kinne et Gerlach, 1953 (Chromadoria, Monhysterida)). ____ Byull. Mosk. Obshch. Ispyt. Priv. (Otd. Biol.) 91, 73-83. (In Russian, not seen).

- CHERGUL, H. & E. PATTEE, 1988. The impact of benthic invertebrates on the breakdown of poplar leaves in the network of a large European river. ____ Arch Hydrobiol. 113, 15-25. (The river Rhone near Lyon, France).
- CIOLPAN, O., 1987. The reproduction of Dikerogammarus villosus (Sow.) Mart. (Crustacea, Amphipoda) in the Iron Gate artificial lake (Bahna Gulf, 1983-1984). ____ Rev. roum. Biol. 32, 70-74.
- CLAMP, J.C., 1987. Five new species of Lagenophrys (Ciliophora, Peritricha, Lagenophryidae) from the United States, with observations on their developmental stages. ____ J. Protozool. 34, 382-392. (L. foxi n. sp., from Gammarus pseudolimnaeus and related spp., and L. missoriensis n. sp., also from G. pseudolimnaeus, have amphipods as hosts).
- CLARK, J. & J.L. BARNARD, 1988. Tonocote introflexidus, a new species of marine amphipod from Argentina (Crustacea: Gammaridea: Amphipoda). ____ Proc. biol. Soc. Wash. 101, 354-365. (T. introflexidus n. sp. comes from the eastern Straits of Magellan. A key to zobrachoid genera is provided).
- COHEN, A.S. & M.R. JOHNSTON, 1987. Speciation in brooding and poorly dispersing lacustrine organisms. ____ Pelaios 2, 426-435. (Not seen).
- COMELY, C.A. & A.D. ANSELL, 1988. Invertebrate associates of the sea urchin, Echinus esculentus L., from the Scottish west coast. ____ Ophelia 28, 111-137. (i.a. Euonyx chelatus and Pariambus typicus inermis).
- CONLAN, K.E., 1988. Phenetic and cladistic methods applied to a small genus (Corophioidea: Ischyroceridae: Microjassa) and a larger outgroup. ____ Crustaceana, Suppl. 13, 143-166. (A methodically most important paper!).
- COSTELLO, M.J. & A.A. MYERS, 1987. Amphipod fauna of the sponges Halichondria panicea and Hymeniacidon perleve in Lough Hyne, Ireland. ____ Mar. Ecol. Progr. Ser. 41, 115-121.
- CROSS, W.E., C.M. MARTIN & D.H. THOMSON, 1987. Effects of experimental releases of oil and dispersed oil on Arctic nearshore macrobenthos. II. Epibenthos. ____ Arctic 40 (Suppl. 1), 201-210.
- CROSS, W.E. & D.H. THOMSON, 1987. Effects of experimental releases of oil and dispersed oils on Arctic near shore macrobenthos - Infauna. ____ Arctic 40 (Suppl. 1), 184-200.
- CULTER, J.H., 1986. Manual for identification of marine invertebrates: A guide to some common estuarine macroinvertebrates of the Big Bend region, Tampa Bay, Florida. ____ Ecol. Res. Ser. U.S. environm. Prot. Agency. EPA/600/4-86/002, 207 pp. (Not seen. Can anybody help me with further data on amphipods treated here?).
- CULVER, D.C., 1987. The role of gradualism and punctuation in cave adaptation. ____ Int. J. Speleol. 16, 17-32.
- CULVER, D.C. & D.W. FONG, 1986. Why all cave animals look alike. ____ Stygologia 2, 208-216.
- CURTIS, L.A., 1987. Vertical distribution of an estuarine snail altered by a parasite. ____ Science 235, 1509-1511. (Trematodes cause the snail Ilyanassa obsoleta to crawl into the high intertidal, where they perish and are scavenged upon by beach-hoppers, the second intermediate host).
- DAUVIN, J-C., 1988. Biologie, dynamique, et production de populations de crustacés amphipodes de la Manche occidentale. 1. Ampelisca tenuicornis Liljeborg. ____ J. exp. mar. Biol. Ecol. 118, 55-84.
- DAUVIN, J-C., 1988. Biologie, dynamique, et production de crustacés amphipodes de la Manche occidentale. 2. Ampelisca brevicornis (Costa). ____ J. exp. mar. Biol. Ecol. 119, 213-233.
- DAUVIN, J-C., 1988. Biologie, dynamique, et production de populations de crustacés amphipodes de la Manche occidentale. 3. Ampelisca typica (Bate). ____ J. exp. mar. Biol. Ecol. 121, 1-22.
- DAUVIN, J-C., & L. CABIOCH, 1988. Nouvelles espèces pour l'inventaire de la faune marine de Roscoff. Crustacés Amphipodes: Siphonocetes striatus Myers & McGrath, et Annélides Polychètes Paraonidae, et nouvelles données sur la répartition du spatangide Echinocardium pennatifidum Norman. ____ Cah. Biol. mar. 29, 215-219.
- DAUVIN, J-C. & A. TOULEMONT, 1988?. Données préliminaires sur les Amphipodes de l'Iroise et de ses abords, leurs affinités biogéographiques. ____ Aspects récents Biol. Crust., Act. Colloq. 8, 217-222.
- DAVIES, B.R., 1984. The zoobenthos of the Touw River floodplain Part 1: The benthos of the Wilderness Lagoon, Touw River and the Serpentine, and the effects of submerged plant cutting. ____ J. limnol. Soc. S. Afr. 10, 62-73. (Melita zeylanica, Corophium triaenonyx and Grandidierella lignorum among the dominant species).
- DE BROYER, C., 1986. Approche d'une revision des Lysianassoidea (Crustacea Amphipoda): definition du groupe Uristidien. ____ Cah. Biol. mar. 27, 328. (Abstract only. Thirty-seven genera comprise the uristid group, the most plesiomorphic Lysianassoidea).
- DE BROYER, C. & M.H. THURSTON, 1987. New Atlantic material and redescription of the type specimens of the giant abyssal amphipod Alicella gigantea (Crustacea). ____ Zool. Scr. 16:335-350.
- DE DECKER, H.P., 1987. Breaching the mouth of the Bot River estuary, South Africa: Impact on its benthic macrofaunal communities. ____ Trans. R. Soc. S. Afr. 46, 231-250.

- DE MARCH, B.G.E., 1988. Acute toxicity of binary mixtures of five cations (Cu^{++} , Cd^{++} , Zn^{++} , Mg^{++} and K^+) to the freshwater amphipod Gammarus lacustris (Sars): alternative descriptive models. ____ Can. J. Fish. aq. Sci. 45, 625-633.
- DESROCHES, M. & P. BRUNEL, 1986. Rhachotropis oculata, Crustacé Amphipode prédateur et migrateur suprabenthique saisonnier dans l'estuaire du St-Laurent. ____ Annls ACFAS 54, 466. (Abstract only).
- DEWITT, T.H., G.R. DITSWORTH & R.C. SWARTZ, 1988. Effects of natural sediment features on survival of the phoxocephalid amphipod, Rhepoxynius abronius. ____ Mar. Environm. Res. 25, 99-124.
- DEXTER, D.M., 1988. Sandy beach fauna of Mediterranean and Red Sea coastlines of Israel and the Sinai peninsula. ____ Israel J. Zool. 34 (1986/87), 125-138. (Amph. - Red Sea coast only - pp. 132-133).
- DEXTER, D.H., 1988. The sandy beach fauna of Portugal. ____ Arq. Mus. Bocage, N.S. 1, 101-110.
- DIEBEL, C.E., 1988. Observations on the anatomy and behavior of Phronima sedentaria (Forskål) (Amphipoda: Hyperiidea). ____ J. crust. Biol. 8, 79-90.
- DITTRICH, B., 1987. Postembryonic development of the parasitic amphipod Hyperia galba ____ Helgol. Meeresunters. 41, 217-232.
- DIVIACCO, G. & W. VADER, 1988. The genus Normanion Bonnier in the Mediterranean Sea, with the description of two new species (Amphipoda, Lysianassoidea). ____ Crustaceana 54, 117-128. (N. chevreuxi n. sp. (= N. quadrimanus s. C & F.) and N. ruffoi n. sp., both from the Gulf of Naples).
- DONNER, K.O., A. LINDSTRÖM & M. LINDSTRÖM, 1987. Seasonal variation in the vertical migration of Pontoporeia affinis (Crustacea, Amphipoda). ____ Ann. zool. fenn. 24, 305-314.
- DOROGOSTAISKAYA, E.V. & H.S. FOREST, 1985. Vitaly Cheslavovich Dorogostaisky: A pioneer investigator of the flora and fauna of Lake Baikal. ____ J. Great Lakes Res. 11, 512-519.
- ELWOOD, R.W. & S.J. NEIL, 1986. Asymmetric contests involving two resources. ____ J. theor. Biol. 120, 237-249.
- ENGLISH, D.S., T.M. ALAN & D.W. BLINN, 1986. Electrophoretic characterization of Hyaella montezuma, an endemic filter-feeding amphipod. ____ J. Hered. 77, 284-285.
- ENGLISH, W.R., 1987. Three inexpensive aquatic invertebrate samplers for the benthos, drift and emergent fauna. ____ Entomol. News 98, 171-179. (Not seen).
- FALCIAI, L. & V. SPADINI, 1986. (The amphipods of the infralittoral plain of the central-northern Tyrrhenian). ____ Atti Soc. tosc. Sci. nat. Mem. B. 92 (1985), 145-163. (In Italian, not seen).
- FALK-PETERSEN, S., J.R. SARGENT & K.S. TANDE, 1987. Lipid composition of zooplankton in relation to the subarctic food web. ____ Polar Biol. 8, 115-120. (i.a. Parathemisto abyssorum).
- FISHELSON, L. & T. HARAN, 1988. Epifauna of algae on a rocky platform near Mikhmoret (Mediterranean Sea, Israel): composition and dynamics. ____ Israel J. Zool. 34 (1986/87), 105-123. (Amph. pp. 122-123).
- FOSTER, B.A., J.M. CARGILL & J.C. MONTGOMERY, 1987. Planktivory in Pagothenia borchgrevinki (Pisces: Nototheniidae) in McMurdo Sound, Antarctica. ____ Polar Biol. 8, 49-54 (Orchomene plebs and Epimeriella macronyx important in diet).
- FRANCE, R.L., 1987. Differences in H- ion sensitivity among Hyaella azteca populations: An illative? hypothesis invoking natural selection. ____ Ann. Soc. R. zool. belg. 117, Suppl. 1. 129-137.
- FRANCE, R.L., 1987. Aggregation in littoral amphipod populations: tranformation controversies revisited. ____ Can. J. Fish. aq. Sci. 44, 1510-1515.
- FRANCE, R.L. & B.D. LAZERTE, 1987. Empirical hypothesis to explain the restricted distribution of Hyaella azteca (Amphipoda) in anthropogenically acidified lakes. ____ Can. J. Fish. aq. Sci. 44, 1112-1121.
- FRANCE, R.L. & P.M. STOKES, 1987. Influence of manganese, calcium, and aluminum on hydrogen ion toxicity to the amphipod Hyaella azteca. ____ Can. J. Zool. 65, 3071-3078.
- FRICKE, H. & J. OEHLenschläger, 1988. Fatty acid and sterol composition of the Antarctic amphipod Themisto gaudichaudii Guérin 1828. ____ Comp. Biochem. Physiol. B. 89, 39-42.
- GABLE, M.F. E.A. LAZO-WASEM & A.J. BALDINGER, 1988. A description of the pigmented and non-stygobiontic females of Podobothrus bermudensis Barnard & Clark, 1985 (Crustacea: Amphipoda: Dulichiidae). ____ Proc. biol. Soc. Wash. 101, 146-150.
- GALAT, D.L., M. COLEMAN & R. ROBINSON, 1988. Experimental effects of elevated salinity on three benthic invertebrates in Pyramid Lake, Nevada. ____ Hydrobiologia 158, 133-144. (i.a. Hyaella azteca).
- GEE, J.H.R., 1988. Population dynamics and morphometrics of Gammarus pulex L.: evidence of seasonal food limitation in a freshwater detritivore. ____ Freshw. Biol. 19, 333-344.

- GERACI, S. & D.J. ST. AUBIN, 1987. Effects of parasites on marine mammals. ____ Intern. J. Parasitol. 17, 407-414. (Not seen).
- GINET, R., 1988. Rejet du taxon Niphargus minutus (Gervais, 1835) et suppression de Niphargus moniezi Wrzesniowski, 1890 (Crustacés Amphipodes). ____ Bull. Soc. linn. Lyon 57, 147- 149. (N. minutus is a nomen nudum, N. moniezi a junior synonym of Crangonyx subterraneus).
- GINSBURGER-VOGEL, T., 1986. Un cas d'ectosymbiose chez un Crustacé? ____ Cah. Biol. mar. 27, 275-276. (Microorganisms on Orchestia gammarellus).
- GINSBURGER-VOGEL, T., 1986. Essai d'analyse de l'arrhenogénie et de la thélygénie indépendantes de l'intersexualité chez Orchestia gammarellus. ____ Cah. Biol. mar. 27, 309-310.
- GODFREY, R.B., J.R. HOLSINGER & K.A. CARSON, 1988. A comparison of the morphology of calceoli in the freshwater amphipods Crangonyx richmondensis s. lat. (Crangonyctidae) and Gammarus minus (Gammaridae). ____ Crustaceana, Suppl. 13, 115-121.
- HACKSTEIN, E., M. SCHIMER & H. LEIBISCH, 1986. Untersuchungen zur Populationsdynamik von Gammarus tigrinus Sexton (Crustacea: Amphipoda) in der Weser bei Bremen. ____ Arch. Hydrobiol. 105, 443-458.
- HAGEN, W., 1988. Zur Bedeutung der Lipiden in antarktischen Zooplankton. ____ Ber. Polarforschung 49, 1-129. (Amph. 64- 73, 109-110).
- HALCROW, K., 1988. SEM studies of the diversity of non-sensory surface microstructure in Amphipoda (Poster presentation). ____ Crustaceana, Suppl. 13, 279.
- HALCROW, K., 1988. TEM investigation of the organisation of the pore canal system of Hyale nilssoni ____ Crustaceana, Suppl. 13, 278. (Abstract only).
- HALCROW, K., 1988. Absence of epicuticle from the repair cuticle produced by four malacostracan crustaceans. ____ J. crust. Biol. 8, 346-354. (l.a. Gammarus oceanicus).
- HARGEBY, A. & R.C. PETERSEN, 1988. Effects of low pH and humus on the survivorship, growth and feeding of Gammarus pulex (L.) (Amphipoda). ____ Freshw. Biol. 19, 235-248.
- HARRIS, G.J. & E. MORGAN, 1986. Seasonal and semi-lunar modulation of the endogenous swimming rhythm in the estuarine amphipod Corophium volutator (Pallas). ____ Mar. Behav. Physiol. 12, 303-314.
- HASEGAWA, M., Y. KUROHJI, S. TAKAYANGI, S. SAWADAISMI & M. YAO, 1986. (Collection of fish and Amphipoda from abyssal sea-floor at 30 N - 147 E using traps tied to 10000 m wire of research vessel). ____ Bull. Tokai reg. Fish. Res. Lab. 119, 65-76. (In Japanese, not seen).
- HAUTUS, T. & S. PINKSTER, 1987. Range extension in the period 1985-1986 of the alien amphipods, Gammarus tigrinus, 1939 (sic!), and Crangonyx pseudogracilis Bousfield, 1958, in the Netherlands (Crustacea, Amphipoda). ____ Bull. zool. Mus. A'dam 11, 57-64.
- HAY, M.E., J. E. DUFFY & C.A. FFISTER, 1987. Chemical defence against different marine herbivores: are amphipods insect equivalents? ____ Ecol. 68, 1567-1580. (The suggested answer is 'yes'. A most interesting paper).
- HAY, M.E., P.E. RENAUD & W. FENICAL, 1988. Large mobile versus small sedentary herbivores and their resistance to seaweed chemical defences. ____ Oecologia 75, 246-252.
- HILL, C. & R. ELMGREN, 1987. Vertical distribution in the sediment in the co-occurring amphipods Pontoporeia affinis and P. femorata. ____ Oikos 49, 221-229. (P. femorata burrows significantly deeper).
- HIROKI, M., 1988. Relation between diel vertical migration and locomotor activity of a marine hyperiidean amphipod, Themisto japonica (Bovallius). ____ J. crust. Res. 8, 48-52.
- HOBERG, E.P., 1986. Aspects of ecology and biogeography of Acanthocephala in antarctic seabirds. ____ Ann. Parasitol. hum. comp. 61, 199-214. (Pontogeneiella sp. as intermediate host).
- HOLMES, J.M.C., 1985. Crustacean records from Lough Hyne, Co Cork 3 ____ Bull. Ir. biogeogr. Soc. 8, 19-25. (Not seen. Seventeen amphipod species on pp. 23-24).
- HOLMES, J.M.C., 1987. Crustacean records from Lough Hyne, Co Cork 4. ____ Bull. Ir. biogeogr. Soc. 10, 99-106. (Not seen. Eulimnogammarus obtusatus and Pontocrates arenarius).
- HOLMES, J.M.C. & J.P. O'CONNOR, 1988. A portable light-trap for collecting marine crustaceans. ____ J. mar. biol. Ass. U.K. 68, 235-238.
- HOLSINGER, J.R., 1988. Trogllobites: the evolution of cave-dwelling organisms. ____ Am. Scient. 76, 147-153.
- HOLSINGER, J.R. & D.C. CULVER, 1988. The invertebrate cave fauna of Virginia and a part of eastern Tennessee: zoogeography and ecology. ____ Bromleyana 14, 1-162.
- HOLSINGER, J.R. & D.P. SHAW, 1987. Stygobromus quatsinensis, a new amphipod crustacean (Crangonyctidae) from caves on Vancouver Island, British Columbia, with remarks on zoogeographic relationships. ____ Can. J. Zool. 65, 2202-2209. (S. q. n. sp. belongs to the hubbsi - group of species).

- HULL, S.C., 1987. Macroalgal mats and species abundance; a field experiment. ____ *Estuar. coast. Shelf Sci.* **25**, 519-533(i.a. Corophium volutator).
- HÜPPOP, K., 1985. The role of mutualism in the evolution of cave animals. ____ *NBS Bull.* **47**, 136-146. (Not seen).
- INGRAM, C.L. & R.R. HESSLER, 1987. Population biology of the deep-sea amphipod Eurythenes gryllus: inferences from instar analyses. *Deep-Sea Res.* **34 A**, 1889-1910.
- ISHIMARU, S-I, 1987. Description of two new species of Guerne (Crustacea, Amphipoda, Dexaminidae) from Japan, with tentative revision of subfamily Prophiantinae. ____ *J. nat. Hist.* **21**, 1395-1414. (Guerne minor n. sp. and G. ezoensis n. sp. The genus Haustoriopsis is revived, while Prinassus is considered invalid even at subgenus level).
- JACOBI, C.M., 1987. The invertebrate fauna associated with intertidal beds of the Brown Mussel Perna perna (L.) from Santos, Brazil. ____ *Stud. neotrop. Fauna Environm.* **22**, 57-72.
- JANGOUX, M., 1987. Diseases of Echinodermata, 3. Agents metazoans (Annelida to Pisces). ____ *Dis. aq. Org.* **3**, 59-83.
- JARAMILLO, E., R.A. CROKER & E.B. HATFIELD, 1987. Long-term structure, disturbance, and recolonization of macrofauna in a New Hampshire sand beach. ____ *Can. J. Zool.* **65**, 3024-3031. (Acanthohaustorius milssi and Amphiporeia virginiana among the dominant species).
- JAVED, W., 1987. Elasmopus japonicus Stephensen, 1932, obtained from the northern Arabian Sea. ____ *Biologia (Lahore)* **33**, 33-38. (Not seen).
- JAZDZEWSKI, K. & A. KONOPACKA, 1988. Notes on the gammaridean Amphipoda of the Dniester River basin and eastern Carpathians. ____ *Crustaceana, Suppl.* **13**, 72-89. (Deals with 25 amphipod taxa).
- JAZDZEWSKI, K. & E. PRESLER, 1988. Hyperiid amphipods collected by the Polish antarctic expedition in the Scotia Sea and in the South Shetlands Islands areas. ____ *Crustaceana, Suppl.* **13**, 61-71.
- JAZDZEWSKI, K. & A.-L. ROUX, 1988. Biogeographie de Gammarus roeseli Gervais en Europe, en particulier repartition en France et en Pologne. ____ *Crustaceana, Suppl.* **13**, 272-277.
- JENSEN, F., 1987. Notes on the benthic stream fauna of the Faroe Islands. ____ *Natura jutlandica* **22**, 73-80.
- JOHNSON, M.G., 1988. Production by the amphipod Pontoporeia hoyi in South Bay, Lake Huron. ____ *Can. J. Fish. aq. Sci.* **45**, 617-624.
- JONES, A.R., 1987. Temporal patterns in the macrobenthic communities of the Hawkesbury estuary, New South Wales. ____ *Austr. J. mar. Freshw. Res.* **38**, 607-624. (Amph. p.623).
- JUST, J., 1985. Siphonoecetinae (Crustacea: Amphipoda: Corophiidae) 4: Australoecetes Just, 1983, including Stebbingoecetes n. subgen. ____ *Rec. austr. Mus.* **37**, 325-341. (Deals with A. sellicki, A. (S) australis (type of Stebbingoecetes n. subgenus), and A. (S) jervisi n. sp. from Jervis Bay, NSW, Australia).
- JUST, J., 1987. Siphonoecetinae (Crustacea, Amphipoda, Corophiidae): 5. Concholestes omani, new species from the coast of Oman. ---- *Steenstrupia* **13**, 93-99. (With remarks on C. dentalii).
- JUST, J. 1988. Siphonoecetinae (Crustacea, Amphipoda, Corophiidae): 6. A survey of phylogeny, distribution, and biology. ____ *Crustaceana, Suppl.* **13**, 193-208.
- KAARTVEDT, S., 1986. Diel activity patterns in deep-living cumaceans and amphipods. ____ *Mar. Ecol. Progr. Ser.* **30**, 243-249. (A study from a west Norwegian fjord).
- KARAKIRI, M. & A. NICOLAIDOU, 1987. Population studies on the Amphipoda of Mazoma Lagoon (Greece). ____ *Helgol. Meeresunters.* **41**, 453-464. (Gammarus insensibilis Dexamine spinosa, Microdeutopus gryllotalpa and Corophium insidiosum).
- KARAKIRI, M. & A. NICOLAIDOU, 1988. On a collection of Amphipoda from the northern Sporades, Aegean Sea. ____ *Crustaceana, Suppl.* **13**, 107-114. (70 amphipod spp., with ecological data).
- KARAMAN, G.S., 1985? Contribution to the knowledge of the Amphipoda. 144. Degocheirocratus spani, new genus and species from Adriatic Sea, with remarks to the Cheirocratus complex of genera (Gammaridea). ____ *Glas. Republ. Zavoda Zast. Prirode- Prirodnjackog Muzeja Titograd* **17** (1984), 5-28. (The new taxon was collected off Dubrovnik in the Adriatic Sea. A key to the Cheirocratus complex of genera, diagnoses of the genera Casco, Cheirocratella, Cheirocratus, Degocheirocratus, Incratella and Prosocratus, and lists of species are given).
- KARAMAN, G.S., 1985? Contribution to the knowledge of the Amphipoda 145. Two subterranean species of the family Niphargidae, Niphargus lunaris, n. sp. and N. timavi S. Kar. 1954. ____ *Glas. Republ. Zavoda Zast. Prirode-Prirodnjackog Muzeja Titograd* **17** (1984), 29-46. (N. lunaris stems from a Bosnian cave. N. timavi, often treated as subsp. of N. stygius, from the Timavo river in Istria).
- KARAMAN, G.S., 1986. Discovery of Niphargus delamarei Ruffo, 1954 in Spain, with first description of females (Gammaridea: Niphargidae). (Contribution to the knowledge of the Amphipoda 154) ____ *Poljoprivreda i Sumarstvo* **33**, 29-42.

- KARAMAN, G.S., 1986. First discovery of genus Niphargus Sch. in Iraq, Israel and adjacent region, with description of N. itus, new species (fam. Niphargidae) (Contribution to the knowledge of the Amphipoda 153). ____ Poljoprivreda i Sumarstvo 32, 13-36. (Deals with N. nadarini from Iraq, Israel and Lebanon, and N. itus n. sp. from Israel. Both species, as also N. altaqahizi from Lebanon, belong to the orcinus-group).
- KARAMAN, G.S., 1986. Syrrhoites barnardi, new marine amphipod from the Mediterranean Sea, with remarks to genus Synopia Dana (Gammaridea, Synopiidae) (Contribution to the knowledge of the Amphipoda 155). ____ Studia mar., Kotor 17-18, 159-178. (The genera Syrrhoites and Latacnga are still kept apart, though S. barnardi n. sp. from the Golfo di Napoli is nearly intermediate. The genus Synopia is divided into Synopia s. str. with deeply incised telson and Telsosynopia n. subgen. with entire telson. Type of T. is Synopia variabilis, further species S. rotunda and S. triangula).
- KARAMAN, G.S., 1986. First discovery of genus Arculfia J.L. Barnard in the Mediterranean Sea, with remarks on two other members of family Pandaliscidae (Contribution to the knowledge of the Amphipoda 156). ____ Acta adriatica 27, 51-66. (Deals with Arculfia trago mediterranea n. ssp. from off S. France, Paradaliscella boeckii and Pandaliscoides tenellus).
- KARAMAN, G.S., 1986. Description of Haploginglymus mateusi, new species of subterranean Gammaridea from Iberian peninsula, with remarks to other taxa of this genus (fam. Niphargidae) (Contribution to the knowledge of the Amphipoda 157). ____ Poljoprivreda i Sumarstvo 32, 75-90. (H. mateusi n. sp. is from a cave in the Guadalupe province of Spain).
- KARAMAN, G.S., 1986. Several poorly known or new species of families Synopiidae and Phoxocephalidae from the Mediterranean Sea (Contribution to the knowledge of the Amphipoda 158). ____ Montenegrin Acad. Sci. Glasnik Sect. nat. Sci. 5, 117-166. (Deals with the synopiids Bruzella typica, Illeustroa i. ilergetes, Syrrhoe angulipes, S. affinis, Pseudotiron bouvieri and Syrrhoites pusilla, and the phoxocephalid Metaphoxus gruneri n. sp. from Malta).
- KARAMAN, G.S., 1985. First discovery of genus Phoxocephalus Stebb. 1888 in the Mediterranean Sea, P. aquosus, n. sp. (fam. Phoxocephalidae) (Contribution to the knowledge of the Amphipoda 159). ____ Bull. Mus. Hist. nat. Belgrade 40, 113-129. (The genera Eusyrophoxus, Cephalophoxus and Cephalophoxoides, all of Gurjanova, are again submerged in Phoxocephalus, and a key to all species provided. P. aquosus n. sp. has Malta as type locality, but has also been found in the Bay of Naples).
- KARAMAN, S.G. (sic!), 1987. On some freshwater gammaridean species new or interesting to the fauna of Italy (Contribution to the knowledge of the Amphipoda 160). ____ Biol. Vestnik 35, 29-44. (Deals with Niphargus arbieter (new to Italy), N. steueri (also new to Italy), N. orcinus and Gammarus roeselii (again new to Italy, but possibly introduced)).
- KARAMAN, G.S., 1986. Redescription of subterranean gammaridean species Niphargus longicaudatus (Costa 1851), based on topotypic material (Contribution to the knowledge of the Amphipoda 161). ____ Fragm. balc. Mus. maced. Sci. nat.
- KARAMAN, G.S., 1986. The genus Gammarus Fabr. in Japan (fam. Gammaridae) (Contribution to the knowledge of the Amphipoda 162). ____ Poljoprivreda i Sumarstvo 32, 81-97. (Redescription of G. nipponensis and G. sobaegensis, the latter new to Japan).
- KARAMAN, G.S., 1987. Two new species of genus Harpinia Boeck (fam. Phoxocephalidae) from the Mediterranean Sea. (Contribution to the knowledge of the Amphipoda 163). ____ Acta adriat. 28, 103-119 (H. agna n. sp. and H. zavodniki n. sp., both from the Bay of Naples).
- KARAMAN, G.S., 1987. Two new species of family Gammaridae from Tunisia and Madagascar (Contribution to the knowledge of the Amphipoda 164). ____ Poljoprivreda i Sumarstvo 35, 17-38. (Echinogammarus dactylus n. sp. from springs in Tunisia, Gammarus ledoyeri n. sp. from deep water off Madagascar. The author reiterates his view that no consistent differences exist between the genera Gammarus and Echinogammarus).
- KARAMAN, G.S., 1987. Taxonomical investigation of the genus Harpinia Boeck in the Mediterranean Sea (fam. Phoxocephalidae) (Contribution to the knowledge of the Amphipoda 165). ____ Poljoprivreda i Sumarstvo 33, 13-44. (H. ala n. sp. is described from the Gulf of Naples. A key to all Mediterranean Harpinia is provided, and H. antennaria, H. crenulata, H. dellavallei, H. pectinata and H. truncata redescribed from Mediterranean material).
- KARAMAN, G.S., 1988. The genus Accubogammarus G. Kar. in Yugoslavia with remarks to the genus Typhlogammarus Schäf. (fam. Gammaridae) (Contribution to the knowledge of the Amphipoda 167) ____ Poljoprivreda i Sumarstvo 34, 63-77. (Describes Accubogammarus albor jalzici n. ssp. from a cave near Dubrovnik. New material of Typhlogammarus mrazeki makes clear that the ssp. heteropalpus can not be maintained).
- KARAMAN, G.S. & S. PINKSTER, 1987. Freshwater Gammarus species from Europe, North Africa and adjacent regions of Asia. 3. Gammarus balcanicus - group and related species. ____ Bijdr. Dierk. 57, 207-260. (Deals with G. balcanicus, G. bosniacus, G. anatoliensis, G. pseudanatoliensis n. sp. (Sivas prov. Turkey), G. abscisus, G. accolae, G. dulensis, G. longipedis n. sp. (Konya prov. Turkey, a grammatically strange name), G. ochridensis, G. macedonicus, G. stankokaramani, G. parechiniformis, G. solidus, G. lychnidensis and G. salemaai).
- KHMELEVA, N.N., 1984. (Features of generative growth of crustaceans on the example of model species). ____ Pp. 71-72 in V.N. OLSHVANG (ed.). (The species and its productivity within its distribution range. Part 4. Invertebrates). Urals scient. Centre, Akad. Nauk USSR, Sverdlovsk (In Russian).
- KHMELEVA, N.N., T.V. MIKHAEVICH & A.I. NESTEROVICH, 1984. ____ Pp. 73-74 in V.N. OLSHVANG (ed.). (The species and its productivity within its distribution range. Part 4. Invertebrates). Urals scient. centre, Akad. Nauk USSR, Sverdlovsk (In Russian).
- KHMELEVA, N.N. & Y. F. MUKHIN, 1984. (Survival and silicon content of Gammarus lacustris from waters of varying temperature. ____ Pp. 72-73 in V.N. OLSHVANG (ed.). (The species and its productivity within its distribution range. Part 4. Invertebrates). Urals scient. Centre, Akad. Nauk USSR, Sverdlovsk (In Russian).

- KITCHING, J.A., 1987. The flora and fauna associated with Himanthalia elongata (L.) S.F. Grey in relation to water current and wave action in the Lough Hyne Marine Nature Reserve. ____ Est. coast. Shelf Sci. 25, 663-676.
- KLUMPP, D.W., A.D. MCKINNON & C.N. MUNDY, 1988. Motile cryptofauna of a coral reef: abundance, distribution and trophic potential. ____ Mar. Ecol. Progr. Ser. 45, 95-108.
- KOLAKOWSKA, A., 1987. Lipids of some Antarctic animals of the Admiralty Bay (King George Island, South Shetland Islands). ____ Polish polar Res. 8 (1987), 391-402. (i.a. Paramoera sp. and Orchomene sp.).
- KUSANO, H. & T. KUSANO, 1988. Seasonal shift in allocation of reproductive resources of Jesogammarus spinopalpus (Amphipoda). ____ Oikos 51, 227-232.
- KUSANO, H., T. KUSANO & Y. WATANABE, 1987. Life history and reproduction of Jesogammarus spinopalpus (Anisogammaridae: Amphipoda) inhabiting in a lowland pond. ____ Jpn J. Limnol. 48, 117-126.
- LAGARDE, G., 1987. (Amphipoda Gammaridea of the Golfo Triste coast, Venezuela, and neighbouring areas). ____ Caribb. J. Sci. 23, 260-277. (In Spanish, not seen).
- LAMARCHE, G. & P. BRUNEL, 1987. Cycle de developpement, ecologie et succès d'Hippomedon propinquus (Amphipoda, Gammaridea) dans deux écosystèmes du Golfe du Saint-Laurent. ____ Can. J. Zool. 65, 3116-3132.
- LANDRUM, P.F., 1988. Toxicokinetics of organic xenobiotics in the amphipod, Pontoporeia hoyi: role of physiological and environmental variables. ____ Aquat. Toxicol. (Amst.) 12, 245-272.
- LANTOS, G., 1986. Data in the Amphipoda- and Isopoda-fauna of Toserdo and its environs in the Tisza Valley 1. Amphipoda, Asellota (Crustacea, Peracarida), ____ Tiscia 21, 81-87.
- LEE, K.S., 1986. Systematic study of Amphipoda (Crustacea) in Korea. 6. Gammarus hoonsooi, a new species of freshwater gammarid (Gammaridae) from South Korea. ____ Korean J. Zool. 29, 165-170.
- LEE, K.S., 1986. Systematic study of Amphipoda (Crustacea) in Korea. 5. Descriptions of one hitherto unrecorded species and two known species from Korean waters. ____ Korean J. Zool. 29, 159-164. (Three Caprella spp. Not seen).
- LEGRAND, J.J., E. LEGRAND-HAMELIN & P. JUCHAULT, 1987. Sex determination in Crustacea. ____ Biol. Rev. 62, 439-470.
- LEONARDSSON, K., T. SÖRLIN & H. SAMBERG, 1988. Does Pontoporeia affinis (Amphipoda) optimize age at reproduction in the Gulf of Bothnia? ____ Oikos 52, 328-336.
- LITT, R., 1986. Observation sur le crustacé aquatique Niphargus aquilex S. ____ Rev. verbiét. Hist. nat. 1986 (spring), 1-3. (Not seen).
- LOBEL, P.S. & J. E. RANDALL, 1986. Swarming behavior of the hyperiid amphipod Anchylomera blossevillei. ____ J. Plankton Res. 8, 253-262.
- LØNNE, O.J., 1988. A diver-operated electric suction sampler for sympagic (=under-ice) invertebrates. ____ Polar Res. 6, 135-136.
- LOWRY, J.K. & H. E. STODDART, 1987. A new South American genus and species in the amaryllid group of lysianassoïd Amphipoda. ____ J. nat. Hist. 21, 1303-1309. (Eriks dahl n. gen., n. sp. from Chile).
- MACDONALD, C.R. & J.B. SPRAGUE, 1988. Cadmium in marine invertebrates and arctic cod in the Canadian Arctic. Distribution and ecological implications. ____ Mar. Ecol. Progr. Ser. 47, 17-30. (Several amphipods. Parathemisto libellula shows consistently high cadmium levels).
- MACKIE, G.O., P-R. PUGH & J.E. PURCELL, 1987. Siphonophore biology. ____ Adv. mar. Biol. 24, 97-262. (Associations with amphipods on pp. 205-207).
- MACQUART-MOULIN, C., A. BOURDILLON, F. CUBIZOLES, F. PASSELAIGUE & R. RASOANARIVO, 1987. Un cas type de migration verticale 'retard' chez l' Amphipode Ampelisca typica ____ J. Plankton Res. 9, 785-790. (Not seen).
- MALICKY, H., 1985. Fraszversuche mit einigen Flieswasser-tieren an verschiedenen Fallaub. ____ Jahresh. biol. Stn Lunz, österr. Akad. Wiss. 8 (1984), 59-63. (Not seen, i.a. Gammarus).
- MALMQUIST, B. & P. SJÖSTRÖM, 1987. Stream drift as a consequence of disturbance by invertebrate predators. Field and laboratory experiments. ____ Oecologia 74, 396-403.
- MARMONIER, P., 1985. Spatial distribution and temporal evolution of Gammarus fossarum, Niphargus sp. (Amphipoda) and Proasellus slavus (Isopoda) in the Seebach sediments (Lunz, Austria). ____ Jahresber. biol. Stn. Lunz, österr. Akad. Wiss. 8 (1984), 40-54. (Not seen).
- MARMONIER, P. & M.J. DOLE, 1986. Les amphipodes des sédiments d'un bras court-circuité du Rhone. Logique de repartition et réaction aux crues. ____ Sciences Eau 5, 461-486. (Not seen).
- MARQUES, J-C. & D. BELLAN-SANTINI, 1987. Crustacés Amphipodes des côtes du Portugal: faune de l'estuaire du Mira (Alenteja. Côte sud-ouest). ____ Cah. Biol. mar. 28, 465-480. (Thirty-five spp., of which Dexamine thea is new to Portugal).

- MATSUMIYA, Y., T. KOKUCHI & H. SUDO, 1986. (A consideration on sampling efficiency and quantitative sampling for benthos, especially gammaridean amphipods). ____ Bull. Fish. Nagasaki Univ. 60, 41-51. (In Japanese. Not seen).
- McCAHON, C.P.A., A.F. BROWN & D. PASCOE, 1988. The effect of the acanthocephalan *Pomphorhynchus laevis* (Mueller 1776) on the acute toxicity of cadmium to its intermediate host, the amphipod *Gammarus pulex* (L.). ____ Arch. environm. Contam. Toxicol. 17, 239-244.
- McMAHON, C.P. & D. PASCOE, 1988. Cadmium toxicity to the freshwater amphipod *Gammarus pulex* (L.) during the molt cycle. ____ Freshw. Biol. 19, 197-204.
- MEURS, H.-G. & G.P. ZAUKE, 1988. Regionale und zeitliche Aspekte der Besiedlung des Elbe-, Weser- und Emsästuars mit euryhalinen Gammariden (Crustacea: Amphipoden). ____ Arch. Hydrobiol. 113, 213-230.
- MEUSY, J.-J., G. MARTIN, D. SOYEZ, J.E. VAN DEYNEN & J.M. GALLO, 1987. Immunochemical and immunocytochemical studies of the crustacean vitellogenesis - inhibiting hormone (VIH). ____ Gen. comp. Endocrinol. 67, 333-341. (Not seen).
- MEYERING, M.P.D., 1987. Die *Gammarus*-Fauna im Pfuhlgraben-Bachsystem bei Wehrda- ein längerfristiger Vergleich. ____ Beitr. Naturk. Osthessen 23, 71-79.
- MEYRAN, J.-C., J. FRANÇOIS & F. GRAF, 1988. Analysis of the protein content of the calcareous deposits in the posterior caeca of the crustacean *Orchestia cavimana*. ____ Comp. Biochem. Physiol. B 89, 213-220.
- MEYRAN, J.-C., F. GRAF & J. FOURNIE, 1987. Carbonic anhydrase activity in a calcium - mobilizing epithelium of the crustacean *Orchestia cavimana* during molting. ____ Histochemistry 87, 419-430.
- MEYRAN, J.-C., F. GRAF & G. NIÇAISE, 1986. Pulse discharge of calcium through a demineralizing epithelium in the crustacean *Orchestia*: ultrastructural cytochemistry and x-ray microanalysis. ____ Tissue Cell 18, 267-283.
- MINCHIN, D. & J.M.C. HOLMES, 1987. *Phronima sedentaria* (Forsk.) (Crustacea: Amphipoda) in Irish waters. ____ Irish Nat. JI 22, 202-203.
- MOORE, P.G., 1988. New and little-known Amphipoda (Crustacea) from Tasmania and Western Australia. ____ J. nat. Hist. 22, 149-174. (Describes *Amphilochus rupertii* n. sp., *Ceradocopsis hamondi* n. sp. and *Gammaropsis insignis* n. sp., all from Tasmanian algal habitats. *Photis dolichomata*, and *Parawaldeckia yamba* are redescribed. *Yulumara tricuspidis* n. sp. was collected from seagrasses in Western Australia).
- MOORE, P.G., 1988. Taxonomic observations on the genera *Xenocheira* Haswell and *Erichthonius* Milne Edwards (Crustacea: Amphipoda) from Australian coastal waters. ____ J. nat. Hist. 22, 705-732. (*Xenocheira fasciata* is redescribed from Tasmanian specimens. Material from W. Australia is tentatively identified as *X. seurati*, while Pirlot's male '*X. fasciata*' from the Aru Islands represents a third species, *X. pirloti* n. sp. (erroneously 'nom. nov.' in paper). In the genus *Erichthonius* 2 new spp. are described, *E. tacitus* n. sp. from Tasmania and *E. coxacanthus* n. sp. from W. Australia. Also *E. pugnax* is redescribed and illustrated.
- MOORE, P.G. & P.J. SCHEMBRI, 1986. Notes concerning the semi-terrestrial and freshwater amphipods (Crustacea: Peracarida) of the Maltese Islands. ____ Animalia, Catania 13, 65-75. (Five talitrids and 4 gammarids, with a nice survey of the habitats on the islands).
- MORINO, H., 1986. A new species of the subgenus *Annanogammarus* (Amphipoda: Anisogammaridae) from Lake Suwa, Japan. ____ Publ. Itako hydrobiol. Stn 3, 1-11. (*Jesogammarus* (A). *suwaensis* n. sp.)
- MORINO, H. & H. MIYAMOTO, 1988. Redefinition of *Talorchestia* (Amphipoda: Talitridae) with description of a new species from the tropical West Pacific. ____ J. crust. Res. 8, 91-98. (The genus *Talorchestia* is restricted to *T. gracilis* (type), *T. spinipalma*, *T. martensii*, and *T. palawanensis* n. sp. from the Philippine Islands. *T. spinipalma* is redescribed).
- MORRIS, R.J., A.P.M. LOCKWOOD, D. DYBALL & S.R.L. BOLT, 1987. Changes in the fatty acid composition of the gill phospholipids in *Gammarus duebeni* during molt: Evidence for reduced permeability of the gill membrane. ____ Comp. Biochem. Physiol. B 88, 257-260. (Not seen)
- MORRISEY, D.J., 1988. Differences in effects of grazing by deposit-feeders *Hydrobia ulvae* (Pennant) (Gastropoda: Prosobranchia) and *Corophium arenarium* Crawford (Amphipoda) on sediment microalgal populations. 1. Qualitative differences. ____ J. exp. mar. Biol. Ecol. 118, 33-42.
- MORRISEY, D.J., 1988. Differences in effects of grazing by deposit-feeders *Hydrobia ulvae* (Pennant) (Gastropoda: Prosobranchia) and *Corophium arenarium* Crawford (Amphipoda) on sediment microalgal populations. 2. Quantitative differences. ____ J. exp. mar. Biol. Ecol. 118, 43-53.
- MUSKO, I.B., 1988. Ultrastructure of the midgut gland of *Gammarus roeselii* Gervais (Amphipoda, Gammaridea). ____ Crustaceana 54, 207-217.
- MYERS, A.A., 1988. A cladistic and biogeographic analysis of the Aorinae subfam. nov. ____ Crustaceana, Suppl. 13, 167-193. (In this important paper, the new subfamily Aorinae has the following composition: *Aora* (A. *typica* + 13), *Aorella* (A. *multiplax*), *Aoroides* (A. *columbiae* + 6), *Autonoe* (A. *longipes* + 11), *Bemlos* (B. *macromanus* + 34), *Columbaora* (C. *cyclocoxa*), *Globosolembos* (G. *smithi* + 7), *Lemboides* (L. *afer* + 1), *Meridiolembos* n. gen. (*Lembos hippocrenes* further spp. L. *acherontis*, L. *pertinax*), *Microdeutopus* (M. *gryllotalpa* + 11), *Paramicrodeutopus* n. gen. (*Microdeutopus schmitti*, further spp. M. *hancocki*, M. *myersi* & M. *trichopus*), *Plesiolembos* n. gen. (*Lembos rectangulatus* (with L. *habanensis* as synonym)), further sp. L. *ovalipes*), *Protolembos* n. gen. (*Lembos chiltoni*, further spp. L. *kidoli* and L. *philacanthus*), and *Tethylembos* n. gen. (*Lembos vigueri*), *Lemboides caecus* and *Microdeutopus thumellinus* are removed to the Neomegamphipodidae. The new genus *Australomicrodeutopus* n. gen. (*Microdeutopus haswelli*, further sp. M. *apopo*) is an aorid, but not in the subfamily Aorinae).

- NAGATA, K., 1986. Amphipod crustaceans found near Syowa Station, Antarctica (1). ____ *Mems. natn. Inst. polar Res., spec. Issue* **40**, 249-258. (Deals with Orchomene rossii, O. plebs, Uristes murrayi and Waldeckia obesa)
- NAGATA, K., 1986. Amphipod crustaceans from surface waters of the southern Ocean during 1983-84 summer. ____ *Mems. natn. Inst. pol. Res., spec. Issue* **40**, 259-276. (Deals with Eusirus microps, Paramoera walkeri, Vibilia stebbingi and Hyperiella antarctica).
- NESIS, K.N., 1986. (An amphipod which pretends to be the eggs of its prey). *Priroda, Moskva* 1986 (4), 112. (In Russian. Tells the story of Paracyphocaris praedator as found by Bowman & Wassmer).
- NOTENBOOM, J., 1987. Species of the genus Pseudoniphargus Chevreux, 1901 (Amphipoda) from the Betic Cordillera of southern Spain. ____ *Bijdr. Dierk.* **57**, 87-150. (This important study describes and illustrates P. branchiatus, P. nevadensis n. sp. (Granada), P. granadensis n. sp. (Granada), P. grandis n. sp. (Malaga), P. affinis n. sp. (Granada), P. stocki n. sp. (Malaga), P. vomeratus n. sp. (Jaen), P. illustris n. sp. (Jaen), P. margalefi n. sp. (Alicante), P. cazorlae n. sp. (Jaen), P. latipes n. sp. (Jaen), P. gracilis n. sp. (Almeria), P. sorbasensis n. sp. (Almeria), P. sp. 2, P. fragilis n. sp. (Malaga), P. gibraltarius n. sp. (Cadiz) and P. ssp 3, 4 and 5).
- NOTENBOOM, J., 1987. Lusitanian species of the amphipod Pseudoniphargus, 1901, with a key to all known Iberian species. ____ *Bijdr. Dierk.* **57**, 191-206. (Describes and illustrates P. calliaicus n. sp. (La Coruna, Spain), P. mateusorum and P. brevipedunculatus).
- NOTENBOOM, J., 1988. Parapseudoniphargus baetis, new genus, new species, a stygobiont amphipod crustacean from the Guadalquivir river basin (Southern Spain), with phylogenetic implications. ____ *J. crust. Biol.* **8**, 110-121. (With a discussion of the phylogenetic position of Pseudoniphargus, a close relative of the new genus).
- NOTENBOOM, J., 1988. Biogeographical observations on the genera of Iberian stygobiont Amphipoda. ____ *Crustaceana, Suppl.* **13**, 122-133.
- ORMEROD, S.J. & S.J. TYLER, 1988. The diet of Green Sandpipers Tringa ochropus in contrasting areas of their winter range. ____ *Bird Study* **35**, 25-30. (Gammarus locally important).
- ORTIZ, M. & T. VELEDO, 1985. (A new species of amphipod of the genus Garosyrhoe (Synopiidae, Gammaridea) from Cuban waters). ____ *Rev. Invest. mar.* **6** (1), 14-18. (In Spanish; G. luquei n. sp.).
- OSHEL, P.E. & D.H. STEELE, 1988. Comparative morphology of amphipod setae, and a proposed classification of setal types. ____ *Crustaceana Suppl.* **13**, 90-99.
- OSHEL, P.E. & D.H. STEELE, 1988. SEM morphology of the foreguts of gammaridean amphipods compared to Anaspides tasmaniae (Anaspidacea; Anaspididae), Gnathophausia ingens (Mysidacea: Lophogastridae) and Idotea balthica (Isopoda: Idoteidae). ____ *Crustaceana. Suppl.* **13**, 209-219.
- OSHEL, P.E., V.J. STEELE & D.H. STEELE, 1988. Comparative SEM morphology of amphipod microtrich sensilla. ____ *Crustaceana, Suppl.* **13**, 100-106.
- PEARSON, R.G. & N.V. JONES, 1987. Short-term movements of chalk stream invertebrates. ____ *Freshw. Biol.* **18**, 559-568. (Many data on Gammarus pulex).
- PECK, S.B. 1988. A review of the cave fauna of Canada, and the composition and ecology of the invertebrate fauna of caves and mines in Ontario. ____ *Can. J. Zool.* **66**, 1197-1213.
- PIEPENBURG, D., 1988. Zur Zusammensetzung der Bodenfauna in der westlichen Fram-Strasse. ____ *Ber. Polarforsch.* **52**, 1-118. (Amph. p. 43).
- PINKSTER, S., 1988. Problems in the taxonomy of the freshwater gammarids with special emphasis on the genus Echinogammarus in Italy. ____ *Crustaceana, Suppl.* **13**, 245-255. (A cautionary tale, in which the author convincingly demonstrates seasonal differences in morphology in freshwater amphipods. As one result, E. bolo and E. roco turn out to be junior synonyms of E. tibaldii. Pinkster closes his paper with an urgent plea 'not to describe new species on the basis of occasional samples', a procedure of very common occurrence hitherto).
- PLATVOET, D., 1988. Side-line organs in amphipods (poster presentation). ____ *Crustaceana, Suppl.* **13**, 281.
- PLESHA, P.D., J.E. STEIN, M.H. SCHIEWE, B.B. McCAIN & U. VARANASI, 1988. Toxicity of marine sediments supplemented with mixtures of selected chlorinated and aromatic hydrocarbons to the infaunal amphipod Rhepoxynius abronius. ____ *Mar. environm. Res.* **25**, 85-97.
- PYATAKOVA, G.M., 1987. (Study of the effect of hydraulic fluids on aquatic organisms on the Caspian Sea). ____ *Izv. Akad. Nauk Az. SSR. Ser. Biol., Nauk* 1986 (5), 41-45. (In Russian, not seen. Deals i.a. with Pontogammarus maeoticus).
- RAGA, J.A., 1988. On some morphological variations of Syncyamus aequus Lincoln & Hurley, 1981 (Amphipoda, Cyamidae) from the Mediterranean Sea. ____ *Crustaceana* **54**, 149-152. (Material from Stenella coerulescens).
- READ, P., 1987. The intertidal benthos and sediments of particulate shores in the Firth of Forth, Scotland, with particular reference to waste water discharges. ____ *Proc. R. Soc. Edinb.* **93** B, 401-413.
- RODRIGUEZ, A.M. & J. C. DAUVIN, 1987. Crustacés peracarides de la 'Ria de Alvor' (Côte du Sud du Portugal). ____ *Cah. Biol. mar.* **28**, 207-223. (Amph. listed pp. 210-211 and 218-219).

- RONN, C., E. BONSDORF & W.G. NELSON, 1988. Predation as a mechanism of interference within infauna in shallow brackish water soft bottoms: experiments with an infauna predator, Nereis diversicolor O.F. Müller. ____ J. exp. mar. Biol. Ecol. 116, 143-158.
- RONNEBERGER, D., 1987. Zur Frage der Besiedlungsabhängigkeit von Untergrundstruktur und Fließgeschwindigkeit in Grundwasser-biotopen Thüringens (DDR). ____ Rocznik Muz. Okregawego w Czestochowie 3, 50-58.
- ROSILLON, D., 1987. About the separation of benthos from stream samples. ____ Arch. Hydrobiol. 110, 469-476.
- RUFFO, S., 1987. (Studies on amphipod crustaceans 103. The Mediterranean species of Lysianassa H. Milne-Edwards, 1830 and description of Pardia, new genus (Crustacea, Amphipoda, Lysianassidae). ____ Monit. zool. ital., Suppl. 32, 31-58. (In Italian. Pardia n. gen. is erected for Callisoma punctatum; this species is for the first time reported from outside the Mediterranean, viz. from Senegal. Lysianassa caesarea n. sp. is described from the Mediterranean coast of Israel. Descriptions and illustrations are also provided of L. longicornis apparently a Med. endemic), L. pilicornis and L. insperata (new to the Mediterranean). A key to Mediterranean Pardia and Lysianassa spp. concludes this useful paper).
- RUSSO, A.R., 1987. Role of habitat complexity in mediating predation by the gray damselfish Abudefduf sordidus on epiphytal amphipods. ____ Mar. Ecol. Progr. Ser. 36, 101-105.
- RYER, C.H., 1988. Pipefish foraging: effects of fish size, prey size and altered habitat complexity. ____ Mar. Ecol. Progr. Ser. 48, 37-45. (Syngnathus fuscus, a predator of amphipods in Zostera meadows).
- SABATER, F., 1988. (Some interstitial species of the crustacean communities of the Ter and Ebro river mouths (northeastern Spain). ____ Misc. Zool. 10 (1986), 113-120. (In Spanish, not seen; i.a. two amphipod spp.)
- SAINTE-MARIE, B. & B.T. HARGRAVE, 1987. Estimation of scavenger abundance and distance of attraction to bait. ____ Mar. Biol. 94, 431-443.
- SALEMAA, H., 1988. Chromosomes in Gammaridea. ____ Crustaceana, Suppl. 13, 281-282. (Abstract only. Data from Baltic Sea and Lake Ohrid).
- SALMAN, S.D., 1985. Stenothoe irakiensis, a new species of stenothoid amphipod from the Arabian Gulf. ____ Crustaceana 49, 244-250.
- SCAPINI, F., 1986. Inheritance of solar direction finding in sandhoppers, 4. Variation in the accuracy of orientation with age. ____ Monit. zool. ital. 20, 53-61.
- SCAPINI, F. & M. BUIATTI, 1985. Inheritance of solar direction finding in sandhoppers. 3. Progeny tests. ____ J. comp. Physiol. 157 A, 433-440.
- SCAPINI, F., A. ERCOLINI & R. BOCCACI, 1988. Laboratory experiments on geotaxis, phototaxis, and anemotaxis in two species of littoral amphipods. ____ Monit. zool. ital. 22, 89-103. (Talitrus saltator and Orchestia mediterranea).
- SCHEEPMAKER, M., 1987. Morphological and genetic differentiation of Gammarus stupendus Pinkster, 1983 in the Massif de la Sainte Baume, France. ____ Bijdr. Dierk. 57, 1-18.
- SCHODEL, H., 1986. Epizoische Einzeller auf Flohkrebse. 3. Besiedler der Coxalplatten und der Mundwerkzeuge. ____ Mikrokosmos 75, 5-11.
- SCHRAMM, H.L., K. J. JIRKA & M.V. HOYER, 1987. Epiphytic macroinvertebrates on dominant macrophytes in two central Florida lakes. ____ J. Freshw. Ecol. 4, 151-162.
- SCHUCHARDT, B., U. HÄSLOOP & M. SCHIRMER, 1987. (On the distribution of Gammarus tigrinus Sexton in the lower Weser (West Germany)). ____ Drosera 87, 129-134. (In German, not seen).
- SCONFIETTI, R., 1982. (Record of Elasmopus pecteniscus (Bate) (Crustacea, Amphipoda) in the Lagoon of Venice (Italy)). ____ Boll. Mus. Civ. Stor. Nat. Venezia 33(1981), 91-92. (In Italian, not seen).
- SEBASTIEN, R.J., D.M. ROSENBERG & A. P. WIENS, 1988. A method for subsampling unsorted benthic macroinvertebrates by weight. ____ Hydrobiologia 157, 69-75. (Not seen).
- SELDEN, P.A., 1986. A new identity for the Silurian arthropod Necrogammarus. ____ Palaeontology 29, 629-631. (Not seen).
- SHANKS, A.L. & W.G. WRIGHT, 1987. Internal-wave-mediated shoreward transport of cyprids, megalopae and gammarids, and correlated longshore differences in the settling rate of intertidal barnacles. ____ J. exp. mar. Biol. Ecol. 114, 1-14.
- SHAW, D.P., 1988. Redescription of Bouvierella carcinophila (Chevreux, 1889) (Eusiroidea, Calliopidae) from northern British Columbia and its proposed synonymy with Leptamphopus paripes Stephensen, 1931. ____ Can. J. Zool. 66, 939-943.
- SHILLAKER, R.O. & P.G. MOORE, 1987. Tube-emergence behaviour in the amphipods Lembos websteri Bate and Corophium bonnellii Milne Edwards. ____ J. exp. mar. Biol. Ecol. 111, 231-241.
- SKALSKI, A.W., 1988. Redescription of Synurella ambulans ssp. tenebrarum (Wrzesniowski, 1888), status n., with notes on its relatives. ____ Crustaceana, Suppl. 13, 220-237. (A redescription of Boruta tenebrarum, here considered a valid subspecies of Synurella ambulans. Also S. intermedia montenigrina is transferred to S. ambulans as a valid subspecies).

- SLATTERY, P.N. & J.S. OLIVER, 1986. Scavenging and other feeding habits of lysianassid amphipods (*Orchomene* spp.) from McMurdo Sound, Antarctica. ____ *Polar Biol.* **6**, 171-177.
- SMITH, D.G., 1987. The genus *Synurella* in New England (Amphipoda, Crangonyctidae). ____ *Crustaceana* **53**, 304-306.
- SNOW, N.B., W.E. CROSS, R.H. GREEN & J.N. BUNCH, 1987. The biological setting of the BIOS site at Cape Hatt, Northwest Territories (Canada), including the sampling design, methodology, and baseline results for macrobenthos. ____ *Arctic* **40**, Suppl. 1, 80-99.
- SPICER, J.I., P.G. MOORE & A.C. TAYLOR, 1987. The physiological ecology of land invasion by the Talitridae (Crustacea: Amphipoda). ____ *Proc. R. Soc. Lond. B* **232**, 95-124.
- SPICER, J.I. & A.C. TAYLOR, 1987. Carbon dioxide transport and acid-base regulation in the blood of the beach-hopper *Orchestia gammarellus* (Pallas) (Crustacea: Amphipoda). ____ *Ophelia* **28**, 49-61.
- STATZNER, B., 1987. Growth and Reynolds number of lotic macroinvertebrates: a problem for adaptation of shape to drag. ____ *Oikos* **51**, 84-87.
- STAUDE, C.P., 1987. Suborder Gammaridea. ____ Pp. 346-391 in E.N. Kozloff (ed.), *Marine Invertebrates of the Pacific Northwest*. Univ. Washington Press, Seattle & London, 509 pp. (A fine set of keys, and an annotated checklist).
- STEELE, D.H., 1988. What is the amphipod life style? ____ *Crustaceana*, Suppl. **13**, 134-142. ('Amphipods are clinging aquatic animals whose primary locomotion is by swimming'. A very important paper!)
- STOCK, J.H., 1987. Stygofauna of the Canary Islands, 5. A hypogean population of *Parhyale* (Amphipoda) in the Jameodel Agua lava tunnel (Lanzarote), a supposed case of recent evolution. ____ *Stygologia* **3**, 167-184. (The Lanzarotan material belongs to the *P. hawaiiensis* complex, but is here described as a new species, *P. multispinosa* n. sp. Material of *P. hawaiiensis* from the West Indies, Hawaii and La Palma (intertidal, first record for Canary Islands) is also described. The type material of *P. injacka* K.H. Bnd also belongs to *P. hawaiiensis*).
- STOCK, J.H., 1988. The amphi-atlantic and insular distribution of *Pseudoniphargus*. ____ *Crustaceana*, Suppl. **13**, 278. (Abstract only).
- STOCK, J.H., 1988. Stygofauna of the Canary Islands. 6. A new *Rhipidogammarus* (Crustacea: Amphipoda) from Tenerife: first record of the genus outside the Mediterranean region and its biogeographic implications. ____ *Hydrobiologia* **169**, 279- 292. (*R. nivariae* n. sp. from Tenerife).
- STOCK, J.H., 1988. Stygofauna of the Canary Islands. 8. Amphipoda (Crustacea) from inland groundwaters of Fuerteventura. ____ *Bull. zool. Mus. A'dam* **11**, 105-113. (*Bogidiella* (*Stygogidiella*) *purpuriae* n. sp., and *Pygocrangonyx repens*).
- STOCK, J.H., 1988. Two new stygobiont Amphipoda (Crustacea) from Polynesia. ____ *Stygologia* **4**, 79-100. (*Fiha schminkei* n. gen., n. sp., (Melitidae, *Psammoniphargus* - group,) is described from riverbank-interstitial in Fiji, and *Josephosella hamata* n. sp. from a marine cave on the Tonga islands. The preoccupied genus name *Quadrus* in the same species group is replaced by *Sriha* n. nom. (Melitidae)).
- STOCK, J.H. & T.M. ILIFFE, 1987. The status of *Bogidiella balearica* Dancau, 1973, a stygobiont amphipod from Madeira ____ *Endins* **13**, 39-46. ('A good species', clearly different from *B. (Metagidiella) chappuisi*, and belonging to the subgenus *Bogidiella* s. str.).
- STOCK, J.H. & J.L. MARTIN, 1988. A new cavehopper (Amphipoda: Talitridae) from lava tubes in La Palma, Canary Islands. ____ *J. nat. Hist.* **22**, 1121-1133. (*Palmorchestia hypogaea* n. gen., n. sp. from lava caves on La Palma).
- STOCK, J.H. & J. NOTENBOOM, 1988. Five new bogidiellid Amphipoda from Spain - the first freshwater records in the Iberian peninsula. ____ *Hydrobiologia* **164**, 75-95. (*Bogidiella* (*B.*) *hispanica* n. sp. (prov. Cuenca), *B. (B.) glabra* n. sp. (prov. Cuenca), *B. (B.) convexa* n. sp. (prov. Madrid), *B. (Medigidiella) uncinata* n. sp. (prov. Valencia), and *B. (M.) antennata* n. sp. (prov. Valencia). A key to Iberian *Bogidiella* is provided).
- STOCK, J.H. & B.L.M. RONDE-BROEKHUIZEN, 1987. Stygofauna of the Canary Islands, 3. The genus *Bogidiella* (Crustacea, Amphipoda). ____ *Rev. Zool. afr.* **101**, 439-461. (Deals with *B. (Xystriogidiella) spathulata* n. sp., *B. (Stygogidiella) uniramosa* n. sp. and *Bogidiella* sp.).
- STOCK, J.H. & E. SANCHEZ, 1987. Stygofauna of the Canary Islands 7. *Psammogammarus initialis* n. sp. a new mediolittoral interstitial amphipod crustacean from Tenerife. ____ *Stygologia* **3**, 264-277. (In a discussion of generic taxonomy, *Roropisa* is re-united with *Victoriopisa*, and *Confodiopisa*, *Flagitopisa* and *Impertiopisa* all with *Psammogammarus*. A key to all *Psammogammarus* (s.l.) species is provided).
- STORCH, V. & P. BURKHARDT, 1986. (The response of the midgut glands of *Orchestia cavimana* to different foods). ____ *Carolinea* **44**, 149-152. (In German, not seen).
- SUDO, H., M. AZUMA & M. AZETA, 1987. Diel changes in predator-prey relationships between red sea bream and gammaridean amphipods in Shijiki Bay, Japan. ____ *Bull. Jpn. Soc. scient. Fish.* **53**, 1567-1575. (Not seen).
- SVESHNIKOV, V.A. & G.M. VINOGRADOV, 1987. (Life forms of Amphipoda Hyperidea) ____ *Dokl. Akad. Nauk SSSR* **293**, 1011-1015. (In Russian).
- TARAMELLI, E. & L. VENANZANGELL, 1988. Amphipoda of Torvaldaliga (Civitavecchia - Roma). ____ *Crustaceana*, Suppl. **13**, 283. (Abstract only).
- TEDENGREN, M., M. ARNÉR & N. KAUTSKY, 1988. Ecophysiology and stress response of marine and brackish water *Gammarus* species (Crustacea, Amphipoda) to changes in salinity and exposure to cadmium and diesel-oil. ____ *Mar. Ecol. Progr. Ser.* **47**, 107-116.

- THOMAS, J.D. & J.L. BARNARD, 1988. Elasmopus balkomanus, a new species from the Florida Keys (Crustacea, Amphipoda). ____ Proc. biol. Soc. Wash. 101, 838-842.
- TIMMS, B.V., U.T. HAMMER & J.W. SHEARD, 1986. A study of benthic communities in some saline lakes in Saskatchewan and Alberta, Canada. ____ Int. Rev. ges. Hydrobiol. 71, 759-777. (i.a. Hyaella azteca).
- TJØNNELAND, A., S. ØKLAND & A. NYLUND, 1987. Evolutionary aspects of the arthropod heart. ____ Zool. Scripta 16, 167-176.
- TULLY, O. & P. O'CEIDIGH, 1987. Investigations of the plankton of the west coast of Ireland. 8. The neustonic phase and vertical migratory behavior of benthic Peracarida in Galway Bay. ____ Prov. R. Ir. Acad. B 87, 43-64.
- TZVETKOVA, N.L., 1987. New species of lysianassid (Amphipoda, Gammaridea, Lysianassidae) from the upper sublittoral zone of Simushir Island (the Kurile Islands). ____ Zool. Zh. 66, 1739-1744. (Ocosingo kussakini n. sp.).
- UGOLINI, A. & F. SCAPINI, 1988. Orientation of the sandhopper Talitrus saltator (Amphipoda, Talitridae) living on dynamic sandy shores. ____ J. comp. Physiol. A 162, 453-462.
- ULIAN, G. & E.G. MENDES, 1987. Preferences of a terrestrial amphipod, Talitrus pacificus Hurley, 1955, towards some environmental factors. ____ Rev. Brasil. Biol. 47, 247-256.
- UNDERWOOD, A.J. & P.H. VERSTEGEN, 1988. Experiments on the association between the intertidal amphipod Hyaella media Dana and the limpet Cellana tramoserica (Sowerby). ____ J. exp. mar. Biol. Ecol. 119, 83-98.
- VALTER, E.D., 1987. (Marinogammarus obtusatus (Amphipoda) as a new intermediate host of the nematode Pseudoterranova decipiens) ____ Biol. Nauki (Mosc.) 1987 (6), 28-32. (In Russian, not seen).
- VAN SENS, P., 1988. Reproduction of the sandhopper, Talorchestia capensis (Dana) (Amphipoda, Talitridae). ____ Crustaceana 55, 93-103.
- VAWTER, A.T., D.W. FONG & D.C. CULVER, 1987. Negative phototaxis in surface and cave populations of the amphipod Gammarus minus ____ Stygologia 3, 83-88.
- VENUGOPALAN, V.P. & A.B. WAGH, 1986. Fouling Gammaridea (Amphipoda) from Bombay offshore waters. ____ Mahasagar 19, 213-215. (Not seen).
- VERMEER, K. & K. DEVITO, 1988. The importance of Paracalliosoma coecus and myctophid fishes to nesting Fork-tailed and Leach's Storm-petrels in the Queen Charlotte Islands, British Columbia. ____ J. Plankton Res. 10, 63-75.
- VERMEER, K., I. SZABO & P. GREISMAN, 1987. The relationship between plankton-feeding Bonaparte's and Mew gulls and tidal upwelling at Active Pass, British Columbia. ____ J. Plankton Res. 9, 483-501. (Feed on i.a. Parathemisto and Calliopius).
- VINOGRADOV, G.M., 1988. (Life forms of amphipods-hyperiid Hyperia and Parathemisto at different stages of ontogeny) ____ Zool. Zh. 67, 346-352. (In Russian).
- VINOGRADOV, M.E., M.V. FLINT, E.A. SHUSHKINE, V.N. TUTUBALIN & E. G. UGER, 1987. (On the comparative catchability of big volume bottles and plankton nets for vertical hauls). ____ Okeanologiya 27, 329-337. (In Russian).
- VIRNSTEIN, R.W., 1987. Seagrass-associated invertebrate communities of the southeastern USA: a review. ____ Florida mar. Res. Publs 42, 89-116.
- VIRNSTEIN, R.W. & R.V. HOWARD, 1987. Motile epifauna of marine macrophytes in the Indian River Lagoon, Florida. 1. Comparisons among three species of seagrasses from adjacent beds. ____ Bull. mar. Sci. 41, 1-12.
- VIRNSTEIN, R.W. & R. V. HOWARD, 1987. Motile epifauna of marine macrophytes in the Indian River Lagoon, Florida. 2. Comparisons between drift algae and three species of seagrasses. ____ Bull. mar. Sci. 41, 13-26.
- VONK, R. & J.H. STOCK, 1987. Amsterdam expeditions to the West Indian islands. Report 53. Psammogammarus longidactylus n. sp., a new cave amphipod (Crustacea) and other stygobiont amphipods from Bonaire. ____ Stygologia 3, 241-251. (In addition to P. longidactylus n. sp., also P. caesiculus, Saliweckelia emarginata and S. holsingeri are found in Bonaire. The genus Confodiopisa is here submerged in Psammogammarus).
- VOROB'YEVA, A.A. & R.S. NIKONOVA, 1987. (Rearing of the gammarids Dikerogammarus haemobaphes and Niphargoides maeoticus) ____ Gidrobiol. Zh. 23 (6), 50-54. (In Russian).
- VORONIN, V.N., 1986. (The Microsporidia of crustaceans). ____ Protozoologiya 10, 137-166. (In Russian, not seen).
- VOSZ, J., 1988. Zoogeographie und Gemeinschafts-analyse des Makrozoobenthos des Weddellmeeres (Antarktis). ____ Ber. Polarforsch. 45, 1-144. (Amph. 53-56).
- WAGNER, V.T. & D.W. BLINN, 1987. A comparative study of the maxillary setae for two coexisting species of Hyaella (Amphipoda), a filter feeder and a detritus feeder. ____ Arch. Hydrobiol. 109, 409-420. (H. azteca and H. montezuma).

- WARD, P.I., 1988. Sexual selection, natural selection, and body size in Gammarus pulex (Amphipoda). ____ Am. Natur. 131, 348-359. (Not seen, unfortunately).
- WESTON, D.P., 1988. Macrobenthos - sediment relationships on the continental shelf off Cape Hatteras, North Carolina. ____ Cont. Shelf Res. 8, 267-286.
- WESTIN, L. & G. ANEER, 1987. Locomotor activity patterns of nineteen fish and five crustacean species from the Baltic Sea. ____ Environm. Biol. Fishes 20, 49-66. (The single amphipod studied, Gammarus oceanicus, was considered a nocturnal species).
- WILLIAMS, D.D. & K.A. MOORE, 1986. Microhabitat selection by a stream-dwelling amphipod: A multivariate analysis approach. ____ Freshw. Biol. 16, 115-122. (On Gammarus pseudolimnaeus in S. Ontario).
- WILLIAMS, J.A., 1987. The relationship between antennal segment number and moulting in Talitrus saltator (Montagu, 1808) (Amphipoda, Talitridae). ____ Crustaceana 53, 243-252.
- WILLIAMS, J.A., 1987. A case for hormonal modulation of locomotor rhythmicity in Talitrus saltator (Crustacea: Amphipoda). ____ Comp. Biochem. Physiol. A 86, 1037-1040.
- WILLIAMS, J.A., 1988. Rhythmic locomotor activity and oxygen consumption patterns in two wrack-dwelling Orchestia (Amphipoda: Talitridae). ____ J. crust. Biol. 8, 232-238. (O. gammarellus and O. mediterranea).
- WILLIAMS, R.J., F.B. GRIFFITHS, E.J. van der WAL & J. KELLY, 1988. Cargo vessel ballast water as a vector for the transport of non-indigenous marine species. ____ Est. coast. Shelf Sci. 26, 409-420. (A number of Japanese animals, i.a. the amphipods Melita rylovae and Orchomene pacifica, were collected from sediment in ballast tanks of bulk cargo ships).
- YOUNG, J.W. & D.T. ANDERSON, 1987. Hyperiid amphipods (Crustacea: Paracarida) from a warm-core eddy in the Tasman Sea. ____ Austr. J. mar. Freshw. Res. 38, 711-725.
- ZAUKE, G-P., H-G. MEURS, K. SCHREY & H-P. BÄUMER, 1988. Influences of species, life-history status and sampling techniques on Cd, Pb, Ni, Cu and Zn in estuarine gammarids. ____ Crustaceana, Suppl. 13, 283-285.

BIBLIOGRAPHY

November 1989

- AARSET, A. V. & T. AUNAAS, 1990. Effects of osmotic stress on oxygen consumption and ammonia excretion of the Arctic sympagic amphipod Gammarus wilkitzkii. ____ Mar. Ecol. Progr. Ser. 58, 217-224.
- AARSET, A.V. & J.J. TORRES, 1989. Cold resistance and metabolic responses to salinity variations in the amphipod Eusirus antarcticus and the krill Euphausia superba. ____ Polar Biol. 9, 491-498.
- d'ACAZ, C. d'UDEKEM & P. STROOT, 1988. (Note on the expansion of Corophium curvispinum Sars, 1895, in the Meuse River). ____ Annls Soc. R. zool. Belg. 118, 171-177. (In French, not seen).
- ADAMS, J., P.J. WATT, C.J. NAYLOR & P.J. GREENWOOD, 1989. Loading constraints, body size and mating preference in Gammarus spp. ____ Hydrobiologia 183, 157-164.
- ALEKSANDROV, D.A. & E.A. IVANYUSHINA, 1989. (Life cycles of benthic amphipods in the White Sea). ____ Soviet-J. mar. Biol. 15, 9-15. (translated from Russian).
- AL-HABIB, O.A.M., F.S. HANNA & R.M.K. AL-JAMMAS, 1989. Biochemical acclimation to temperature in the stenothermal gammarid Rivulogammarus syriacus (Chevreux). ____ J. biol. Sci. Res. 20, 55-64. (Not seen).
- ANDRES, H.G., 1989. Antatelson tuberculatum sp. n., ein neuer Vertreter der Thaumatelsoninae aus der Antarktis (Crustacea: Amphipoda: Gammaridea). ____ Mitt. hamb. zool. Mus. Inst. 86, 179-184. (With key to Antatelson spp.).
- ARRESTI, A., 1989. Parhyale explorer, a new species of talitroid amphipod from the Bay of Arcachon, France. ____ Bull. Mus. natn. Hist. nat. Paris (4) 11, 101-115.
- ARRONTES, J., 1989. A new method for estimating digestive efficiency in herbivorous crustaceans. ____ Comp. Biochem. Physiol. A 94, 133-136.
- ARSUFFI, T.L. & K. SUBERKROPP, 1989. Selective feeding by shredders on leaf colonizing stream fungi. Comparison of macroinvertebrate taxa. ____ Oecologia (Berlin) 79, 30-37. (i.a. Gammarus).
- ASHLEY, D.C. & B.B. NICKOL, 1989. Dynamics of the Leptorhynchoides thecatus (Acanthocephala) suprapopulation in a Great Plains reservoir. ____ J. Parasitol. 75, 46-54. (Hyalella azteca intermediate host).

- AUNAAS, T., 1989. The effect of temperature on the osmotic regulation in the marine amphipod Gammarus oceanicus from Spitsbergen waters. ____ Pp. 49-52 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept. BECTOS-progr. 1985-88. Univ. of Trondheim, Norway.
- AUNAAS, T., 1989. Lethality in Gammarus oceanicus following exposure to water soluble fractions and water emulsions of oils and chemicals. ____ Pp. 109-114 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept. BECTOS progr. 1985-88. Univ. of Trondheim, Norway.
- AUNAAS, T., 1989. Metabolic effects of water soluble fractions and water emulsions of oil and chemicals on the amphipod Gammarus oceanicus from arctic waters. ____ Pp. 131-134 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept. BECTOS progr. 1985-88. Univ. of Trondheim, Norway.
- AUNAAS, T., 1989. The effect of water soluble fractions and water emulsions of oil and chemicals on inorganic ions and free amino acids in the amphipod Gammarus oceanicus in arctic waters. ____ Pp. 153-165 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept. BECTOS progr. 1985-88. Univ. of Trondheim, Norway.
- AUNAAS, T., 1989. The effects of pre-exposure to water soluble fractions of oils and chemicals on the osmoregulation of the amphipod Gammarus oceanicus in low salinity seawater. ____ Pp. 201-207 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept. BECTOS progr. 1985-88. Univ. of Trondheim, Norway.
- AUNAAS, T., J.F. BÖRSETH, J-P. DENSTAD & A.V. AARSET, 1989. Energetic costs of osmotic regulation in the amphipod Gammarus oceanicus from arctic waters. ____ Pp. 39-43 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept. BECTOS progr. 1985-88. Univ. of Trondheim, Norway.
- AUNAAS, T., J.F. BÖRSETH, J-P. DENSTAD & A.V. AARSET, 1989. Effect of acute osmotic stress and temperature on the metabolism of Gammarus oceanicus from Svalbard waters. ____ Pp. 45-48 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept. BECTOS progr. 1985-88. Univ. of Trondheim, Norway.
- AUNAAS, T., J.F. BÖRSETH, J-P. DENSTAD & A.V. AARSET, 1989. The effects of a light diesel oil spill on arctic marine amphipods and fish. ____ Pp. 227-233 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept. BECTOS progr. 1985-88. Univ. of Trondheim, Norway (mainly Gammarus oceanicus).
- BARNARD, J.L., 1989. Rectification of Halirages regis and H. huxleyanus (Crustacea: Amphipoda), from marine Antarctica, with description of a new genus Austroregia. ____ Proc. biol. Soc. Wash. 102, 701-715. (The family Gammarellidae is revived for Gammarellus (type), Gondogeneia, Chosroes, and Austroregia n. gen. The family is mainly based on most characteristic and apomorphic calceoli. Austroregia n. gen. has Atylus huxleyanus as type, and also contains Bovallia regis).
- BARNARD, J.L. & J.D. THOMAS, 1989. Four species of Synopiidae from the Caribbean region (Crustacea: Amphipoda). ____ Proc. biol. Soc. Wash. 102, 362-374. (Synopia ultramarina, S. scheeleana, Tiron bellairisi and Garosyrhoe bigorra (transferred from Syrrhoites, with G. disjuncta as junior synonym). With a key to Synopia spp.).
- BARNARD, J.L. & J.D. THOMAS, 1989. A new species, Ampelisca burkei (Crustacea Amphipoda) from Florida. ____ Proc. biol. Soc. Wash. 102, 375-384.
- BEK, T.A., 1988. (New records of amphipod species in the White Sea). ____ Vestn. MGU (Biol.) 1988-2, 42-45. (In Russian, not seen. Argissa hamatipes and Monoculodes packardii).
- BELLAN-SANTINI, D. & J-C. DAUVIN, 1989. Distribution verticale et répartition biogéographique de crustacés holobenthiques filtreurs: exemple des amphipodes du genre Ampelisca, groupe zoologique à forte spéciation. ____ Bull. Soc. géol. Fr. (8) 5, 561-568.
- BENNETT, B.A., 1989. The diets of fish in three southwestern Cape estuarine systems. ____ S. Afr. J. Zool. 24, 163-177.
- BIERNBAUM, C.K., 1989. Distribution and seasonality of branchiopod and malacostracan crustaceans of the Santee National Wildlife refuge, South Carolina. ____ Brimleyana 15, 7-30. (Not seen).
- BOATES, J.S. & P.C. SMITH, 1989. Crawling behavior of the amphipod Corophium volutator and foraging by Semipalmated Sandpipers, Calidris pusilla. ____ Can. J. Zool. 67, 457-462.
- BORGMANN, U. & M. MUNAWAR, 1989. A new standardized sediment bioassay protocol using the amphipod Hyalella azteca (Saussure). ____ Hydrobiologia 188-189, 425-431.
- BOROWSKY, B., 1988. Delaying copulation in the amphipod Gammarus palustris: Effects on female fecundity and consequences for the frequency of amplexus. ____ Mar. Behav. Physiol. 13, 359-368.
- BOROWSKY, B., 1989. The effect of residential tubes on reproductive behaviors in Microdeutopus gryllotalpa (Costa) (Crustacea: Amphipoda). ____ J. exp. mar. Biol. Ecol. 128, 117-125.
- BOROWSKY, R. & B. BOROWSKY, 1987. Feeding by the salt marsh amphipod, Gammarus palustris, on Enteromorpha and Ulva. ____ Am. Zool. 27, 55.
- BOROWSKY, R. & B. BOROWSKY, 1988. Carbohydrate feeding preferences of Gammarus amphipods. ____ Am. Zool. 28, 28.

- BOROWSKY, R. & M.M. GUARNO, 1989. Excess amylase in Gammarus palustris (Crustacea: Amphipoda); its release into and possible roles in the environment. ____ Mar. Biol. 101, 529-534.
- BÖRSETH, J.F. & T. AUNAAS, 1989. Physiological effects of an oil slick and its chemical treatment on marine gammarids, Gammarus locusta, and eggs of plaice, Pleuronectes platessa. ____ Pp. 221-225 in K.E. Zachariassen (ed). Biological effects of chemical treatments of oilspills at sea. Rept. BECTOS progr. 1985-1988. Univ. of Trondheim, Norway.
- BOUHIN, H. & J-C. MEYRAN, 1989. Change of in vitro translation products from the RNA in the posterior caeca of the crustacean Orchestia during the molt cycle. ____ Comp. Biochem. Physiol. B. 93, 905-910.
- BOUSFIELD, E.L. 1989. Revised morphological relationships within the amphipod genera Pontoporeia and Gammaracanthus and the 'glacial relict' significance of their postglacial distributions. ____ Can. J. Fish. aq. Sci. 46, 1714-1725. (The old genus Pontoporeia is here divided into Pontoporeia s. str., Monoporeia n. gen. (type P. affinis) and Diporeia n. gen. (type P. hoyi). The family Gammaracanthidae n. fam. (Eusiroidea) contains the genera Gammaracanthus, with subgenera G. s. str. and Pseudacanthus n. subg. (type: G. aestuariorum), and Relictacanthus n. gen. (type: G. relictus). Several theories on the origin and distribution history of these so-called 'glacial relicts' are reviewed).
- BOUSFIELD, E.L. & J.R. HOLSINGER, 1989. A new crangonyctid amphipod crustacean from hypogean fresh waters of Oregon. ____ Can. J. Zool. 67, 963-968. (Stygonyx courtneyi n. gen. n. sp. from NW Oregon, USA, close to Stygobromus).
- BRINK, F.W.B. v.d., G. v.d. VELDE & A. bij de VAATE, 1989. A note on the immigration of Corophium curvispinum Sars, 1895 (Crustacea: Amphipoda) into the Netherlands. ____ Bull. zool. Mus. Univ. A'dam 11, 211-214.
- BRUCE, L., 1989. Invertebrates associated with the thinstripe hermit Clibanarius vittatus (Bosc) (Crustacea: Decapoda: Dlogenidae) from the barrier islands of Mississippi. ____ Gulf Res. Repts 8, 213-217. (Very few amphipods: Talorchestia sp. and Hyale sp.).
- BYERS, T. & R.W. PRACH, 1988. Diet of the Kelp Snailfish, Liparis tunicatus, in Jones Sound, Canadian High Arctic. ____ Can. Field-Nat. 102, 242-245. (Amph. abundant in diet).
- CAINE, E.A., 1989. Caprellid amphipod behavior and predatory strikes by fish. ____ J. exp. mar. Biol. Ecol. 126, 173-180.
- CAINE, E.A., 1989. Relationships between wave activity and robustness of caprellid amphipods. ____ J. crust. Biol. 9, 425-431.
- CAMILLERI, J., 1989. Leaf choice by crustaceans in a mangrove forest in Queensland. ____ Mar. Biol. 102, 453-460. (i.a. Orchestia and Melita).
- CARLTON, C.E. & E.H. SCHMITZ, 1989. Anatomy of the extrinsic gut musculature of Gammarus minus (Crustacea: Amphipoda). ____ J. Morphol. 200, 87-92.
- CHAMIER, A-C., D.W. SUTCLIFFE, & J.P. LISHMAN, 1989. Changes in sodium, potassium, calcium, magnesium and aluminum content of Gammarus pulex (L.). ____ Freshw. Biol. 21, 181-190.
- CHESS, J.R., 1989. Aciconula acanthosoma, new species, a caprellid amphipod from southern California, with notes on its biology. ____ J. crust. Biol. 9, 662-665.
- CLANZIG, S., 1987. Inventaire des invertébrés d'une lagune méditerranéenne des côtes de France, biocenoses et confinement: l'Etang de Salses-Leucate (Roussillon). ____ Thèse, Ecole Pratique des Hautes Etudes, Paris. 468 pp. (Not seen).
- CLARK, M.R., K.J. KING & P.J. McMILLAN, 1989. The food and feeding relationships of black oreo, Allocyttus niger, smooth oreo, Pseudocyttus maculatus, and eight other fish species from the continental slope of the south-west Chatham Rise. New Zealand. ____ J. Fish Biol. 35, 465-484. (Amphipods important prey).
- COLEMAN, C.O., 1989. Gnathiphimedia mandibularis K.H. Barnard, 1930, an Antarctic amphipod (Acanthonotozomatidae, Crustacea) feeding on Bryozoa. ____ Antarct. Sci. 1, 343-344.
- COLEMAN, C.O. 1989. On the nutrition of two Antarctic Acanthonotozomatidae (Crustacea: Amphipoda). Gut contents and functional morphology of mouthparts. ____ Polar Biol. 9, 287-294. (Echiniphimedia hodgsoni and Maxilliphimedia longipes).
- COLEMAN, C.O., 1989. Burrowing, grooming, and feeding behaviour of Paraceradocus, an antarctic amphipod genus. ____ Polar Biol. 10, 43-48.
- CONLAN, K.E., 1989. Delayed reproduction and adult dimorphism in males of the amphipod genus Jassa (Corophioidea: Ischyroceridae): an explanation for systematic confusion. ____ J. crust. Biol. 9, 601-625.
- CORKUM, L.A., 1989. Patterns of benthic invertebrate assemblages in rivers of northwestern North America. ____ Freshw. Biol. 21, 191-206.
- CORKUM, L.D. & E.C. HANES, 1989. A laboratory aeration system for rearing aquatic invertebrates. ____ Entom. News 100, 169-172. (Not seen).
- COSTELLO, M.J. & A.A. MYERS, 1989. Breeding periodicity and sex ratios in epifaunal marine Amphipoda in Lough Hyne, Ireland. ____ Est. coast. Shelf Sci. 29, 409-419.
- CRESPI, B.J., 1989. Causes of assortative mating in amphipods. ____ Anim. Behav. 38, 980-1000.

- CULVER, D.C. & D.W. FONG, 1986. Why all cave animals look alike. ____ *Stygologia* **2**, 208-216.
- CYRUS, D.P., 1988. Episodic events and estuaries: effects of cyclonic flushing on the benthic fauna and diet of *Solea bleekeri* (Teleostei) in Lake St. Lucia on the south-eastern coast of Africa. ____ *J. Fish Biol.* **33** Suppl. 1, 1-7.
- CYRUS, D.P. & T.J. MARTIN, 1988. Distribution and abundance of the benthos in the sediment of Lake Cubhu: a freshwater coastal lake in Zululand, South Africa. ____ *J. limnol. Soc. S. Afr.* **14**, 93-101. (*Grandidierella lignorum* and *Corophium triaenonyx* codominant).
- DAUVIN, J.-C., 1989. Life cycle, dynamics and productivity of Crustacea-Amphipoda from the western English Channel. 5. *Ampelisca sarsi* Chevreux. ____ *J. exp. mar. Biol. Ecol.* **128**, 31-56.
- DAUVIN, J.-C. & D. BELLAN-SANTINI, 1988. Illustrated key to *Ampelisca* species from the north-eastern Atlantic. ____ *J. mar. biol. Ass. UK* **68**, 659-676. (Deals with 52 spp. from W. Africa to N. Norway. The paper also contains a phenogram illustrating the phenetic similarities among the spp.).
- DE BROYER, C. & M. KLAGES, 1990. Studies in amphipod biology. ____ Pp. 113-115 in W. Arntz, W. Ernst & I. Hempel (eds). The expedition Antarktis VII/4 (Epos leg 3) and VII/5 of RV "Polarstern" in 1989. ____ *Ber. Polarf.* **68**, 1-214.
- DETHIER, M., 1988. (Benthic macroinvertebrates of the Rhone river in Geneva (Switzerland). 2. Faunistical aspects). ____ *Rev. suisse Zool.* **95**, 1117-1132. (In French, not seen).
- DEXTER, D.M., 1989. The sandy beach fauna of Egypt. ____ *Est. coast. Shelf Sci.* **29**, 261-271.
- DICK, J.T.A. & R.W. ELLWOOD, 1989. Assessments and decisions during mate choice in *Gammarus pulex* (Amphipoda). ____ *Behaviour* **109**, 235-246.
- DICK, J.T.A., D.E. IRVINE & R.W. ELWOOD, 1990. Differential predation by males on moulted females may explain the competitive displacement of *Gammarus duebeni* by *G. pulex* (Amphipoda). ____ *Behav. Ecol. Sociobiol.* **26**, 41-45.
- DITTRICH, B., 1987. Populationsdynamikk der parasitischen Amphipoden *Hyperia galba* (Montagu). ____ *Verh. dtsh. zool. Ges.* **80**, 293-294.
- DUNHAM, P.J. & A. HURSHMAN, 1988. Amphipod mate guarding decisions: deprivation versus uncertainty. ____ *Anim. Behav.* **36**, 609-611.
- DUNTON, K.H., S.M. SAUPE, A.N. GOLIKOV, D.M. SCHELL & S.V. SCHONBERG, 1989. Trophic relationships and isotopic gradients among arctic and subarctic marine fauna. ____ *Mar. Ecol. Progr. Ser.* **56**, 89-97.
- DUTRA, R. Roxo Coutinho, 1988. (The vagile fauna of the phytal of *Pterocladia capillacea* (Rhodophyta, Gelidiaceae) from Ilha do Mel, Parana, Brazil) ____ *Rev. bras. Biol.* **48**. (In Portuguese).
- EMELYANOVA, L.V. & Yu.V. PLIGIN, 1987. (Finding of *Talorchestia deshayesi* (Crustacea, Talitridae) in Kokhovskaya Reservoir). ____ *Vestn. Zool.* 1987 (6), 51- (In Russian, not seen).
- ESSELINK, P., J. van BELKUM & K. ESSINK, 1989. The effect of organic pollution on local distribution of *Nereis diversicolor* and *Corophium volutator*. ____ *Neth. J. Sea Res.* **23**, 323-332.
- FONG, D.W., 1988. Morphological evolution of the amphipod *Gammarus minus* in caves: Quantitative genetic analysis. ____ *Am. Midl. Nat.* **121**, 361-378.
- FONG, D.W. & D.C. CULVER, 1985. A reconsideration of Ludwig's differential migration theory of regressive evolution. ____ *NSS Bull.* **47**, 123-127.
- FOSTER, J.M., 1989. *Acanthohaustorius uncinus*, a new species of sand-burrowing amphipod from the northern Gulf of Mexico, with notes on its ecology (Haustoriidae: Haustoriinae). ____ *Gulf Res. Rep.* **8**, 189-197.
- FRANZ, D.R., 1989. Population density and demography of a fouling community amphipod. ____ *J. exp. mar. Biol. Ecol.* **125**, 117-130. (*Jassa marmorata*).
- FRANZ, D.R. & W.H. HARRIS, 1988. Seasonal and spatial variability in macrobenthos communities in Jamaica Bay, New York - An urban estuary. ____ *Estuaries* **11**, 15-28.
- FRANZ, D.R. & Y. MOHAMED, 1989. Short-distance dispersal in a fouling community amphipod crustacean, *Jassa marmorata* Holmes. ____ *J. exp. mar. Biol. Ecol.* **133**, 1-13.
- GARTSHORE, N.A., W.K. STEELE & N.T. KLAGES, 1988. Summer diet of the Salvin's Prion at sub-antarctic Marion Island. ____ *S. Afr. J. Sci.* **23**, 309-313. (Main prey *Parathemisto gaudichaudii*).
- GIBBONS, M.J., 1988. The impact of wave exposure on the meiofauna of *Gelidium pristoides* (Turner) Kuetzing (Gelidiales: Rhodophyta). ____ *Est. coast. Shelf Sci.* **27**, 581-594.
- GINET, R., 1988. Description d'un neotype et choix d'une nouvelle localité-type pour le crustacé stygobie *Niphargus ciliatus* Chevreux, 1906 (Amphipoda). ____ *Bull. mens. Soc. linn. Lyon* **57**, 215-231.
- GOOCH, J.L., 1989. Genetic differentiation in relation to stream distance in *Gammarus minus* (Crustacea, Amphipoda) in Appalachian watersheds. ____ *Arch. Hydrobiol.* **114**, 505-519.

- GROSSO, L.E. & H.R. FERNANDEZ, 1988. A case of sympatry of three species in the genus Bogidiella (Crustacea, Amphipoda) in the northwest of Argentina, with descriptions of two new species. ____ Stygologia **4**, 64-78. (Not seen. Deals with B. tucumanensis, B. (Stygogidiella) horcomollensis n. sp. and B. (Dyctiogidiella) ringueleti n. sp., all from the Tucuman area of Argentina).
- GULLIKSEN, B. & O.J. LÖNNE, 1989. Distribution, abundance, and ecological importance of marine sympagic fauna in the Arctic. ____ Rapp. P-v. Réunion Cons. int. Explor. Mer. **188**, 133-138.
- HACKSTEIN, E., 1988. Die Veränderung populations- dynamischer Parameter bei Gammarus tigrinus Sexton (Crustacea: Amphipoda) als Ausdruck sublethaler Effekte durch die Wechselwirkung von Temperatur und cadmiumkontaminierten Futter. ____ Int. Rev. ges. Hydrobiol. **73**, 213-227.
- HAEFNER, J.W. & L. C. ABBOTT, 1989. Extrapolation of laboratory pH dose-response data to fluctuating environments: comparisons with a null model. ____ Can. J. Fish. aq. Sci. **46**, 1499-1509. (The amphipod Gammarus lacustris was chosen as a pH sensitive species).
- HALLBERG, E. & R. ELOFSSON, 1989. Construction of the pigment shield of the crustacean compound eye: a review. ____ J. crust. Biol. **9**, 359-372.
- HARDING, P.T. & S.L. SUTTON, 1988. The spread of the terrestrial amphipod Arcitalitrus dorrieni in Britain and Ireland: watch this niche. ____ Isopoda **1988-2**, 7-10. (Not seen. Can somebody get me a copy?).
- HEUSS, K., 1986. Erstfunde des Flohkrebsses Gammarus tigrinus Sexton (Crustacea, Amphipoda) in Mittelfranken. ____ Natur Mensch (Nuremburg) **1986**, 95-96. (Not seen).
- HILL, C., 1988. Life cycle and spatial distribution of the amphipod Pallasea quadrispinosa in a lake in northern Sweden. ____ Holarct. Ecol. **11**, 298-304.
- HILL, C. & R. ELMGREN, 1987. Vertical distribution in the sediment in the co-occurring benthic amphipods Pontoporeia affinis and P. femorata. ____ Oikos **49**, 221-229.
- HIWATARI, T. & T. KAJIHARA, 1988. Experimental studies on the growth and breeding of Hyale barbicornis (Amphipoda, Crustacea) at different temperatures. ____ Jap. Soc. scient. Fish. **54**, 39-43.
- HOLMQUIST, J.G., 1989. Grooming structure and function in some terrestrial Crustacea. ____ Pp. 95-114 in B.E. FELGENHAUER, L. WATLING & A.B. THISTLE (eds). Functional morphology of feeding and grooming in Crustacea. A.A. Balkema, Rotterdam. (i.a. Talitroidea).
- HOLSINGER, J.R., 1989. Allocrangonyctidae and Pseudocrangonyctidae, two new families of holarctic subterranean amphipod crustaceans (Gammaridea), with comments on their phylogenetic and zoogeographic relationship. ____ Proc. biol. Soc. Wash. **102**, 947-959. (Both families are split off from the Crangonyctidae s.l. The Allocrangonyctidae are monotypic, for Allocrangonyx, with the North American spp. A. pellucidus (type) and A. hubrichti, here redescribed. The Allocrangonyctidae are of uncertain affinities, but may be aberrant hadzioids. The East Asian Pseudocrangonyctidae consist of Pseudocrangonyx (type P. asiaticus, redescribed) and Procrangonyx (type P. japonicus, no material extant). The Pseudocrangonyctidae are allied to the Crangonyctidae).
- HUGHES, R.G. & I.M. HORSFALL, 1990. Differences in the swimming behaviour of the amphipod Corophium volutator from different populations ____ J. mar. biol. Ass. UK **70**, 143-148.
- IKEDA, T., 1989. Are Antarctic zooplankton metabolically more cold-adapted than Arctic zooplankton? An intra-genetic comparison of oxygen consumption rates. ____ J. Plankton Res. **11**, 619-624. (Not seen. The author's answer to the question is 'Probably not').
- INGLIS, G., 1989. The colonization and degradation of stranded Macrocystis pyrifera (L.) C. Ag. by the macrofauna of a New Zealand sandy beach. ____ J. exp. mar. Biol. Ecol. **125**, 203-218.
- JACQUES, F., 1989. The setal system of crustaceans: Types of setae, groupings, and functional morphology. ____ Pp. 1-13 in B.E. FELGENHAUER, L. WATLING & A.B. THISTLE (eds). Functional morphology of feeding and grooming in Crustacea. A.A. Balkema, Rotterdam.
- JANEVA, I.J., 1987. Gammarus arduus Kar. & Pink. (Amphipoda, Gammaridae), a new bioindicator of saprobity of Bulgarian river fauna. ____ Khidrobiol. **30**, 60-61. (Not seen).
- JARAMILLO, E., 1987. Sandy beach macroinfauna from the Chilean coast: zonation patterns and zoogeography. ____ Vie Milieu **37**, 165-174.
- JAZDZEWSKI, K., & A. KONOPACKA, 1989. Gammarus leopoliensis, new species (Crustacea, Amphipoda) from eastern Carpathians. ____ Bull. zool. Mus. Univ. A'dam **11**, 185-196. (Close to G. kischineffensis).
- JEWETT, S.C., R.H. DAY & H.M. FEDER, 1989. Feeding biology of the Blackfin Sculpin (Malacocottus kincaidj Gilbert and Thompson, 1905) and the Spinyhead Sculpin (Dasycottus setiger Bean, 1980) in the Northeastern Gulf of Alaska. ____ Pacif. Sci. **43**, 144-151.
- JO, Y.W., 1989. Shallow-water phoxocephalid Amphipoda (Crustacea) of Korea. ____ Bijdr. Dierk. **59**, 97-125. (Deals with Mandibulophoxus mai n. sp., M. hongae n. sp., Grandifoxus malipoensis n. sp., G. cuspis n. sp. and G. bangpoensis n. sp. A key to Grandifoxus is provided (excl. G. nasuta, which does not belong in this genus). Mandibulophoxus latipes is provisionally transferred to Basuto).
- JO, Y.W., 1990. Oedicerotid Amphipoda (Crustacea) from shallow waters of Korea. ____ Beaufortia **39**, 155-200. (Deals with Monoculodes koreanus n. sp., M. muwoni n. sp., M. dentimanus n. sp., Periculodes seohae n. sp., Synchelidium lenorostratum n. rank (was ssp. of S. miraculum), S. carinorostrum n. sp., S. triostegitum n. sp., and Chitonomandibulum emargicoxa n. gen., n. sp. (close to Synchelidium). Keys to Pacific oedicerotid genera and Monoculodes spp. and to worldwide Periculodes and Synchelidium spp. are provided).

- JOHNSON, I. & M.B. JONES, 1989. Effects of zinc/salinity combinations on zinc regulation in Gammarus duebeni from the estuary and the sewage treatment works at Looe, Cornwall. ____ J. mar. biol. Ass. UK 69, 249-260.
- JOHNSON, R.K. & T. WIEDERHOLM, 1989. Long-term growth oscillation of Pontoporeia affinis Lindström (Crustacea: Amphipoda) in Lake Mälaren. ____ Hydrobiologia 175, 183-194.
- JONES, M.B. & G.D. WIGHAM, 1988. Colonization by estuarine amphipods of a sewage treatment area. ____ Bull. est. brackish Water Sci. Ass. 50, 29-33. (Not seen).
- JONES, R. & D.C. CULVER, 1989. Evidence for selection on sensory structures in a cave population of Gammarus minus (Amphipoda). ____ Evolution 43, 688-693.
- KARAMAN, G.T., 1987. New species of family Niphargidae and new localities of some other subterranean gammaridean species from Yugoslavia (Contribution to the knowledge of the Amphipoda 166). ____ Glasn. Republ. Zavoda Zast. Prirodo-Prirodnjackog Muz. Titograd 19 (1986), 15-32. (Niphargus carcerarius n. sp., in 'Orniphargus' group, from Crna Gora, Yugoslavia. Faunistic data on 5 other Niphargus spp. and Hadzia f. fragilis).
- KARAMAN, G.S., 1988. The genus Paraphoxus Sars, 1891 (fam. Phoxocephalidae) in the Mediterranean Sea (Contribution to the knowledge of the Amphipoda 171). ____ Glasnik Sect. nat. Sci. Montenegrin Acad. Sci. Arts 6, 139-161. (With descriptions of P. oculatus (incl. P. maculatus) and P. lincolni n. sp. (Adriatic Sea). P. oculatus auct. may consist of two species, but lack of topotypic Arctic material prevented the author from deciding this).
- KARAMAN, G.S. 1988-89. Two cavernicolous species of the genus Niphargus Schiödte from Yugoslavia, Niphargus jalzici, n. sp. and N. lunaris G. Kar. 1985 (fam. Niphargidae). (Contribution to the knowledge of the Amphipoda 176). ____ Bull. Mus. nat. Belgrade 43-44, 48-61. (Type locality of N. jalzici is Croatia. It belongs to the orcinus-group).
- KARAMAN, G.S. 1988. The new species of the genus Niphargus Schiödte (Gammaridea, fam. Niphargidae) from Italy and Yugoslavia (Contribution to the knowledge of the Amphipoda 177). ____ Poljoprivreda i Sumarstvo 34, 11-31. (N. poianoi n. sp. from Emilia Romagna, Italy; N. arcanus n. sp. from Croatia, Yugoslavia, both in the orcinus-group).
- KARAMAN, G.S., 1989. Metacrangonyx ortalii, n. sp., a new subterranean member of the family Crangonyctidae from Dead Sea region. Contribution to the knowledge of the Amphipoda 178. ____ Studia marina, Kotor 20, 33-49. (The genera Metacrangonyx and Afrocrangonyx are provisionally upheld, pending revision).
- KARAMAN, G.S., 1988. New genera and species of the subterranean family Bogidiellidae from the Near East. (Contribution to the knowledge of the Amphipoda 179). ____ Studia marina, Kotor 19, 31-51. (Hebraegidiella bromleyana n. gen. n. sp., close to Bogidiella, from the Dead Sea area. Bogidiella (?B.) capia n. sp. from the same area. The new genus Nubigidiella is erected for Bogidiella nubica; it is close to Maghrebiella).
- KARAMAN, G.S., 1988. Two new species of genus Bogidiella Hert. from Sardinia and France, with remarks to B. vandeli Coineau 1968 (Gammaridea, fam. Bogidiellidae) (Contribution to the knowledge of the Amphipoda 180). ____ Poljoprivreda i Sumarstvo 34, 25-41. (B. vandeli is redescribed from type locality, B. (B.) calicali n. sp. from same locality in Sardinia includes the presumed males of B. vandeli s. Coineau. B. (B.) nicolae n. sp. from the Tech in S. France is B. skopljensis s. Coineau 1968).
- KARAMAN, G.S., 1988. The new genus of the family Bogidiellidae (Gammaridea) from coastal seawater (mesopsammon) of Italy, Aurobogidiella, n. gen. (Contribution to the knowledge of the Amphipoda 182). ____ Poljoprivreda i Sumarstvo 34, 93-105. (Aurobogidiella n. gen. is erected for Bogidiella italica, here redescribed).
- KARAMAN, G.S., 1987. New species of the family Bogidiellidae (Gammaridea) from Yugoslavia, Bogidiella serbica, n. sp. (Contribution to the knowledge of the Amphipoda 183). ____ Bull. Mus. Hist. nat. Belgrade 42, 37-50. (near Pec, Serbia).
- KARAMAN, G.S., 1987. A new species of genus Melita Leach (fam. Melitidae) from Bermuda and Fiji Islands. Contribution to the knowledge of the Amphipoda 173. ____ Bull. Mus. Hist. nat. Belgrade 42, 19-35. (Melita stocki n. sp. from Bermuda, M. persona n. sp. (= M. lanaterga s. Kar. 1981), also from Bermuda, M. myersi n. sp. (= M. zeylanica s. Myers 1985) from Fiji).
- KARAMAN, G.S., 1989. Bogidiella sketi, new freshwater species of the family Bogidiellidae from Dalmatia (Yugoslavia) with remarks to some other Bogidiella species (Contribution to the knowledge of the Amphipoda 188). ____ Poloprivreda i Sumarstvo 35, 49-60.
- KARAMAN, G.S. & B. SKET, 1989. Niphargus species (Crustacea: Amphipoda) of the Kvarner - Velebit Islands (NW Adriatic, Yugoslavia). ____ Biol. Vestn. 37, 19-36. (Descriptive notes on N. longicaudatus, N. wolffi and N. steueri liburnicus n. ssp. (Island Krk)).
- KAWABARA, R. & Y. AKIMOTO, 1986. The offshore environment of Tungkong, Southwest Taiwan. 2. Macrobenthos. ____ Proc. 1. Asian Fish. Forum, Manila, 26-31 May 1986, 193-198. (Not seen. Apparently many Amph).
- KEVREKIDIS, T. & A. KOUKOURAS, 1988. Life cycle and reproduction of Gammarus aequicauda (Crustacea - Amphipoda) in the Evoros Delta (northeastern Greece). ____ Isr. J. Zool. 35, 137-150.
- KIERSTEAD, W.G. & F. BÄRLOCHER, 1989. Ecological effects of pentachlorophenol on the brackishwater amphipod Gammarus tigrinus. ____ Arch. Hydrobiol. 115, 149-156.

- KIM, C.B. & W. KIM, 1989. A new species of the genus Ceradocus (Crustacea, Amphipoda, Melitidae) from Korea. ____ Korean J. syst. Zool. 5, 173-181. (Ceradocus (Denticeradocus) koreanus n.sp from near Pusan).
- KIM, H.S. & C.B. KIM, 1987. Marine gammaridean Amphipoda (Crustacea) of Cheju Island and its adjacent waters, Korea. ____ Korean J. syst Zool. 3, 1-23. (Deals with Ampithoe lacertosa, A. valida, Aoroides columbiae, Jassa falcata, Paramoera koreana, Pontogeneia rostrata, Colomastix lunailo, Elasmopus pecteniscus, Maera pacifica, Melita koreana, Hyalae crassicornis, H. punctata, H. rubra, Parhyalella pietschmanni, Platorchestia platensis and P. pachypus, many of them new to Korea).
- KIM, H.S. & C.B. KIM, 1988. Marine gammaridean Amphipoda (Crustacea) of the family Ampithoidae from Korea. ____ Korean J. syst. Zool., spec. Issue 2, 107-134. (Deals with Ampithoe lacertosa, A. valida, A. v. shimizuensis (reduced to subspec.-status), A. brevipalma n. sp. (Pangchukp'o), A. koreana n. sp. (Ayajin), A. youngsanensis n. sp. (Youngsando Ist.), A. ramondi, Peramphithoe baegryeongensis n. sp. (Paengnyondo Isl.), P. namhaensis n. sp. (Upper Chujade Isl.), P. tea, P. orientalis and Sunamphithoe plumulosa).
- KIM, S.L. & J.S. OLIVER, 1989. Swarming benthic crustaceans in the Bering and Chukchi Seas and their relation to geographic patterns in Gray whale feeding. ____ Can. J. Zool. 67, 1531-1549.
- KOCH, H., 1989. Desiccation resistance of the supralittoral amphipod Traskorchestia traskiana (Stimpson, 1857). ____ Crustaceana 56, 162-175.
- KOCH, H., 1989. The effect of tidal inundation on the activity and behavior of the supralittoral talitrid amphipod Traskorchestia traskiana (Stimpson, 1857). ____ Crustaceana 57, 295-303.
- KÖHN, J. & F. GOSSELCK, 1989. Bestimmungsschlüssel der Malakostraken der Ostsee. ____ Mitt. zool. Mus. Berlin 65, 3-114. (Amph. 39-98. With data on biology and distribution, and illustrations of all species. A most useful and welcome paper).
- KORCZYNSKI, R.E., M.J. LAWRENCE & G.E. HOPKY, 1989. Range extensions of some peracaridan crustaceans in the Beaufort Sea. ____ Crustaceana 56, 215-219.
- KUSANO, H., T. KUSANO & Y. WATANABE, 1987. Life history and reproduction of Jesogammarus spinipulps (sic!) (Anisogammaridae: Amphipoda) inhabiting a lowland pond in Tokyo City. ____ Jap. J. Limnol. 48, 117-126. (Not seen).
- LABERGE, R.J. & J.D. McLAUGHLIN, 1989. Hyalella azteca (Amphipoda) as an intermediate host of the nematode Streptocara crassicauda. ____ Can. J. Zool. 67, 2335-2340.
- LALANA RUEDA, R., 1985. (Study of the epifauna from the coastal lagoons 'Tolete' and 'El Basto'). ____ Rev. Invest. mar. 6, 19-28. (In Spanish)
- LAM, P.K.S. & H.H.T. MA, 1989. Some observations on the life cycle and population dynamics of Talitroides topitotum (Burt) (Amphipoda: Talitridae) in Hong Kong. ____ J. nat. Hist. 23, 1087-1092.
- LANCRAFF, T.M., J.J. TORRES & T.L. HOPKINS, 1989. Micronekton and makrozooplankton in the open waters near Antarctic Ice Edge zones (AMERIEZ 1983 and 1986). ____ Polar Biol. 9, 225-233.
- LAND, M.F., 1989. The eyes of hyperiid amphipods : Relations of optical structures to depth. ____ J. comp. Physiol. A. 164, 751-762.
- LANDRUM, P.F., 1988. Toxicokinetics of organic xenobiotics in the amphipod, Pontoporeia hoyi: role of physiological and environmental variables. ____ Aquat. Toxicol. 12, 245-271.
- LANDRUM, P.F., 1988. Bioavailability and toxicokinetics of polycyclic aromatic hydrocarbons sorted to sediments for the amphipod Pontoporeia hoyi. ____ Environm. Sci. Toxicol. 23, 588-595.
- LANDRUM, P.F., W.R. FAUST & B.J. EADIE, 1989. Bioavailability and toxicity of a mixture of sediment-associated chlorinated hydrocarbons to the amphipod Pontoporeia hoyi. ____ Am. Soc. Testing & Materials, spec. techn. Publ. 1027, 315-329.
- LANDRUM, P.F., S.R. NIHART, B.J. EADIE & L.R. HERCHE, 1987. Reduction in bioavailability of organic contaminants by the amphipod Pontoporeia hoyi by dissolved organic matter of sediment interstitial waters. ____ Environm. Toxicol. Chem. 6, 11-20.
- LAZIM, M.N., A.D. SHUKER-KHAN & I.S. SAEED, 1988. Some of them acrobenthic invertebrates from springs and streams of northern parts of Iraq. ____ J. biol. Sci. Res. 19, 425-427. (Not seen. Amph. ?).
- LEITE, F.P.P. & Y. WAKABARA, 1989. Aspects of marsupial and post-marsupial development of Hyalae media (Dana) 1853 (Hyalidae, Amphipoda). ____ Bull. mar. Sci. 45, 85-97.
- LIN, J.H. & R.X. CHEN, 1988. (Distribution of planktonic Amphipoda in western Taiwan Strait). ____ J. Oceanogr. Taiwan Strait 7, 324-330. (In Chinese, not seen).
- LLAZE, R. & M. DAUMAS, 1985. De la presence de Neocyamus physeteris (Pouchet, 1888), ectoparasite du cachalot Physeter macrocephalus, en Méditerranée. ____ Actes Congr. natn. Soc. Savantes, Sect. Sci. 110, 33-42.
- LOCKE, A. & S. COREY, 1989. Amphipods, isopods and surface currents: A case for passive dispersal in the Bay of Fundy, Canada. ____ J. Plankton Res. 11, 419-430.

- LOP, A.F. & J.L. OLIVER, 1989. Isozyme differentiation among sibling species and among populations of the *Echinogammarus berilloni* - group (Crustacea, Amphipoda). ____ Z. zool. Syst. Evol. f. 27, 282-296.
- LOPEZ, G. & R. ELMGREN, 1989. Feeding depths and organic absorption for the deposit-feeding benthic amphipods *Pontoporeia affinis* and *Pontoporeia femorata*. ____ Limnol. Oceanogr. 34, 982-991.
- LOVETT, D.L. & D.L. FELDER, 1989. Application of regression techniques to studies of relative growth in crustaceans. ____ J. crust. Biol. 9, 529-539.
- LOWRY, J.K. & P.B. BERENTS, 1989. A redescription of *Cerapus tubularis* Say, 1817, based on material of the first reviewer, S.I. Smith, 1880. (Crustacea: Amphipoda: Corophioidea). ____ J. nat. Hist. 23, 1341-1352. (A neotype is selected and the genus redefined. *C. 'tubularis'* s. Barnard 1962, from S. California, is a different, as yet undescribed species).
- LOWRY, J.K. & G.C.B. POORE, 1989. First Ingolfiellids from the Southern Pacific (Crustacea: Amphipoda) with a discussion of their systematics. ____ Proc. biol. Soc. Wash. 102, 933-946. (Deals with *Ingolfiella australiana* n. sp. and *I. bassiana* n. sp., both from marine shelly bottoms in Bass Strait, Australia. The authors are skeptical as to the current division of *Ingolfiella* in subgenera, but support the placement of the family within the Gammaroidea and not as a special suborder of its own).
- LOWRY, J.K. & H.E. STODDART, 1989. The scopelocheirid genus *Aroui* (Crustacea: Amphipoda: Lysianassoidea) with notes on the association between scopelocheirid amphipods, cassid gastropods, and spatangoid echinoids. ____ Rec. austr. Mus. 41, 111-120. (*Aroui setosus* is fully redescribed and a second species, *A. hamatopodus* n. sp., described from NSW-waters. The authors state that scopelocheirids are specialized predators on spatangid echinoids, and speculate on the possible role of cassid gastropods in making the spatangids available to the amphipods).
- LOWRY, J.K. & H.E. STODDART, 1989. *Shoemakerella* Pirlot, 1936 (Crustacea, Amphipoda): proposed designation of *Lysianax cubensis* Stebbing, 1897, as the type species. ____ Bull. zool. Nomencl. 46, 236-238. (This proposal, i.e. to change the type species of *Shoemakerella* from *Lysianassa nasuta* to *L. cubensis* (two spp. Pirlot thought to be synonymous), will formalize the generally accepted framework of the genus *Shoemakerella*. This becomes important, as the authors consider *Shoemakerella* to be a valid genus).
- MACKIE, G.L., 1989. Tolerance of five benthic invertebrates to hydrogen ions and metals (Cd, Pb, Al). ____ Arch. environm. Contam. Toxicol. 18, 215-223. (*Hyalella azteca* most sensitive).
- MAGALHAES, C., J.C.O. MALTA, B. ROBERTSON & A. VARELLA & A. VARELLA, 1988. A catalogue of type specimens of Crustacea in the invertebrate collection of the Instituto Nacional de Pesquisas da Amazonia, Manaus, Brazil, up to January, 1988. ____ Amazoniana 10, 267-282. (Not seen, Amph ?).
- MATTHIESEN, P. & J.E. THAIN, 1989. A method for studying the impact of polluted marine sediments on intertidal colonising organisms; tests with diesel-based drilling mud and tributyltin antifouling paint. ____ Hydrobiologia 188/189, 477-485. (i.a. *Urothoe poseidonis*).
- MATTSON, S. & T. CEDHAGEN, 1989. Aspects of the behaviour and ecology of *Dyopodes monacanthus* (Metzger) and *D. porrectus* Bate, with comparative notes on *Dulichia tuberculata* Boeck (Crustacea, Amphipoda: Podoceridae). ____ J. exp. mar. Biol. Ecol. 127, 253-272. (Most interesting paper!).
- McCLATCHIE, S., D. HUTCHINSON & K. NORDIN, 1989. Aggregation of avian predators and zooplankton prey in Otago shelf waters, New Zealand. ____ J. Plankton Res. 11, 361-374. (i.a. *Parathemisto* spp.).
- McDERMOTT, J.J., 1987. Laboratory observations on the feeding biology of nemertean from northwest Wales. ____ Proc. Pennsylvania Acad. Sci. 61, 38-40.
- McDERMOTT, J.J., 1988. The role of hoplonemertean in the ecology of seagrass communities. ____ Hydrobiologia 156, 1-11.
- McDERMOTT, J.J. & R.L. SNYDER, 1988. Food and feeding behaviour of the hoplonemertean *Oerstedtia dorsalis*. ____ Hydrobiologia 156, 47-51.
- McGRATH, D. & A.A. MYERS, 1989. The drift amphipod *Hyale grimaldii* in Irish and British waters. ____ J. mar. biol. Ass. UK 69, 913-918.
- MENIOUI, M. & S. RUFFO, 1988. Considerations sur quelque amphipodes intéressants trouvés sur les côtes du Maroc. ____ Atti Soc. Tosc. Sci. Nst., Mem B 95, 161-175. (Contains a number of records of zoogeographical interest both from the Mediterranean and Atlantic coasts of Morocco. *Elasmopus vachoni* and *Leucothoe procer* are illustrated).
- MESCHENMOSER, M., 1989. Ultrastructure of the embryonic dorsal organ of *Orchestia cavimana* (Crustacea: Amphipoda); with a note on localization of chloride and on the change in calcium deposition before the embryonic molt. ____ Tissue & Cell 21, 431-442.
- MEUSY, J.-J. & G.G. PAYEN, 1988. Female reproduction in malacostracan Crustacea. ____ Zool. Sci. 5, 217-265. (A review. Not seen).
- MEYER-ROCHOW, B. & T. SUZUKI, 1986. A study of visual pigments in the two Antarctic crustaceans *Orchomene plebs* (Amphipoda) and *Glyptonotus antarcticus* (Isopoda). ____ Comp. Biochem. Physiol. B 83, 75-78.
- MICHAUT, P. & F. GRAF, 1988. (Dynamics of the posterior caeca epithelium of *Orchestia* (Crustacea, Amphipoda): 1. Growth). ____ Ann. Sci. nat. Zool. Biol. anim. 9, 67-76. (In French, not seen).

- MOORE, P.G., 1989. Three new amphipod species related to Hippomedon (Crustacea; Amphipoda; Lysianassoidea) from Tasmanian inshore waters. ____ J. nat. Hist. **23**, 1443-1460. (Deals with Hippomedon rodericki n. sp., ?H. adentatus n. sp. and H. dentatus n. sp., all from Burnie, Tasmania. With a key to southern hemisphere Hippomedon s. lat).
- MORRIS, R.J., A.P.M. LOCKWOOD, D. DYBALL & S.R.L. BOLT, 1987. Changes in the fatty acid composition of the gill phospholipids in Gammarus duebeni during moult: Evidence for reduced permeability of the gill membrane. ____ Comp Biochem. Physiol. B **88**, 257-260.
- MORRITT, D., 1989. Ionic regulation in littoral and terrestrial amphipods (Crustacea: Amphipoda: Talitridae). ____ J. exp. mar. Biol. Ecol. **123**, 53-67.
- MUKAI, H. & M. TAKEDA, 1987. A giant amphipod Crustacea from the Miocene Morozaki Group in the Chita peninsula, Central Japan. ____ Bull. natn. Sci. Mus. C (Geol. Paleontol.) **13** (1), 35-39. (Not seen).
- MULLER, H-G., 1987. Zur Biologie der Gespensterkrebse. ____ Mikrokosmos **76**, 180-185. (Not seen. Caprellidae?)
- MURDOCH, R.C., 1989. The effect of a headland eddy on surface macro-zooplankton assemblages North of Otago peninsula, New Zealand. ____ Est. coast. Shelf Sci. **29**, 361-383. (i.a. Parathemisto and Cylopus spp.).
- MUSKO, I.B., 1989. Amphipoda (Crustacea) in the littoral zone of Lake Balaton (Hungary). Qualitative and quantitative studies. ____ Int. Rev. ges. Hydrobiol. **74**, 195-205. (Deals with Corophium curvispinum, Dikerogammarus haemobaphes and D. villosus).
- MYERS, A.A., 1989. Amphipoda from the South Pacific: The Society Islands. ____ Rec. austr. Mus. **41**, 63-82. (Not seen. New spp. Elasmopus integer, Maera lindsae, M. mooreana, Cheiriphotis rotui, Jassa socia and Polynesocetes kekeae n. gen., n. sp.).
- MYERS, A.A., D. McGRATH & P. CUNNINGHAM, 1989. A presumed male of the parthenogenetic amphipod Corophium bonnellii (Milne-Edwards). ____ J. mar. biol. Ass. UK **69**, 319-321.
- NALEPA, T.F., 1989. Estimates of macroinvertebrate biomass in Lake Michigan. ____ J. Great Lakes Res. **15**, 437-443. (Pontoporeia hoyi dominant).
- NAPOLITANO, G.E. & R.G. ACKMAN, 1989. Lipid and hydrocarbons in Corophium volutator from Minas Basin, Nova Scotia, Canada. ____ Mar. Biol. **100**, 333-338.
- NAYLOR, C., L. MALTBY & P. CALOW, 1989. Scope for growth in Gammarus pulex, a freshwater benthic detritivore. ____ Hydrobiologia **188/189**, 517-523.
- NEBEKER, A.V., W.L. GRIFFIS, C.M. WISE, E. HOPKINS & J.A. BARBITTA, 1989. Survival, reproduction and bioconcentration in invertebrates and fish exposed to hexachlorobenzene. ____ Environm. Toxicol. Chem. **8**, 601-611.
- NEBEKER, A.V. & C.E. MILLER, 1988. Use of the amphipod crustacean Hyalella azteca in freshwater and estuarine sediment toxicity tests. ____ Environm. Toxicol. Chem. **7**, 1027-1033.
- NEBEKER, A.V., G.S. SCHUYTEMA, W.L. GRIFFIS, J.A. BARBITTA & L.A. CAREY, 1989. Effect of sediment organic carbon on survival of Hyalella azteca exposed to DDT and endrin. ____ Environm. Toxicol. Chem. **8**, 705-718.
- NICOLAIDOU, A. & M. KARAKIRI, 1989. The distribution of Amphipoda in a brackish-water lagoon in Greece. ____ Mar. Ecol. **10**, 131-140.
- NYLUND, A., S. ÖKLAND & A. TJÖNNELAND, 1987. The crustacean heart ultrastructure and its bearing upon the position of the isopods in eumalacostracan phylogeny. ____ Zool. Scripta **16**, 235-241.
- OLSEN, A., 1989. Physiological effects of cyanide and mercury in the gammarid Marinogammarus marinus. ____ Pp. 213-218 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept BECTOS- progr. 1985-88. Univ. of Trondheim, Norway.
- OLSEN, A., T. AUNAAS, J.F. BÖRSETH & J-P. DENSTAD, 1989. Methods and procedures used in the present studies on gammarids, blue mussels, and fish eggs and alevins. ____ Pp. 19-36 in K.E. Zachariassen (ed.). Biological effects of chemical treatment of oilspills at sea. Rept. BECTOS-progr. 1985-88. Univ. of Trondheim, Norway.
- PANOV, V.E., 1986. Growth and production of Gammarus lacustris in the Neva inlet. ____ Hidrobiol. Zh. **22** (5), 37-42. (In Russian, not seen).
- PARDI, L., 1987. Wenn die Strandflohkrebse den Himmel anschauen: Ein Beitrag zur tierischen Orientierung an der Meeres-Landesgrenze. ____ Pp. 7-54 in M. LINDAUER (ed.). Information processing in animals, vol. 4. Gustav Fischer Verlag, Stuttgart.
- PARDI, L. & A. ERCOLINI, 1986. Zonal recovery mechanisms in talitrid crustaceans. ____ Bull. Zool. **53**, 139-160.
- PATARNELLO, T., P.M. BISOL & B. BATTAGLIA, 1989. Studies in differential fitness of PGI genotypes with regard to temperature in Gammarus insensibilis (Crustacea: Amphipoda). ____ Mar. Biol. **102**, 355-360.
- PLATVOET, D., M. SCHEEPMACKER & S. PINKSTER, 1989. The position of two introduced amphipod crustaceans, Gammarus tigrinus and Crangonyx pseudogracilis in the Netherlands during the period 1987-88. ____ Bull. zool. Mus. Univ. A'dam **11**, 197-200.
- POOVACHIRANON, S., K. BOTO & N. DUKE, 1986. Food preference studies and ingestion rate measurements of the mangrove amphipod Parhyale hawaiiensis (Dana). ____ J. exp. mar. Biol. Ecol. **98**, 129-140.

- POSTAN, T.M., R.M. BEAN, D.R. KALKWARF, B.L. THOMAS, M.L. CLARK & B.W. KILLAND, 1988. Photooxidation products of smoke generator fuel (SGF), No. 2 fog oil and toxicity to Hyalella azteca. ____ Environm. Toxicol. Chem. 7, 753-762.
- POULTON, M.J. & D.J. THOMPSON, 1987. The effects of the acanthocephalan parasite Pomphorhynchus laevis on mate choice in Gammarus pulex. ____ Anim. Behav. 35, 1577-1579.
- PRESING, M., 1989. Data to the toxic effect of K-Othrine on crustaceans. ____ Arch. Hydrobiol. 114, 621-629. (i.a. Gammarus roeseli).
- PRESTON, A. & P.G. MOORE, 1989. Seasonal cycles of abundance of the flora and fauna associated with Cladophora albida (Huds.) Kütz. in rockpools. ____ J. nat. Hist. 23, 983-1002.
- QUIGLEY, M.A. & G.A. LANG, 1989. Measurement of amphipod body length using a digitizer. ____ Hydrobiologia 171, 255-258.
- RAFFAELLI, D., A. CONACHER, H. McLACHLAN & C. EMES, 1989. The role of epibenthic crustacean predators in an estuarine food web. ____ Est. coast. Shelf Sci. 28, 149-160.
- RAFFAELLI, D., S. HULL & H. MILNE, 1989. Long-term changes in nutrients, weed mats and shorebirds in an estuarine system. ____ Cah. Biol. mar. 30, 259-270. (i.a. Corophium volutator).
- RAINBOW, P.S., 1989. Copper, cadmium and zinc concentrations in oceanic amphipod and euphausiid crustaceans, as a source of heavy metals to pelagic seabirds. ____ Mar. Biol. 103. (i.a. Themisto spp.).
- RAUSCHERT, M., 1989. Atylopsis fragilis n. sp. (Crustacea, Amphipoda, Gammaridea, Eusiridae) aus dem Sublitoral von King George (Süd-Shetland-Inseln). ____ Mitt. zool. Mus. Berlin 65, 127-138.
- READ, G.H.L. & A.K. WHITFIELD, 1989. The response of Grandidierella lignorum (Barnard) (Crustacea: Amphipoda) to episodic flooding in three eastern Cape estuaries (S. Africa). ____ S. Afr. J. Zool. 24, 99-105.
- REDMOND, M.S. & K.J. SCOTT, 1989. Amphipod predation by the infaunal polychaete, Nephtys incisa. ____ Estuaries 12, 205-207. (Ampelisca abdita and Microdeutopus gryllotalpe).
- REHAGE, H.D., 1987. Zum weiteren Vordringen von Orchestia cavimana Heller, 1865 (Crustacea, Talitridae) in Westfalen. ____ Natur Heimat 47, 41-44. (Not seen).
- ROBERTSON, M.R., S.J. HALL & A. ELEFThERIOU, 1989. Environmental correlates with amphipod distribution in a Scottish sea loch. ____ Cah. Biol. mar. 30, 243-258.
- RODGERS, K.A. & R. OLERÖD, 1988. A catalog of zoological specimens collected from Tuvalu (Ellice Islands) by Sixten Bock, 1917. ____ Pacif. Sci. 42, 300-306. (Amph. p. 302. The holotype of Paranamixis bocki could not be found).
- ROUCH, R., 1988. Spatial distribution of crustaceans in the hyporheic habitat of a Pyrenean brook (France). Annls. Limnol. 24, 213-234. (In French, not seen).
- ROUCH, R. & D.L. DANIELOPOL, 1987. The origin of the subterranean freshwater fauna, between the refugium paradigm and the model of active colonization. ____ Stygologia 2, 345-372. (Not seen).
- RUFFO, S. (ed.), 1989. The Amphipoda of the Mediterranean. Part. 2. Gammaridea (Haustoriidae to Lysianassidae). ____ Mem. Inst. océanogr., Monaco 13, 365-576. (This second part of this monumental monograph deals with the families Haustoriidae (3 genera, 17 species), Isaeidae (6-20), Ischyroceridae (3-4), Lafystiidae (1-1), Leucothoidae (1-10), Liljeborgiidae (2-5) and Lysianassidae (30-56). Of the 113 spp. 48 are new to the Mediterranean; all new species have been described in advance in separate papers. The contributing authors are D. Bellan-Santini (Haust.), G. Krapp-Schickel (Leuc., Lilj.), A.A. Myers (Is., Isch.) and S. Ruffo (Laf., Lys., with G. Diviacco).
- RUFFO, S. & A. VIGNA TAGLIANTI, 1988. Gammaropisa arganoi, new genus new species from the phreatic waters of southern Anatolia (Crustacea, Amphipoda, Gammaridae sensu lato). ____ Boll. Mus. Civ. Stor. Nat. Verona 14 (1987), 241-254. (The new genus is intermediate between 'gammarids' and 'melitids'. It was found in wells near Antalya, S. Turkey).
- RUFFO, S. & A. VIGNA TAGLIANTI, 1989. Description of a new cavernicolous Ingolfiella species from Sardinia, with remarks on the systematics of the genus (Crustacea, Amphipoda, Ingolfiellidae). ____ Ann. Mus. Civ. Stor. Nat. Genova 87, 237-261. (Ingolfiella cottarellii n. sp. from a freshwater cave at sea level on an islet off Sardinia. The ocular lobes of I. ruffoi are illustrated. The authors give a review of the taxonomy of Ingolfiella, which they provisionally divide into 7 subgenera: Ingolfiella (abyssi, britannica, atlantisi), Hanseniella (littoralis, berrisfordi, ischitana), Tethydiella n. subgen. (fuscina (type), xariffae, kapuri, grandispina, quadridentata, longipes), Antilleella n. subgen. (tabularis (type), putealis, fontinalis, margaritae, similis), Gevgeliella (petkovskii, vandeli), Balcanella (possibly not monophyletic) (acherontis, macedonica, manni, uspallatae), and Tyrrenidiella n. subgen. (cottarellii (type), catalanensis, thibaudi). A key to the subgenera is provided. This taxonomy differs in many particulars from that of Stock, and a number of spp. is transferred to different subgenera. The subgenus Trianguliella is provisionally synonymized with Balcanella, but may well prove distinct.
- RUSSO, A.R., 1989. Fluctuations of epiphytal gammaridean amphipods and their seaweed hosts on a Hawaiian algal turf. ____ Crustaceana 57, 25-37.
- SABATES, A., J.M. GILL & F. PAGES, 1989. Relationship between zooplankton distribution, geographic characteristics and hydrographic patterns off the Catalan coast (western Mediterranean). ____ Mar. Biol. 103, 153-160.

- SAINTE-MARIE, B., J.A. PERCY & J.R. SHEA, 1989. A comparison of meal size and feeding rate of the lysianassid amphipods *Anonyx nugax*, *Onisimus littoralis*, and *Orchomenella pinguis*. ____ Mar. Biol. **102**, 361-368.
- SANTOS, J. dos & S. FALK-PETERSEN, 1989. Feeding ecology of cod (*Gadus morhua* L.) in Balsfjord and Ullsfjord, northern Norway, 1982-1983. ____ J. Cons. int. Explor. Mer **45**, 190-199.
- SCAPINI, F., 1986. Inheritance of direction finding in sandhoppers. ____ Pp. 111-119 in G. BEUGNON (ed.). Orientation in space. Editions Privat, Toulouse.
- SCAPINI, F., 1987. Erblisches und erlerntes in der Orientierung der Strandfloh. ____ Pp. 55-74 in M. LINDAUER (ed.). Information processing in animals, vol. 4. Gustav Fischer Verlag, Stuttgart.
- SCAPINI, F., 1988. Heredity and learning in animal orientation. ____ Monit. zool. ital. **22**, 203-234. (Talitrid work on pp. 210-220).
- SCAPINI, F., M. BUIATTI & O. OTTAVIANO, 1988. Phenotypic plasticity in sun orientation of sandhoppers. ____ J. comp. Physiol. A **162**, 739-749.
- SCAPINI, F., A. UGOLINI & L. PARDI, 1988. Aspects of direction finding inheritance in natural populations of littoral sandhoppers (*Talitrus saltator*). ____ Pp. 93-103 in G. CHELAZZI & M. VANNINI (eds.). Behavioural adaptation to intertidal life. Plenum Publ. Corp.
- SCHUYTEMA, G.S., D.F. KRAWCZYK, W.L. GRIFFIS, A.V. NEBEKER, M.L. ROBIDEAUX, B.J. BROWNAWELL & J.C. WESTALL, 1988. Comparative uptake of hexachlorobenzene by fathead minnows, amphipods and oligochaete worms from water and sediment. ____ Environm. Toxicol. Chem. **7**, 1035-1045. (*Hyalella azteca*).
- SEDBERRY, G.R., 1988. Food and feeding of black sea bass, *Centropomus striata*, in live bottom habitat in the South Atlantic bight. ____ J. Elisha Mitchell scient. Soc. **104**, 35-50. (Amph. very important; caprellids and *Ericthonius* selected for).
- SELLEM, E., F. GRAF & J.-C. MEYRAN, 1989. Some effects of salmon calcitonin on calcium metabolism in the crustacean *Orchestia* during the molt cycle. ____ J. exp. Zool. **249**, 177-181.
- SERGEEV, V.N., S.M. CLARKE & S.A. SHEPHERD, 1988. Motile macroepifauna of the seagrasses, *Amphibolis* and *Posidonia*, and unvegetated sandy substrates in Holdfast Bay, South Australia. ____ Trans. R. Soc. S. Austr. **112**, 97-108.
- SHAW, D.P., 1989. Redescription of *Clarencia chelata* K.H. Barnard, 1932 (Amphipoda, Eusiroidea) ____ Crustaceana **57**, 201-207.
- SHEADER, M. & A. SHEADER, 1987. The distribution of the lagoonal amphipod, *Gammarus insensibilis* Stock, in England. ____ Porcupine Newsl. **3** (9), 220-223. (Not seen).
- SKADSHEIM, A., 1989. Regional variation in amphipod life history: effects of temperature and salinity on breeding. ____ J. exp. mar. Biol. Ecol. **127**, 25-42.
- SOKOLOV, V.E. & T.G. EVGEN'EVA, 1988. (Response of the skin epidermis of the gray whale *Eschrichtius gibbosus* to parasitizing Crustacea). ____ Dokl. Akad. Nauk SSSR **303**, 484-487. (In Russian, not seen. Deals with i.a. *Cyamus scammoni*).
- SPICER, J.I. & A.C. TAYLOR, 1989. Oxygen-binding by the blood of the landhopper, *Arcitalitrus dorrieni* (Hunt) (Crustacea: Amphipoda). ____ Comp. Biochem. Physiol. A **92**, 421-422.
- STATZNER, B. & T.F. HOLM, 1989. Morphological adaptation of shape to flow: Microcurrents around lotic macroinvertebrates with known Reynolds numbers at quasi-natural flow conditions. ____ Oecologia **78**, 145-157. (i.a. *Gammarus*).
- STEELE, D.H., 1989. The genus *Anonyx* (Crustacea, Amphipoda) in the North Pacific and Arctic oceans: *Anonyx compactus* group. ____ Can. J. Zool. **67**, 1945-1954. (Deals with *Anonyx compactus*, *A. attenuatus* n. sp. (Kyska Harbor, Alaska), *A. stappersi* n. sp. (= *Chironesimus debruyni* s. Stappers, holotype from unknown loc., distr. from Kara to Norwegian Seas), *A. oculatus*, *A. stebbingi* n. sp. (= *A. ampulloides* s. Stebbing, off Japan), *A. ochoticus* and *A. derjugini* (= *A. ampulloides* s. Gurjanova 1951)).
- STEELE, V.J. & P.E. OSHEL, 1989. Ultrastructure of the attachment cells of the organ of Bellonci in *Gammarus setosus* (Crustacea, Amphipoda). ____ J. Morphol. **200**, 93-119.
- STEPHENSON, M. & G.L. MACKIE, 1988. Multivariate analysis of correlations between environmental parameters and cadmium concentrations in *Hyalella azteca* (Crustacea: Amphipoda) from central Ontario lakes. ____ Can. J. Fish. aq. Sci. **45**, 1705-1710.
- STOCK, J.H., 1989. A new genus and species of Talitridae (Amphipoda) from a cave in Terceira, Azores. ____ J. nat. Hist. **23**, 1109-1118. (*Macarorchestia martinii* n. gen., n. sp.).
- STOCK, J.H. & G.A. BOXSHALL, 1989. Comparison between the landhoppers (Amphipoda: Talitridae) of the genus *Orchestia* from Tenerife (Canary Islands) and the Azores. ____ Beaufortia **39**, 45-54. (*O. chevreuxi* is confined to the Azores, while the Tenerife populations belong to a new species, *O. guanacha* n. sp.).
- STORCH, V. & BURKHARDT, 1986. Di Reaktion der Mitteldarmdrüse des Süsswasserstrandfloh *Orchestia cavimana* auf verschiedener Nahrung. ____ Carolinea **44**, 149-152.

- SWARTZ, R.C., P.F. KEMP, D.W. SCHULTS & J.O. LAMBERTSON, 1988. Effects of mixtures of sediment contaminants on the marine infaunal amphipod, Rhepoxynius abronius. ____ Environm. Toxicol. Chem. 7, 1013-1020.
- THEODORIDES, J., 1989. Parasitology of marine plankton. ____ Adv. mar. Biol. 25, 117-177. (Amph. pp. 154-156).
- THURSTON, M.H., 1989. A new genus and species of fossorial amphipod from the Falkland Islands (Crustacea, Amphipoda, Phoxocephalopsidae), with notes on Phoxocephalopsis. ____ J. nat. Hist. 23, 299-310. (Eophoxocephalopsis rhachianensis n. gen., n. sp. from intertidal sandy beach in E. Falklands. Phoxocephalopsis deceptionis is transferred to Eophoxocephalopsis. Haustoriella psammophila is shown to be a synonym of Phoxocephalopsis zimneri, but P. zimneri s. Ruffo 1956 represents an undescribed species. A key to the Phoxocephalopsidae is provided).
- THURSTON, M.H., 1989. A new species of Valettia (Crustacea: Amphipoda) and the relationship of the Valettidae to the Lysianassoidea. ____ J. nat. Hist. 23, 1093-1108. (Valettia hystrix n. sp. The Valettidae are re-established, close to, but outside the Lysianassoidea. V. hystrix was found in the guts of abyssal holothuroids).
- TJÖNNELAND, A., S. ÖKLAND & A. NYLUND, 1987. Evolutionary aspects of the amphipod heart. ____ Zool. Scripta 16, 167-175.
- UGOLINI, A. & T. MACCHI, 1988. Learned component in the solar orientation of Talitrus saltator Montagu (Amphipoda: Talitridae). ____ J. exp. mar. Biol. Ecol. 121, 79-87.
- UGOLINI, A., F. SCAPINI, G. BEUGNON & L. PARDI, 1988. Learning in zonal orientation of sandhoppers. ____ Pp. 105- ____ in CHELAZZI & M. VANNINI (eds). Behavioral adaptation to intertidal life. Plenum Publ. Corp.
- ULIAN, G.B. & E.G. MENDES, 1987. Preferences of a terrestrial amphipod, Talitrus (Talitroides) pacificus Hurley, 1955, towards some environmental factors. ____ Rev. Bras. Biol. 47, 247-256.
- VÄINÖLÄ, R. & S.-L. VARVIO, 1989. Molecular divergence and evolutionary relationships in Pontoporeia (Crustacea: Amphipoda). ____ Can. J. Fish. aq. Sci. 46, 1705-1613.
- VALDECASAS, A.G. & A. BALTANAS, 1989. A note on the use of Angelier's fluid for freshwater invertebrates. ____ Arch. Hydrobiol. 115, 313-316. ('Close to an ideal fixative').
- VAWTER, A.T., D.W. FONG & D.C. CULVER, 1987. Negative phototaxis in surface and cave populations of the amphipod Gammarus minus. ____ Stygologia 3, 83-88.
- VIGNA-TAGLIANTI, A., 1988. A cave amphipod of the hadziid group from Turkey (Crustacea, Amphipoda, Melitidae). ____ Boll. Mus. Civ. Stor. Nat. Verona 14 (1987), 439-452. (Parhadzia sbordonii n. gen., n. sp. from S. Anatolia, a'hadziid'. The author opines that the hadziids are oversplit at the generic level, and provisionally only recognizes Hadzia, Metaniphargus, Bahadzia and the new Parhadzia).
- VONK, R., 1988. The Niphargus kochianus - group (Crustacea, Amphipoda) in the Aquitanian Basin, south-western France, and a supposed case of introgressive hybridization. ____ Stygologia 4, 177-191. (Deals with N. k. kochianus, N. pachypus, and their possible hybrid. First description of N. pachypus male).
- VONK, R., 1988. Amsterdam expeditions to the West Indian Islands: Report 55, Psammomelita uncinata new genus new species (Crustacea, Amphipoda, Melitidae) from infralittoral and interstices on Curacao. ____ Stygologia 4, 166-176. (Not seen).
- WÄGELE, J.W., 1989. On the influence of fishes on the evolution of benthic crustaceans. ____ Z. zool. Syst. Evol. f. 27, 297-309.
- WARD, P., 1989. The distribution of zooplankton in an Antarctic fjord at South Georgia during summer and winter. ____ Antarct. Sci. 1, 141-150.
- WATLING, L., 1989. A classification system for crustacean setae bound on the homology concept. ____ Pp. 15-26 in B.E. FELGENHAUER, L. WATLING & A.B. THISTLE (eds). Functional morphology of feeding and grooming in Crustacea. A.A. Balkema, Rotterdam.
- WATLING, L. & M.H. THURSTON, 1989. Antarctica as an evolutionary incubator: evidence from the cladistic biogeography of the amphipod family Iphimedilidae. ____ Pp. 297-313 in J.A. CROME (eds). Origins and evolution of Antarctic biota. Geol. Soc. spec. Publ. 47.
- WEIGMANN-HAAS, R., 1989. Zur Taxonomie und Verbreitung der Gattung Hyperiella Bovallius 1887 im antarktischen Teil des Atlantik. ____ Senckenberg. biol. 69, 177-191.
- WESAWSKI, J.M. & W. KULINSKI, 1989. Notes on fishes in Hornsund fjord area (Spitsbergen). ____ Polish polar Res. 10, 241- 250. (With data on diets of several spp., i.a. with many amph.).
- WESAWSKI, J.M., M. ZAJACZKOWSKI, S. KWASNIEWSKI, J. JEZIERSKI & W. MOSKAL, 1988. Seasonality in an Arctic fjord ecosystem: Hornsund, Spitsbergen. ____ Polar Res. 6, 185-190.
- WILLIAMS, D.D. & K.A. MOORE, 1989. Environmental complexity and the drifting behaviour of a running water amphipod. ____ Can. J. Fish. aq. Sci. 46, 1520-1530. (Gammarus pseudolimnaeus).
- WILSON, W.H., 1988. Shifting zones in a Bay of Fundy soft-sediment community: patterns and processes. ____ Ophelia 29, 227-245. (Corophium volutator dominant in lower zone).

- WILSON, W.H., 1989. Predation and the mediation of intraspecific competition in an infaunal community in the Bay of Fundy. ____ J. exp. mar. Biol. Ecol. 132, 221-245.
- YAMATO, S., 1988. Two species of the genus *Melita* (Crustacea: Amphipoda) from brackish waters in Japan. ____ Publ. Seto mar. Biol. Lab. 33, 79-95. (*Melita setiflagella* n. sp. and *M. shimizui*).
- ZACHARIASSEN, K.E. (ed.), 1989. Biological effects of chemical treatment of oilspills at sea. ____ Rept. BECTOS progr. 1985-88, 330 pp. Univ. of Trondheim, Trondheim, Norway. (Contributions listed separately).

BIBLIOGRAPHY

January 1990

- AARSET, A.V. & F.V. WILLUMSEN, 1985. Hydraulic based sampling equipment for under-ice fauna. ____ Polar Res. 3, 253-255.
- ABELE, L.G. & B.E. FELGENHAUER, 1985. Observation on the ecology and feeding behavior of the anchialine shrimp *Procaris ascensionis*. ____ J. crust. Biol. 5, 15-24. (A predator on *Melita*).
- ADAMS, J. & P.J. GREENWOOD, 1985. Environmental constraints on mate choice in *Gammarus pulex* (Amphipoda). ____ Crustaceana 50, 45-52.
- ADAMS, J. & P.J. GREENWOOD, 1987. Loading constraints, sexual selection and assortative mating in peracarid Crustacea. ____ J. Zool. 211, 35-46.
- AGNEW, D.J. & M.B. JONES, 1986. Metabolic adaptations of *Gammarus duebeni* (Crustacea, Amphipoda) to hypoxia in a sewage treatment plant. ____ Comp. Biochem. Physiol. A 84, 475-478.
- AGNEW, D.J. & P.G. MOORE, 1986. The feeding ecology of two littoral amphipods (Crustacea), *Echinogammarus pirloti* (Sexton & Spooner) and *E. obtusatus* (Dahl). ____ J. exp. mar. Biol. Ecol. 103, 203-215.
- AGNEW, D.J. & A.C. TAYLOR, 1985. The effect of oxygen tension on the physiology and distribution of *Echinogammarus pirloti* (Sexton and Spooner) and *E. obtusatus* (Dahl) (Crustacea: Amphipoda). ____ J. exp. mar. Biol. Ecol. 87, 169-190.
- AGNEW, D.J. & A.C. TAYLOR, 1986. Effects of oxygen tension, temperature, salinity, and humidity on the survival of two intertidal gammarid amphipods. ____ Mar. Ecol. Progr. Ser. 32, 27-33. (You guessed right: *Echinogammarus pirloti* and *E. obtusatus*).
- AHSANULLAH, M. & G.W. BRAND, 1985. Effects of selenite and seleniferous fly-ash leachate on growth and viability of the marine amphipod *Allorchestes compressus*. ____ Mar. Biol. 89, 245-248.
- AHSANULLAH, M. & A.R. WILLIAMS, 1986. Effect of uranium on growth and reproduction of the marine amphipod *Allorchestes compressus*. Mar. Biol. 93, 459-464.
- AINLEY, D.G., W.R. FRASER, C.W. SULLIVAN, J.C. TORRES, T.L. HOPKINS & W.D. SMITH, 1986. Antarctic mesopelagic micronekton; Evidence from seabirds that pack ice affects community structure. ____ Science (N.Y.) 232, 847-849.
- ALEYEV, YU. G. & S.A. KHVOROV, 1986. (Buoyancy as a function of life forms of hydrobionts). ____ Gidrobiol. Zh. 21(6), 22-26. (In Russian).
- ALI, M.H. & S.D. SALMAN, 1986. The reproductive biology of *Parhyale basrensis* Salman (Crustacea: Amphipoda) in the Shatt al-Arab river. ____ Est. coast. Shelf Sci. 23, 339-352.
- ALIEV, R.A., 1985. (Composition and distribution of makrozoobenthos in lakes Adzhikabul and Nakhalykhchala (Azerbaijan SSR, USSR)). ____ Izv. Akad. Nauk AZ SSR, Ser. Biol. Nauk 1985(4), 70-76. (In Russian, not seen).
- ALLDREDGE, A.L. & J.M. KING, 1985. The distance demersal zooplankton migrate above the benthos: implications for predation. ____ Mar. Biol. (Berl.) 84, 253-260. (Many amphipod data. A study from the Gulf of California).
- ALOUF, N.J., 1986. Biologie de *Gammarus laticoxalis* dans une rivière de Liban. ____ Hydrobiologia 133, 45-57.
- ANDERSSON, K.G., C. BRÖNMARK, J. HERRMANN, B. MALMQUIST, C. OTTO & P. SJÖSTRÖM, 1986. Presence of sculpins (*Cottus gobio*) reduces drift and activity of *Gammarus pulex* (Amphipoda). ____ Hydrobiologia 133, 209-215.
- ANDERSON, R.L. & P. SHUBAT, 1984. Toxicity of flucythrinate to *Gammarus lacustris* (Amphipoda), *Pteronarcys dorsata* (Plecoptera) and *Brachycentrum americanus* (Trichoptera): Importance of exposure duration. ____ Environm. Poll. 35, 353-365.
- ANDRES, H.G., 1985. Die Gammaridea (Crustacea: Amphipoda) der deutschen Antarktis- Expeditionen 1975/76 und 1977/78. 4. Acanthonotozomatidae, Paramphithoidae und Stegocephalidae. ____ Mit. hamb. zool. Mus. Inst. 82, 119-153. (Deals with *Iphimediella margueritei*, *Pariphimedia incisa* n. sp. (Palmer archipelago) (= *P. integricauda* s. Barnard 1932), *Epimeria macrodonta*, *E. similis*, *E. extensa* n. sp. (64° S, 55° W), *E. sp.*, *Epimeriella macronyx* (with discussion of relations between *Epimeriella*, *Eclysis* and *Astyra*), *E. walkeri*, *E. truncata* n. sp. (64° S, 64,5° W) and

- Euandania nonhiata n. sp. (63,5° S, 64,5° W). Lepidepcreum urometacarinatum is a nov. nom. for L. carinatum Andres 1983, non B. & W. 1868).
- ANDRES, H.G., 1986. Atylopsis procerus sp. n. und Cheirimedon solidus sp. n. aus der Weddell See sowie Anmerkungen zu Orchomenella pinguides Walker, 1903 (Crustacea: Amphipoda: Gammaridea). ____ Mitt. hamb. Zool. Mus. Inst. 83, 117-130.
- ANDRES, H.G. & N. LOTT, 1986. Where to place Eclysis similis K.H. Barnard, 1932? Hints at its relationship and remarks on the systematic position of the Astyridae (Crustacea: Amphipoda). ____ Mitt. hamb. zool. Mus. Inst. 83, 131-137. (Eclysis, and also Epimeriella, should be assigned to the Astyridae, an independent family).
- ANONYMOUS, 1985. (The effect of lithium chloride on Pontogammarus robustioides (Grimm)). ____ Trudy Akad. Nauk Litov. SSR C 89, 147. (In Russian, abstract only).
- ANSARI, Z.A., B.S. INGALE & A.H. PARULEKAR, 1984. Macrofauna and meiofauna of two sandy beaches at Mombasa, Kenya. ---- Indian J. mar. Sci. 13, 187-189. (Not seen).
- ARIMOTO, I., 1980. Occurrence of caprellids (Amphipoda, Caprellidae) from the Gelidium region off Misaki-cho, Ehime prefecture, Japan. ____ Bull. biogeogr. Soc. Japan 35, 33-38. (Not seen).
- ARIMOTO, I., 1982. Three species of caprellid amphipods from Ogasowara Islands, with description of a new species. ____ Proc. jap. Soc. syst. Zool. 22, 21-23. (Not seen. Caprella temperativa n. sp.).
- ARNAUD, P.M., K. JAŹDŹEWSKI, P. PRESLER & J. SICINSKI, 1986. Preliminary survey of benthic invertebrates collected by Polish Antarctic expeditions in Admiralty Bay (King George Island, South Shetland Islands, Antarctica). ____ Pol. Res. Z, 7-24. (Amphipods pp. 16-17, i.a. Synopiidae n. gen. n. sp. Eight spp. are new to the South Shetlands area).
- ATKINS, S.M., A.M. JONES & J.A. SIMPSON, 1985. The fauna of sandy beaches in Orkney: a review. ____ Proc. R. Soc. Edinb. 87 B, 27-45. (Amph. listed on p. 32).
- AZUMA, M. & Y. HANO, 1985. (The bottom fauna communities in Shijiki Bay, Hirado Island. 2. Habitat analysis based on gammarid-sediment relations). ____ Benthos Res. 28, 1-11. (In Japanese, not seen).
- BACHELET, G., 1985. Influence de la maille de tamisage sur les estimations d'abondance des stades juveniles du macrobenthos marin. ____ C-r. hebd. Acad. Sci. Paris 301, 795-798.
- BAKUS, G.J., N.M. TARGETT & B. SCHULTE, 1986. Chemical ecology of marine organisms: an overview. ____ J. chem. Ecol. 12, 951-987.
- BALLY, R., 1986. A bibliography of sandy beaches and sandy beach organisms on the African continent. ____ S. Afr. natn. scient. Program. Rept 126, 1-179.
- BÄRLOCHER, F. & C. PORTER, 1986. Digestive enzymes and feeding strategies of three stream invertebrates. ____ J. North Am. benthol. So. 5, 58-66. (Not seen. Deals with i.a. Gammarus tigrinus).
- BARNARD, J.L. & J.B.R. AGARD, 1986. A new species of Ampelisca (Crustacea, Amphipoda) from Trinidad. ____ Bull. mar. Sci. 39, 630-636. (A. parva n. sp., a rostrate Ampelisca).
- BARNARD, J.L. & J. CLARK, 1985. A new sea-cave amphipod from Bermuda (Dulichidae). ____ Proc. biol. Soc. Wash. 98, 1048-1053. (Podobothrus bermudensis n. gen., n. sp., an aberrant dulichiid with very long antennae).
- BARNARD, J.L. & C.L. INGRAM, 1986. The supergiant amphipod Alicella gigantea, Chevreux from the North Pacific gyre. ____ J. crust. Biol. 6, 825-839. (Maximum size 340 mm!).
- BARNARD, J.L. & J.D. THOMAS, 1987. New species of Neomegamphopus from tropical America (Crustacea: marine Amphipoda). ____ Proc. biol. Soc. Wash. 100, 147-163. (With an extensive and critical discussion of the concept Neomegamphopidae, as defined by Myers. The authors provisionally retain Aoridae, Isaeidae and Neomegamphopidae sub Isaeidae s.l. A key to Neomegamphopus ♂♂ is given. The following species are treated: N. roosevelti, N. hiatus n. sp. (Venezuela), N. kalanii n. sp. (Fort Pierce, Florida), N. pachiatius n. sp. (Pacific Panama), N. heardi n. sp. (Pacific Panama), and N. sp. C. (Venezuela)).
- BARR, T. C. & J.R. HOLSINGER, 1985. Speciation in cave faunas. ____ Ann. Rev. Ecol. Syst. 16, 313-337.
- BECHLER, D.L., 1985. Structure and foraging behavior in hypogean crustacean assemblages. ____ Hydrobiologia 127, 203-211. (A study on Gammarus troglophilus, Bactrurus brachycaudus and the isopod Caecidotea stygia).
- BECHLER, D.L. & A.G. FERNANDEZ, 1981. Preliminary observations on foraging behavior in a hypogean crustacean community. Proc. 8. int. Congr. Speleol., Bowling Green, Ky, USA, 66-67. (Not seen).
- BEKMAN, M. Y., 1984. (Deep water amphipod fauna). ____ Pp. 114-123 in A.A. LINEVICH (ed.) (Systematics and evolution of invertebrates from Bajkal). Nauka, Novosibirsk. (In Russian, not seen. New taxa?).

- BELLAN-SANTINI, D., 1985. Etude de la faune profonde de Méditerranée: les amphipodes des trois campagnes Polymède 1, Polymède 2 et Biomède. ____ Rapp. Comm. int. Mer Médit. 29, 333-334.
- BELLAN-SANTINI, D. & J.C. DAUVIN, 1986. Ampelisca remora (Amphipoda): nouvelle espèce des côtes de Galice (Atlantique nord-est). ____ Crustaceana 51, 38-48. (This species has a unique sucker-like dorsal growth on uros. 1, the function of which is unknown).
- BELLAN-SANTINI, D. & J.C. MARQUES, 1984. Contribution à l'étude des amphipodes des côtes du Portugal. ____ Cienc. biol. 5, 131-149. (Not seen).
- BELLAN-SANTINI, D., J. PICARD & M.L. ROMAN, 1984. Contribution à l'étude des peuplements des invertébrés des milieux extrêmes. 2. Distribution des Crustacés de la macrofaune des plages du delta du Rhône. ____ Ecol. médit. 10 (3-4), 1-7.
- BELLAN-SANTINI, D. & S. RUFFO, 1986. Urothoe intermedia espèce nouvelle recoltée dans le Canal du Suez (Crustacea Amphipoda, Haustoriidae s.l.) ____ Boll. Mus. Civ. Stor. Nat. Verona 12 (1985), 85-95.
- BELLAN-SANTINI, D. & S. RUFFO, 1986. Une nouvelle espèce de Talorchestia des côtes du Sardaigne (Crustacea, Amphipoda, Talitridae). ____ Bol. Mus. Civ. Stor. Nat. Verona 12 (1985), 405-410. (T. pelecaniformis n. sp. from Sardinia).
- BELLAN-SANTINI, D. & W. VADER, 1988. Nouvelles espèces de Bathyporeio en Méditerranée (Crustacea, Amphipoda). ____ Boll. Mus. civ. St. nat. Verona 14 (1987), 229-240. (Deals with Bathyporeia pseudopelagic n. sp. (S. France), B. sardoa n. sp. (Sardinia), B. sophiae n. sp. (S. France), and B. sunnivae n. sp. (Sicily)).
- BERENTS, P.B., 1985. Warragaia rintouli n. gen., n. sp. (Amphipoda: Urohaustoriidae) from New South Wales, Australia. ____ Rec. Austr. Mus. 36, 253-258.
- BIANCHI, C.N., G. DIVIACCO & C. MORRI, 1983-84. (Faunal interchanges between a coastal lagoon and the adjacent marine tract in benthic communities of the Po River delta (Northern Adriatic): methodological premises.) ____ Nova Thalassia 6 Suppl. 201-206. (In Italian. Amphipods on p. 203).
- BIESINGER, K.E. & G.N. STOKES, 1986. Effects of synthetic polyelectrolytes on selected aquatic organisms. ____ J. Water Poll. Control Fed. 58, 207-213. (Not seen, i.a. Hyaella).
- BIRD, G.A. & N.K. KAUSHIK, 1985. Processing of decaying maple leaf, Potamogeton and Cladophora packs by invertebrates in an artificial stream. ____ Arch. Hydrobiol. 104, 93-104. (i.a. Gammarus pseudolimnaeus, the only organism tested that preferred maple leaves).
- BIRKHEAD, T.R. & S. PRINGLE, 1986. Multiple mating and paternity in Gammarus pulex. ____ Anim. Behav. 34, 611-613.
- BLOMQUIST, E.-M. & E. BONSDORFF, 1986. Spatial and temporal variations of benthic macrofauna in a sand-bottom area on Åland, northern Baltic Sea. ____ Ophelia, Suppl. 4, 27-36.
- BLUZAT, R. & J. SEUGE, 1983. Effet d'une intoxication par le thirame présent dans le milieu ou la nourriture chez le crustacé Gammarus pulex. ____ Environm. Poll. 31 A, 133-147.
- BOLOGNA, M.A. & A. VIGNA TAGLIANTI, 1985. (Cave fauna of the Ligurian Alps). ____ Ann. Mus. Civ. Stor. Nat. G. Doria 84 bis (1984), 1-389. (In Italian, not seen. Amph. pp. 222-223).
- BOLT, S.R.L., 1985. Urine clearance rates and apparent permeability of Gammarus duebeni exposed to varying conditions. ____ J. exp. Biol. 114 673-6761.
- BONSDORFF, E., J. MATTILA, C. RONN & C-S ÖSTERMAN, 1986. Multidimensional interactions in shallow soft-bottom ecosystems; testing the competitive exclusion principle. ____ Ophelia Suppl. 4, 37-44.
- BORODICH, N.D., L.F. BONDERENKO & V.L. VAVROV, 1985. (Ponto-Caspian Crustacea of the middle Volga.) ____ Informatsionnyi Byull. Biol. vnutr. Vod. (65), 25-28. (In Russian, not seen).
- BOROWSKY, B., 1985. Responses of the amphipod crustacean Gammarus palustris to waterborne secretions of conspecifics and congeners. ____ J. chem. Ecol. 11, 1545-1552. (Not seen, unfortunately).
- BOROWSKY, B., 1985. Differences in reproductive behavior between two male morphs of the amphipod crustacean Jassa falcata Montagu. ____ Physiol. Zool. 58, 497-503.
- BOROWSKY, R., B. BOROWSKY, H. MILANI & P. GREENBERG, 1985. Amylase variation in the salt marsh amphipod, Gammarus palustris. ____ Genetics 111, 311-323.
- BORTEKIVICH, L. V., 1985. (Daily vertical migration of corophiids (Amphipoda, Corophiidae) in estuaries of northwestern Prichernomor'ya). ____ Gidrobiol. Zh. 21 (5), 100-101. (In Russian, not seen).
- BORTEKIVICH, L. V., N.N. KHMELEVA & N.I. SADOVSKAYA, 1984. (Ecological characteristics, caloric value and production of Corophium robustum (Amphipoda, Corophiidae) from the lower Dnieper.) ____ Gidrobiol. Zh. 1984-2, 93-99. (In Russian, not seen).
- BOULIGAN, Y., 1986. Theory of microtomy artifacts in arthropod cuticle. ____ Tissue & Cell 18, 621-643.

- BOULTON, A. J., 1985. A sampling device that quantitatively collects benthos in flowing or standing waters. ____ *Hydrobiologia* **127**, 31-39. (Not seen).
- BOUSFIELD, E.L., 1985. The ubiquitous amphipods, mysterious insects of the sea. ____ *Rotunda* **18**, 31-35. (A popular essay, with beautiful colour pictures).
- BOUSFIELD, E.L. & R.W. HEARD, 1986. Systematics, distributional ecology, and some host-parasite relationships of *Uholorchestia uhleri* (Shoemaker) and *U. spartinophila* new species (Crustacea: Amphipoda), endemic to salt marshes, of the Atlantic coast of North America. ____ *J. crust. Biol.* **6**, 264-274. (*U. spartinophila* n. sp. from the Atlantic seaboard of the USA has Cape Ann, Massachusetts as type locality).
- BOUTIN, C. & N. COINEAU, 1987. *Pseudoniphargus maroccanus* n. sp. (subterranean Amphipoda), the first representation of the genus in Morocco. Phylogenetic relationships and paleogeography. ____ *Crustaceana*, Suppl. **12**.
- BOUTIN, C. & M. MESSOULI, 1987. *Pachypodacrangonyx maroccanus*, n. gen., n. sp., nouveau représentant du groupe *Metacrangonyx* dans les eaux souterraines du Maroc. ____ *Crustaceana*, Suppl. **12**.
- BOWIE, J.Y., 1984. Parasites from an Atlantic bottle-nose dolphin (*Tursiops truncatus*), and a revised checklist of parasites of this host. ____ *N.Z. J. Zool.* **11**, 395-398.
- BRATTEY, J. & A. CAMPBELL, 1986. A survey of parasites of the American lobster, *Homarus americanus* (Crustacea; Decapoda), from the Canadian Maritimes. ____ *Can. J. Zool.* **64**, 1998-2003. (No amphipods!)
- BRINTON, E., 1985. Observations of plankton organisms obtained by bongo nets during the November-December 1983 ice-edge investigations. ____ *Antarct. J. US* **19**, 113-115.
- BROWN, P.W. & L.H. FREDRICKSON, 1986. Food habits of breeding White-winged Scoters. ____ *Can. J. Zool.* **64**, 1652-1654. (85-100% of food was made up by *Hyalella azteca*).
- BUBINAS, A.D., 1985. (Species composition, distribution and biomass of zoomacro-benthos in the Baltic Sea in 1981). ____ *Liet. Tsr Mokslu Akad. Dorb Ser. C Biol. Mokslai* 1985-3, 63-74. (In Russian, with Latvian summary. Not seen).
- BUHL-JENSEN, L., 1986. The benthic amphipod fauna of the West-Norwegian continental shelf compared with the fauna of five adjacent fjords. ____ *Sarsia* **71**, 193-208.
- BULNHEIM, H-P. & A. SCHOLL, 1986. Genetic differentiation between populations of *Talitrus saltator* and *Talorchestia deshayesii* (Crustacea: Amphipoda) from coastal areas of the northwestern European continent. ____ *Mar. Biol.* **92**, 525-536.
- BUSHUEVA, I.V., 1986. (New representatives of the antarctic fauna of scuds (Amphipoda, Gammaridea)). ____ *Zool. Zh.* **65**, 1296-1302. (In Russian. Deals with *Calliopius excellens* n. gen., n. sp. (Calliopidae, with a key to all genera) and *Paramoera incognita* n. sp., both from the Davis Sea. According to the English summary also *Prometopa dorsoundata* n. sp. and *Proloboides* (sic!) *bellansantiniae* n. sp. are described, but this part is absent from the Russian text).
- CAHOON, L.B., C.R. TRONZO & J.C. HOWE, 1986. Notes on the occurrence of *Hyperoche medusarum* (Kroyer) (Amphipoda, Hyperidae), with Ctenophora off North Carolina, USA. ____ *Crustacean* **51**, 95-97.
- CAINE, E.A., 1986. Carapace epibionts of nesting loggerhead sea turtles: atlantic coasts of USA. ____ *J. exp. mar. Biol. Ecol.* **95**, 15-26. (Amph. p. 20. The most common amph. are *Caprella andreae* and *Podocerus cheloniae*, in southern populations also a *Hyale* sp.).
- CARR, W.E.S. & C.D. DERBY, 1986. Chemically stimulated feeding behavior in marine animals. Importance of chemical mixtures and involvement of mixture interactions. ____ *J. chem. Ecol.* **12**, 989-1011.
- CASTROVIEJO, R.A. & J.M. BERMUDEZ, 1979. (On twelve marine amphipod species new for the Spanish littoral zone). ____ *Trab. compostelanos Biol.* **8**, 62-70. (In Spanish. *Bathyporeia pelagica*, *Stenothoe spinimana*, *S. valida*, *Pereionotus testudo*, *Periculodes longimanus*, *Tritaea gibbosa*, *Hyale dollfusi*, *Aora atlantidea*, *Biancolina cuniculus*, *Jassa ocia*, *J. dentex* and *Parajassa pelagica*, all from the coast of Galicia).
- CELLOT, B. & M. BOURNAUD, 1986., (Faunistic modification caused by a small variation in discharge in a large river). ____ *Hydrobiologia* **135**, 223-232. (In French, not seen).
- CHAMBERLAIN, Y., M.W. HOLDGATE & N. WACE, 1985. The littoral ecology of Gough Island, South Atlantic Ocean. ____ *Tethys* **11**, 302-319.
- CHAMIER, A-C. & L.G. WILLOUGHBY, 1986. The role of fungi in the diet of the amphipod *Gammarus pulex*: an enzymatic study. ____ *Freshw. Biol.* **16**, 197-208.
- CIPRIANO, F., 1985. Dusky dolphin research at Kaikura, New Zealand. ____ *Mauri Ora* **12**, 151-158. (Not seen. Data on Cyamidae?)
- CLARK, J. & J.L. BARNARD, 1986. *Tonocote* a new genus and species of Zobrachoidae from Argentina (Crustacea: marine Amphipoda). ____ *Proc. biol. Soc. Wash.* **99**, 225-236. (*Tonocote magellani* n. gen., n. sp. from the Magellan Straits. The genus is related to the Australian (*Prantinus*)).
- CLARK, J. & J.L. BARNARD, 1987. *Chono angustiarum*, a new genus and species of Zobrachoidae (Crustacea: Amphipoda) from Magellan strait, with a revision of the Urohaustoriidae. ____ *Proc. biol. Soc. Wash.* **100**, 75-88. (With an extensive discussion, and a key to all genera, of the Urohaustoriidae and the Zobrachoidae, which the authors for the time being keep separated).

- CLARK, A., A. SKADSHEIM & L.J. HOLMES, 1985. Lipid biochemistry and reproductive biology in two species of Gammaridae (Crustacea: Amphipoda). ____ Mar. Biol. (Berl.) **88**, 247-263.
- CLAVIER, J., 1984. (Vertical distribution of benthic macrofauna in an irretrievable fine sediment). ____ Cah. Biol. mar. **25**, 141-152. (In French, not seen).
- COINEAU, N., 1983. Energetic values in interstitial Isopods and amphipod from sandy beaches as a function of body size and season (western Mediterranean). ____ Dev. Hydrobiol. **19**, 687-692.
- COINEAU, N., 1984. Quelques aspects de l'adaptation des crustacés isopodes, amphipodes et syncarides à la vie dans le milieu interstitiel. ____ Mém. biospeol. **11**, 7-15.
- COINEAU, N. & J.H. STOCK, 1986. Amphipoda: Bogidiellidae. ____ Pp. 574-580 in L. BOTOSANEANU (ed.). Stygofauna mundi. E.J. Brill, Leiden (*Artesia* is by these authors considered to be a bogidiellid).
- COLE, G.A., 1985. Analysis of the *Gammarus pecos* - complex (Crustacea: Amphipoda) in Texas and New Mexico, USA. ____ J. Arizona - Nevada Acad. Sci. **20**, 83-103. (The complex probably contains two undescribed spp., besides *G. pecos*, *G. hyalelloides* and *G. desperatus*).
- COLEBROOK, J.M., 1985. Continuous plankton records: overwintering and annual fluctuations in the abundance of zooplankton. ____ Mar. Biol. (Berl.) **88**, 261-265. (i.a. Hyperiididae, treated as a group).
- CRONIN, T.W., 1986. Optical design and evolutionary adaptation in crustacean compound eyes. ____ J. crust. Biol. **6**, 1-23.
- CULVER, D.C., 1987. Eye morphometrics of cave and spring populations of *Gammarus minus* (Amphipoda: Gammaridae). ____ J. crust. Biol. **7**, 136-147.
- CURTIS, D.J., C.G. GALBRAITH, J.C. SMYTH & D.B.A. THOMPSON, 1985. Seasonal variations in prey selection by estuarine Black-headed gulls (*Larus ridibundus*). ____ Est. coast. Shelf Sci. **23**, 75-89. (i.a. *Corophium volutator*).
- CURTIS, D.J. & D.B.A. THOMPSON, 1985. Spacing and foraging behaviour of Black-headed Gulls *Larus ridibundus* in an estuary. ____ Ornis Scand. **16**, 245-252. (*Corophium volutator* important prey).
- DAUVIN, J.-C., 1985. Sur deux espèces d'Amphipodes nouvelles pour la faune marine de Roscoff. ____ Cah. Biol. mar. **26**, 469-471. (*Synchelidium maculatum* and *Melphidella macra*).
- DAUVIN, J. C. & D. BELLAN-SANTINI, 1985. Collection des Ampelisцидés d'Edouard Chevreux du Muséum National d'Histoire Naturelle: description *Ampelisca melitae* et d'*A. monoculata* n. spp. et redescription d'*A. verga* Reid. ____ Bull. Mus. natn Hist. nat. Paris (4) **7** A, 659-675. (*A. melitae* n. sp. from Tunisia, *A. monoculata* n. sp. from off the coast of the Sahara at 17N. *A. aequicornis* var. *verga* of Reid, 1951 from near Dakar is redescribed as a full species, *A. verga* Reid).
- DE BROYER, C., 1985. Description de *Falklandia* n. gen. de l'Océan Austral et définition des Lysianassoidea Uristidiens (Crustacea, Amphipoda). ____ Zool. Scripta **14**, 303-312. (*Falklandia* n. gen. is erected for *Orchomenopsis reducta*. The uristid group of genera is discussed and tentatively subdivided according to the mandible morphology).
- DE PAUW N., D. ROELS & A.P. FONTOURA, 1986. Use of artificial substrates for standardized sampling of macroinvertebrates in the assessment of water quality by the Belgian Biotic Index. ____ Hydrobiologia **133**, 237-258.
- DE SMET, W.M.A. & R. ASSELBERG, 1985. (A porpoise with skin lesions). ____ Marwin **6**, 123-138. (In Flemish. First Belgian record of *Isocyamus delphinii*, from *Phocaena phocaena*).
- DESROSIERES, G., D. BELLAN-SANTINI & J.-C. BRETHES, 1986. Trophic organization of four rock-bottom assemblages along a gradient of industrial pollution (Gulf of Fos, France). ____ Mar. Biol. **91**, 107-120.
- DESSAIX, J., 1986. Structure et fonctionnement des écosystèmes du Haut-Rhône français. 27. Dynamique des populations des gammares. ____ Arch. Hydrobiol. **106**, 541-558.
- DESSAIX, J., 1986. Structure et fonctionnement des écosystèmes du Haut-Rhône français. 28. Estimation de production de gammares. ____ Arch. Hydrobiol. **107**, 125-141.
- DEWITT, T.H. & LEVINTON, 1985. Disturbance, emigration, and refugia: how the mud snail, *Ilyanassa obsoleta* (Say), affects the habitat distribution of an epifaunal amphipod, *Microdeutopus gryllotalpa* (Costa). ____ J. exp. mar. Biol. Ecol. **92**, 97-113.
- DIAZ-CASTANEDA, V., 1984. Etude de la colonisation benthique de sédiments mis en enceintes expérimentales dans la baie du Lazaret (Toulon, France). ____ Thèse 3. Cycle, Univ. Aix-Marseille, 231 pp. (Not seen. Amph. important especially in first 3 months).
- DIVICACCO, G., 1983. (?) (Considerations on the lysianassid *Socarnes filicornis* (Heller) (Crustacea, Amphipoda). ____ Boll. Mus. civ. Stor. nat. Verona **10**, 539-540. (In Italian. Synonyms of this mediterranean species are *Anonyx schmardae* and *Socarnopsis crenulata*. *S. erythrophthalmus* probably does not occur in the Mediterranean).
- DIVIACCO, G., 1985. (The amphipod crustaceans of the marine caves of Bergeggi (Ligurian Sea)). ____ Olbalia (N.S.) **11**, 785-787. (In Italian, not seen).

- DIVIACCO, G., 1986. (First record of Trischizostoma raschi Esmark & Boeck in the Mediterranean Sea (Crustacea, Amphipoda)). ____ Boll. Mus. civ. Stor. Nat. Verona 12 (1985), 513-515. (In Italian).
- DIVIACCO, G., & S. RUFFO, 1985. New Bogidiellidae of the African subterranean waters. ____ Monit. zool. ital., Suppl. 20, 135-148. (Not seen. Afridiella messanae n. sp. from S. Somalia. Maghrebiella maroccana n. gen., n. sp. (Bogidiellidae) from a well near Marrakesh, Morocco).
- DONN, T.E. & R.A. CROKER, 1983. Production ecology of Haustorius canadensis (Amphipoda: Haustoriidae) in southern Maine. ____ Dev. Hydrobiol. 19, 661-667. (Not seen).
- DONN, T. E. & R.A. CROKER, 1986. Life-history patterns of Haustorius canadensis (Crustacea: Amphipoda) in northern New England. ____ Can. J. Zool. 64, 99-104.
- DONN, T. E. & R.A. CROKER, 1986. Seasonal patterns of production in the sandy-beach amphipod Haustorius canadensis ____ Est. coast. Shelf Sci. 22, 675-688.
- DOWNING, J.-A., 1986. A regression technique for the estimation of epiphytic invertebrate populations. ____ Freshw. Biol. 16, 161-174.
- DOWNING, J.-A. & H. CYR, 1985. Quantitative estimation of epiphytic invertebrate populations. ____ Can. J. Fish. aq. Sci. 42, 1570-1579.
- DUCK, R.W., 1986. Traces made by the amphipod Gammarus in subaerially-exposed marginal sediments of a freshwater lake. ____ Boreas (Oslo) 15, 19-23.
- DUCRUET, J. 1986. Localisation de l'organe Y chez le crustacé amphipode Gammarus pulex (L., 1758). ____ Crustaceana 51, 231-234.
- DUNCAN, K.W., 1985. Cuticular microstructures of terrestrial Amphipoda (Crustacea, family Talitridae). ____ Zool. Anz., Jena 215, 140-146.
- DUNCAN, K.W., 1985. A critique of the concept of genetic assimilation as a mechanism in the evolution of terrestrial talitrids (Amphipoda). ____ Can. J. Zool. 63, 2230-2232.
- DUNHAM, P. J., 1986. Mate guarding in amphipods: a role for brood pouch stimuli. ____ Biol. Bull. 170, 526-531. (A study on Gammarus lawrencianus).
- EDGAR, G.J. & P.G. MOORE, 1986. Macro-algae as habitats for motile macrofauna. ____ Pp. 255-277 in B. SANTELICES (ed.) Usos y Funciones Ecologicas de las Algas Marinas Bentonicas. Monogr. biol. 4.
- EDWARDS, J.M. & E. NAYLOR, 1987. Endogenous circadian changes in orientational behaviour of Talitrus saltator. ____ J. mar. biol. Ass. U.K. 67, 17-26.
- ELEFThERIOU, A., M.R. ROBERTSON & D.J. MURISON, 1986. The benthic fauna of sandy bays, with particular reference to Irvine Bay. ____ Proc. Roy. Soc. Edinb. 90 B, 317-327.
- ELKAIM, B., J.P. IRLINGER & S. PICHARD, 1985. Dynamique de la population d'Orchestia mediterranea L. (Crustacé, Amphipode) dans l'estuaire du Bou Regreg (Maroc). ____ Can. J. Zool. 63, 2800-2809.
- ELMGREN, R., S. ANKAR, B. MARTELEUR & G. EJDUNG, 1986. Adult interference with postlarvae in soft sediments: the Pontoporeia - Macoma example. ____ Ecology 67, 827-836.
- ELSTAD, C.A., 1986. Macrobenthic distribution and community structure in the upper navigation pools of the Upper Mississippi River. ____ Hydrobiologia 136, 85-100.
- ETTEN, J. van & A.M. VOUTE, 1983 (Orchestia gammarella (Pallas)). An arthropod suitable for study in schools and universities for field and laboratory practicals). ____ Valblad Biol. 63, 261-267. (In Dutch, not seen).
- EWELL, W.S., J.W. GORSUCH, R.O. KRINGLE, K.A. ROBILIARD & R.C. SPIEGEL, 1986. Simultaneous evaluation of the acute effects Toxico. Chem. 5, 831-840. (I.a. Gammarus fasciatus).
- FELGENHAUER, B.E., 1987. Techniques for preparing crustaceans for scanning electron microscopy. ____ J. crust. Biol. 7, 71-76.
- FINGERMAN, M., 1987. The endocrine mechanisms of crustaceans. ____ J. crust. Biol. 7, 1-24. (A review paper).
- FIOCCO, G., A. GUERRINI & L. PARDI, 1983. (Spectral differences in sky radiance over land and sea, and orientation of the littoral amphipod Talitrus saltator Montagu). ____ Atti Accad. naz. Linzei Rc. Ser. 8, 74 25-33. (In Italian).
- FOECKLER, F. & E. SCHRIMPF, 1985. Gammarids in streams of northeastern Bavaria, F.R.G. 2. The different hydrochemical habitats of Gammarus fossarum Koch, 1835 and Gammarus roeseli Gervais, 1835. ____ Arch. Hydrobiol. 104, 269-286.
- FORWARD, F.B., 1986. Behavioral responses of a sand-beach amphipod to light and pressure. ____ J. exp. mar. Biol. Ecol. 102, 55-74.
- FOX, R.S. & E.E. RUPPERT, 1985. Shallow-water marine benthic invertebrates of South Carolina: Species identification, community composition and symbiotic association. Belle W. Baruch Libr. mar. Sci. 14, 336 pp. Univ. S. Carol. Press. (Not seen, unfortunately).

- FREDETTE, T. J. & E.J. DIAZ, 1986. Life history of Gammarus mucronatus Say (Amphipoda: Gammaridae) in warm temperate estuarine habitats, York river, Virginia. ____ J. crust. Biol. 61, 57-78.
- FREDETTE, T.J. & E.J. DIAZ, 1986. Secondary production of Gammarus mucronatus Say (Amphipoda: Gammaridae) in warm temperate estuarine habitats, York River, Virginia. ____ J. crust. Biol. 6, 729-741.
- FRIEND, J.A. & P.K.S. LAM, 1985. (Occurrence of the terrestrial amphipod Talitroides topitotum (Burt) on Hong Kong island.) ____ Acta zootaxon. sin. 10, 27-33. (In Chinese and English).
- FRIEND, J.A. & A.M.M. RICHARDSON, 1986. Biology of terrestrial amphipods. ____ Ann. Rev. Entomol. 31, 25-48. (An important review paper).
- FRITHSEN, J.B., D.T. RUDNICK & P.H. DOERING, 1986. The determination of fresh organic carbon weight from formaldehyde preserved macrofaunal samples. ____ Hydrobiologia 133, 203-208.
- FRUTIGER, A., 1985. A versatile flow channel for laboratory experiments with running water macroinvertebrates. ____ Rev. suisse Hydrobiol. 46, 301-305. (Not seen).
- FÜTTERER, D. (ed), 1986. Die Expedition Antarktis IV mit FS 'Polarstern' 1985/86. ____ Ber. Polarforsch. 32, 1-114. (On pp. 28-31 B. Christiansen reports on a collection of Eurythenes gryllus from the North Atlantic).
- GALBRAITH, C.G., J.C. SMYTH & D.B.A. THOMPSON, 1985. Seasonal variations in prey selection by estuarine Black-headed Gulls (Larus ridibundus). ____ Est. coast. Shelf Sci. 21, 75-90. (i.a. Corophium volutator).
- GARDNER, W.S., T.F. NALEPA, W.A. FREZ, E.A. CICHOCKI & P.F. LANDRUM, 1985. Lipid content and energy flow through Lake Michigan macroinvertebrates. ____ Internat. Assoc. Great Lakes Res., Toronto 1985, 40. (Abstract only, i.a. Pontoporeia hoyi).
- GARDNER, W.S., T.F. NALEPA, W.A. FREZ, E.A. CICHOCKI & P.F. LANDRUM, 1985. Seasonal pattern in lipid content of Lake Michigan macroinvertebrates. ____ Can. J. Fish. aq. Sci. 42, 1827-1832.
- GINET, R. & V. AELLEN, 1985. (Synonymy of Niphargus godeti and Niphargus virei (Crustacea Amphipoda) from subterranean waters: preeminence of Niphargus virei.) ____ Rev. suisse Zool. 92, 107-114. (In French, not seen. N. godeti Wrezniewski, though a senior synonym, is considered a nomen oblitum).
- GINET, R., 1985. Presence de l'amphipode hypogé Niphargus dans certains lacs alpins de haute-montagne. ____ Verh. int. Ver. theor. angew. Limnol. 22, 3220-3222. (Not seen).
- GOEKE, G.D., 1985. Amphipods of the family Ampeliscidae (Gammaridea). 5. Ampelisca hawaiiensis, new species. ____ Pacif. Sci. 39, 261-265. (= A. schellenbergi s. Barnard 1967, 1970. Type locality: near Honolulu).
- GOEKE, G.D., 1987. Amphipods of the family Ampeliscidae (Gammaridea). VI Ampelisca macrodonta, a new species from the Falkland Islands. ____ Proc. biol. Soc. Wash. 100, 4-7.
- GOLIKOV, A.A., 1985. (Studies on the distribution and ecology of Onega Bay of the White Sea.) ____ Explor. Fauna SSSR 33, 150-170. (In Russian. Many important biological data).
- GONZALEZ, E., 1986. A new record of Paracorophium hartmannorum Andres, 1975, from the Chilean coast, with a description of the adult (Amphipoda: Corophiidae). ____ Proc. biol. Soc. Wash. 99, 21-28. (P. chilense is considered to be a junior synonym of P. hartmannorum).
- GOOCH, J.L. & D.S. GLAZIER, 1986. Levels of heterozygosity in the amphipod Gammarus minus in an area affected by Pleistocene glaciation. ____ Am. Midl. Nat. 116, 57-63.
- GOOD, L.K., R.C. BAYER, M.L. GALLAGHER & J.H. RITTENBURG, 1982. Amphipods as potential diet for juveniles of the American lobster Homarus americanus (Milne Edwards). ____ J. Shellfish Res. 2, 183-187.
- GOODING, R.U., 1986? Animals associated with the sea urchin, Diadema antillarum. ____ Pp. 334-336 in T. J. BRIGHT & L.H. PEQUEGNAT. Biota of the West Flower Garden Bank. Gulf Publishing Co. Houston. (An undescribed Lysianassa sp. from Puerto Rico, to be described by Gooding, lives in association with Diadema).
- GRAF, F., 1986. Fine determination of the molt cycle stages in Orchestia cavimana Heller (Crustacea: Amphipoda). ____ J. crust. Biol. 6, 666-678.
- GRANEY, R.L. & J.P. GIESEY, 1986. Seasonal changes in the free amino acids pool of the freshwater amphipod Gammarus pseudolimnaeus (Crustacea: Amphipoda). ____ Comp. Biochem. Physiol. A 85, 535-544.
- GREEN, K., 1986. Food of the Emperor Penguin Aptenodytes forsteri on the Antarctic fast ice edge in late winter and early spring. ____ Polar Biol. 6, 187-188. (Many amphipods, esp. in October).
- GREENWOOD, P.J. & J. ADAMS, 1984. Sexual dimorphism in Gammarus pulex: the effect of current flow in precopula pair formation. ____ Freshw. Biol. 14, 203-209.
- GREENWOOD, P.J. & J. ADAMS, 1987. Sexual selection, size dimorphism and a fallacy. ____ Oikos 48, 106-108.

- GREZE, I.I., 1985. (Amphipods). ____ Naukova Dumka, Kiev (UKSSR). Fauna Ukrainy 26, 172 pp. (This Russian monograph deals with *Orchomene humilis*, *Nannonyx goesi reductus*, *Ampelisca diadema*, *Bathyporeia guillamsoniana*, *Harpinia dellavallei*, *H. crenulata*, *Stenothoe monoculoides*, *Colomastix pusilla*, *Perioculodes longimanus*, *Synchelidium maculatum*, *Monoculodes gibbosus*, *Apherusa bispinosa*, *Atylus guttatus*, *Dexamine spinosa*, *Tritaeta gibbosa*, *Gammarus insensibilis*, *G. aeguicauda*, *G. subtypicus*, *G. crinicornis*, *Marinogammarus olivii*, *Chaetogammarus ischnus*, *C. l. behningi*, *C. oliviformis* n. sp., *Gammarellus carinatus*, *Megaluropus agilis*, *Melita palmata*, *Cheirocratus sundevalli*, *Pontogammarus maoticus*, *Cardiophilus baeri* (with which *C. marisnigri* and *C. miloslawskajae* are synonymized), *Pleonexes gammaroides*, *Ampithoe vaillanti*, *Cymadusa crassicornis*, *Microdeutopus gryllotalpa*, *M. anomalus*, *M. damnoniensis*, *Coremapus versiculatus*, *Leptocheirus pilosus*, *Microtopopus longimanus*, *M. minutus* Sowinsky (revived), *Megamphopus cornutus*, *Photis longicaudata*, *Biancolina algicola*, *Jassa oia*, *J. falcata*, *Erichthonius difformis*, *Corophium crassicornis*, *C. bonnellii*, *C. acherusicum*, *C. volutator* (incl. *C. orientale*), *C. runicorne*, *Siphonocetes dellavallei*, *Chelura terebrans*, *Orchestia gammarella* (sic.), *O. mediterranea*, *O. montagui*, *O. bottae*, *Talorchestia deshavesii*, *Talitrus saltator*, *Hyale pontica*, *H. dollfusi*, *H. prevostii*, *H. perieri*, *Phthisica marina*, *Pseudoprotella phasma*, *Caprella a. acanthifera*, *C. a. ferox*, *C. liparotensis*, *C. mitis*, and *C. danilevskii*). All species are fully described and illustrated, and extensive ecological data provided.
- GROSSO, L.E. & L. GLAPS, 1985. (Geographic distribution of the family Bogidiellidae (Crustacea, Amphipoda) in Argentina, with description of a new subgenus and a new species). ____ Physis, Sec. B 43, 49-55. (In Spanish. The new subgenus *Dyticogidiella* is erected for *B. talampuyensis* n. sp. from La Rioja province).
- GUIDI, L.D., 1986. The feeding response of the epibenthic amphipod *Siphonocetes dellavallei* to varying food particle sizes and concentrations. ____ J. exp. mar. Biol. Ecol. 96, 51-64.
- HACKSTEIN, E., H. SCHIRMER & H. LEBSCH, 1985. Die Veränderungen populationsdynamischer Parameter bei *Gammarus tigrinus* Sexton (Crustacea: Amphipoda) durch Cadmium- und Temperaturstress. ____ Verh. Ges. Ökol. 13, 605-612. (Not seen).
- HACKSTEIN, E., H. SCHIRMER & H. LEBSCH, 1986. (Studies on the population dynamics of *Gammarus tigrinus* (Crustacea: Amphipoda) in the river Weser near Bremen). ____ Arch. Hydrobiol. 105, 443-458. (In German, not seen).
- HAESLOOP, U. & M. SCHIRMER, 1985. Accumulation of orally administered cadmium by the eel (*Anguilla anguilla*). ____ Chemosphere 14 1627-1634. (Gammarids used as food).
- HALCROW, K., 1985. The fine structure of the pore canals of the talitrid amphipod *Hyale nilssonii* Rathke. ____ J. crust. Biol. 5, 606-615.
- HAMASHINA, W. & H. MORINO, 1984. (Experimental study on growth, survivorship and breeding activity of *Jesogammarus* sp. in different temperatures). ____ Benthos Res. 26, 43-44. (In Japanese; not seen).
- HAMELKINK, J.L., D.R. BUCKLER, F.L. MAYER, D.H. PALEWSKI & H.O. SANDERS, 1986. Toxicity of fluridane to aquatic invertebrates and fish. ____ Environm. Toxicol. Chem. 5, 87-94. (i.a. *Gammarus pseudolimnaeus*).
- HARGEBY, A., 1986. A simple trickle chamber for rearing aquatic invertebrates. ____ Hydrobiologia 133, 271-274. (Designed for *Gammarus pulex*).
- HARRISON, N.M., 1984. Predation on jellyfish and their associates by seabirds. ____ Limnol. Oceanogr. 29, 1335-1337. (Not seen).
- HARTLEY, J.P. & J.D.D. BISHOP, 1986. The macrobenthos of the Beatrice Oilfield, Moray Firth, Scotland. ____ Proc. roy. Soc. Edinb. 91 B, 221-245. (Amph. pp. 242-243).
- HARTOG, C. den & G. v.d. VELDE, 1987. Invasions by plants and animals into coastal brackish and fresh water of the Netherlands. ____ Proc. kon. Ned. Akad. Wetensch. C 90, 31-37.
- HAWKINS, C.M., 1985. Population carbon budgets and the importance of the amphipod *Corophium volutator* in the carbon transfer on a Cumberland Basin mudflat, upper Bay of Fundy, Canada. ____ Neth. J. Sea Res. 19, 165-176.
- HEPPLESTON, P.B., 1984. *Gammarus pulex* (L.) in Orkney, Scotland (Amphipoda). ____ Crustaceana 46, 220.
- HERRING, P.J., 1985. Bioluminescence in the Crustacea. ____ J. crust. Biol. 5, 557-573.
- HESSLER, R.R., 1985. Swimming in Crustacea. ____ Trans. R. Soc. Edinb., Earth. Sci. 76, 115-122. (Not seen, unfortunately).
- HINES, A.H. & K.T. COMTOS, 1985. Vertical distribution of infauna in sediments of a subestuary of Central Chesapeake Bay. ____ Estuaries 8, 296-304.
- HIPPA, H., S. KOPONEN & R. MANNILA, 1986. Invertebrates of Scandinavian caves. 5. Diplopoda, Chilopoda, Isopoda and Amphipoda. ____ Memor. Fauna Flora fenn. 62, 1-4. (*Gammarus pulex* only).
- HIRAYAMA, A., 1985. Taxonomic studies on the shallow water gammaridean Amphipoda of West Kyushu, Japan. 5. Leucothoidae, Liljeborgiidae, Lysianassidae (Parchynella, Aristias, Waldeckia, Ensayara, Lepidepcreum, Hippomedon, and Anonyx). ____ Publ. Seto mar. biol. Lab. 30, 167-212. (Describes and illustrates *Leucothoides pottsii*, *Leucothoe alata*, *L. bidens* n. sp., *Listriella orientalis* n. sp., *Prachynella lodo*, *Aristias nonspinus* n. sp., *Ensayara dentarius* n. sp., *Lepidepcreum vitjazii*, *Hippomedon pacificus* and *Anonyx simplex* n. sp.).
- HIRAYAMA, A., 1986. Taxonomic studies on the shallow water gammaridean Amphipoda of West Kyushu, Japan. 6. Lysianassidae (*Orchomene*), *Megaluropus* family group, Melitidae (*Cottesloe*, *Jerbarnia*, *Maera*, *Ceradocus*, *Eriopisella*, *Dulichella*). ____ Publ. Seto mar. biol. Lab. 31, 1-35. (Deals with *Orchomene breviceps* n. sp., *O. liomargo* n. sp., *O. tomiokaensis* n. sp., *O. orchospina* n. sp., *Megaluropus massiliensis*, *Jerbarnia aquilopacifica* n. sp. Keys to Japanese *Orchomene* and *melitids* are also provided).

- HIRAYAMA, A., 1986. Two new subspecies of *Synchelidium* (Crustacea: Amphipoda: Oedicerotidae) from the sea shore of North Japan. ____ Zool. Sci. (Tokyo) **3**, 357-366. (*S. americanum latipalpus* n. ssp. and *S. miraculum lenorostratum* n. ssp., both from Otsuchi Bay in NE Japan).
- HOLMES, J.M.C., 1984. Crustacean records from Lough Ine, Co. Cork. Part 3. ____ Bull. Ir. biogeogr. Soc. **8**, 19-25. (Not seen. Amph?).
- HOLSINGER, J.R., 1986. Amphipoda: Holarctic Crangonyctid Amphipods. ____ Pp. 535-349 in L. BOTOSANEANU (ed.). Stygofauna mundi. E.J. Brill, Leiden. (Comprises here the genera *Allocrangonyx*, *Batrurus*, *Crangonyx*, *Lyurella*, *Procrangonyx*, *Pseudocrangonyx*, *Stygobromus* and *Synurella*. In a forthcoming paper by the same author *Allocrangonyx*, *Procrangonyx* and *Pseudocrangonyx*, will be removed from the Crangonyctidae).
- HOLSINGER, J.R., 1986. Amphipoda: Sebidae. ____ Pp. 568-569 in L. BOTOSANEANU (ed.). Stygofauna mundi. E.J. Brill, Leiden.
- HOLSINGER, J.R. & J. YAGER, 1985. A new genus and two new species of subterranean amphipod crustaceans (Hadziidae) from the Bahamas and Turks and Caicos islands (West Indies). ____ Bijdr. Dierk. **55**, 283-294. (*Bahadzia* n. gen., related to *Metaniphargus* and *Saliweckelia*, with type *B. williamsi* n. sp. (great Abaco Isl.) and *B. stocki* n. sp. (Turks and Caicos Isl. Both species are from anchialine caves).
- HOLTHUIS, L.B., 1985. Rafinesque's crustacean genera *Heterelos* and *Yalomus*. ____ Zool. Meded. Leiden ?, 133-147. (The rediscovery of a forgotten ms by Rafinesque show i.a. that *Heterelos* is a synonym of *Phronima*).
- HOLTHUIS, L. B., 1986. The data of publication of Crustaceana Vols. 1-50 and Suppl. 1-10. ____ Crustaceana **51**, 221-222.
- HOOGENBOOM, J. & J. HENNEN, 1985. Etude sur les parasites du macrozooplankton gelatineux dans la rade de Villefranche-sur-Mer (France), avec description des stades de developpement de *Hyperoche mediterranea* Senna (Amphipodes Hyperiidae), ____ Crustaceana **49**, 233-243.
- HOOPER, R.G., 1986. A spring breeding migration of the Snow Crab, *Chionoecetes opilio* (O. Fabr.), into shallow water in Newfoundland. ____ Crustaceana **50**, 257-264. (see pp. 262-263 for data on an unidentified associated amphipod).
- HOPKINS, T. L., 1985. Food web of an Antarctic midwater ecosystem. ____ Mar. Biol., Berl. **89**, 197-212.
- HORNER, R. & D. MURPHY, 1985. Species composition and abundance of zooplankton in the nearshore Beaufort Sea in winter-spring. ____ Arctic **38**, 201-209.
- HOUSTON, K.A. & R.L. HAEDRICH, 1986. Food habits and intestinal parasites of deep demersal fishes from the upper continental slope east of Newfoundland, northwest Atlantic Ocean. ____ Mar. Biol. **92**, 563-574.
- HUDSON, A.V. & D. REYNOLDS, 1984. Distribution of Irish intertidal Talitridae. ____ Bull. Ir. biogeogr. Soc. **8**, 63-78. (Not seen).
- HUH, S.-H., 1986. (Ontogenetic food habits of four common fish species in seagrass meadows). ____ J. oceanol. Soc. Korea **21**, 25-33. (In Korean. Amphipods important prey).
- HUNTER, J. & D. RENDALL, 1986. The sub-littoral fauna of the Inverness, Cromarty and Dornach Firths. ____ Proc. roy. Soc. Edinb. **91** B, 263-274. (Amph. pp. 272-273).
- ICELY, J.D. & J.A. NOTT, 1985. Feeding and digestion in *Corophium volutator* (Crustacea: Amphipoda). ____ Mar. Biol., Berl. **89**, 183-196.
- ILIFFE, T.M., 1986. The zonation model for the evolution of aquatic fauna in anchialine caves. ____ Stygologia **2**, 2-9. (Not seen).
- IMADA, K. & T. KIKUCHI, 1984. (Some reproductive traits and seasonal population fluctuations of three caprellids (Crustacea: Amphipoda) on *Sargassum horneri*). ____ Benthos Res. **26**, 45-48. (In Japanese, not seen).
- ISHIMARU, S., 1985. Taxonomic studies of the family Pleustidae (Crustacea Amphipoda Gammaridea) from coastal waters of northern Japan. 3. The genus *Pleusirus*, with notes on body aesthetascs. ____ J. Fac. Sci., Hokkaido Univ., Ser. 6. Zool. **24**, 103-12. (Redescribes *Pleusirus securus*, and notes the presence and morphology of dorsal 'body aesthetascs' in this species, *Parapleustes tricuspis*, and *Pleusymtes mucidus*).
- ISHIMARU, S., 1985. A new species of the genus *Leucothoe* (Amphipoda Gammaridea Leucothoidae) from Japan. ____ Proc. jap. Soc. syst. Zool. **30**, 46-52. (*L. nagatai* n. sp. (= *L. alata* s. Nagata) from solitary ascidians at Osoro and Misaki, Japan).
- ISHIMARU, S., 1986. Records of *Iphiplateia whiteleggei* (Crustacea Amphipoda Phliantidae) from Hokkaido, Japan. ____ J. Fac. Sci., Hokkaido Univ., Ser. 6 Zool. **24**, 173-181.
- ISHIMARU, S. & K. IKEHARA, 1986. A new genus and species of the subfamily Amphilochinae (Amphipoda, Gammaridea, Amphilochidae) found in the Japan Sea. ____ Zool. Sci. (Tokyo) **3**, 193-198. (*Paramphilochus parachelatus* n. gen. n. sp., closely related to *Amphilochus*).
- IVAJUSHINA, E.A., 1984. (The life cycle of *Atylus carinatus* (Crustacea Amphipoda) in the Kandalaksha Bay (White Sea)). ____ Zool. Zh. **63**, 191-196. (In Russian).
- JACOBI, C.M., 1987. Spatial and temporal distribution of Amphipoda associated with mussel beds from the Bay of Santos (Brazil). ____ Mar. Ecol. Progr. Ser. **35**, 51-58.

- JAGMINIENE, I.B., 1985. (The study of the influence of dimethylacetamide on the viability of the amphipod Pontogammarus robustoides (Grimm)). ____ Akad. Nauk Litov. SSR C 192, 132. (In Russian, not seen).
- JANKE, K., 1986. (Macrofauna and its distribution in the northeastern rocky intertidal zone of the island of Helgoland (North Sea)). ____ Helgol. Meeresunters. 40, 1-56. (In German, not seen).
- JARAMILLO, E., C. BERTRAN, G. AGUILAR, A. TURNER & M. PINO, 1985. Annual fluctuations of the subtidal macroinfauna in an estuary of South of Chile. ____ Stud. neotrop. Fauna Environm. 20, 33-44. (Paracorophium chilense and Cheus sp. codominant).
- JARAMILLO, E., S. MULSOW & R. NAVARRO, 1985. (Intertidal and subtidal macroinfauna in the Queule River Estuary, S. Chile). ____ Rev. chil. Hist. nat. 58, 127-138. (In Spanish, not seen).
- JAŹDŹEWSKI, K., W. JURASZ, W. KITTEL, E. PRESLER, P. PRESLER & J. SICINSKI, 1986. Abundance and biomass estimates of the benthic fauna in Admiralty Bay, King George Island, South Shetland Islands. ____ Polar Biol. 6, 5-16.
- JENSEN, K.T. & J.N. JENSEN, 1985. The importance of some epibenthic predators on the density of juvenile macrofauna in the Danish Wadden Sea. ____ J. exp. mar. Biol. Ecol. 89, 157-174.
- JOHNSON, P.T., 1985. Parasites of benthic amphipods: microsporidians of Ampelisca agassizi (Judd) and some other gammarideans. Fish. Bull. 83, 497-505.
- JOHNSON, P. T., 1986. Parasites of benthic amphipods: ciliates. ____ Fish. Bull. 84, 204-209.
- JOHNSON, P.T., 1986. Parasites of benthic amphipods: dinoflagellates (Duboscquodinia: Synidinidae). ____ Fish. Bull. 84, 605-614.
- JOHNSON, R.K., 1986. Life histories and coexistence of Chironomus plumosus (L.), C. anthracinus Zett. (Diptera: Chironomidae) and Pontoporeia affinis Lindström (Crustacea: Amphipoda) in mesotrophic Lake Erken. ____ Acta Univ. uppsal. 37, 1-20.
- JOHNSON, R.K., 1987. The life history, production and food habitat of Pontoporeia affinis Lindström (Crustacea: Amphipoda) in mesotrophic Lake Erken. ____ Hydrobiologia 144, 277-283. (L. Erken is in Sweden).
- JONES, A.R., 1984. Sedimentary relationships and community structure of benthic crustacean assemblages of reef-associated sediments at Lizard Island, Great Barrier Reef. Coral Reefs 3, 101-112. (Amph. p. 103).
- JONES, A.R., C.J. WATSON-RUSSELL & A. MURRAY, 1986. Spatial patterns in the macrobenthic communities of the Hawkesbury estuary, New South Wales. ____ Austr. J. mar. Freshw. Res. 37, 521-543. (Amph pp. 541-542).
- JUST, J., 1984. Siphonocetinae (Crustacea, Amphipoda, Corophiidae) 3: Concholestes Giles, 1888 and Africoecetes Just, 1983. ____ Steenstrupia 10, 225-234. (Deals with Concholestes dentalii and Africoecetes armatus).
- JUST, J., 1985. Siphonocetinae (Crustacea: Amphipoda: Corophiidae). 4 Australoecetes Just, 1983, including Stebbingoecetes n. subgen. ____ Rec. austr. Mus. 37, 325-341. (Deals with A. selicki, A. (Stebbingoecetes) n. subgen.) australis, and A. jervisi n. sp. (from Jervis Bay, NSW))
- KANAKADURGA, M.R., K. HANUMANTHA RAO & K. SHYAMASUNARI, 1985. A new species of amphipod, Hyale gopalaswamyi sp. nov. - a commensal of sponges. ____ J. Bombay nat. Hist. Soc. 82, 165-170.
- KARAMAN, G.S., 1984. Revision of the Niphargus orcinus - group, part I (fam. Niphargidae) (Contribution to the knowledge of the Amphipoda 130). ____ Montenegrin Acad. Sci. Arts, Glasnik Sect. nat. Sci. 4, 7-79. (Describes and illustrates N. orcinus, N. longiflagellum (was ssp. of orcinus) N. salonitanus, N. arbiter n. sp. (Croatia), N. croaticus, N. s. steueri, N. s. kolombatovici (was ssp. of orcinus), N. podgoriensis, N. hercegovinensis (was ssp. of orcinus), N. v. kusceri (was ssp. of orcinus) and N. v. kusceri f. biletanus (described as N. biletanus)).
- KARAMAN, G.S., 1984. Remarks to the freshwater Gammarus species (fam. Gammaridae) from Korea, China, Japan and some adjacent regions (Contribution to the knowledge of the Amphipoda 134). ____ Montenegrin Acad. Sci. Arts, Glasnik Sect. nat. Sci. 4, 139-162. (Deals with Gammarus galgosensis, G. koreanus Ueno, 1940 (raised to specific rank), G. odaensis, G. s. sobaegensis, G. s. kimi n. ssp. (Cheongsong, Korea), G. s. marginalis n. ssp. (Bongwha, Korea), G. soyoensis, G. zeongogensis, G. gregoryi, G. nekkensis, G. spinipalmus, G. suifunensis, G. taliensis, G. lacustris, 'G. pulex', G. nipponensis, G. chimkentii n. nom (= G. gracilis Martynov 1935, non Rathke 1837), G. songirdaki n. nom. (= G. truncatus Martynov, non Viviani 1805), G. s. hissari n. nom. (= G. truncatus montanus Martynov, non Costa 1851), G. matienus f. stagnalis and G. belli n. nom. (= G. fluvialilis Bell 1921, non M. Edw. 1830)).
- KARAMAN, G.S., 1984. The genus Laurogammarus n. gen. (fam. Gammaridae) in Yugoslavia. (Contribution to the knowledge of the Amphipoda 135). ____ Bilten, Sarajevo B (3) 2, 29-35. (Incomplete ref. in AN 16-20).
- KARAMAN, G.S., 1984. Critical remarks to the fossil Amphipoda with description of some new taxa (Contribution to the knowledge of the Amphipoda 137). ____ Poljoprivreda i Šumarstvo 30-34, 87-104. (Alsacomelita n. gen. is erected for A semipalmata n. sp. (= 'Melita palmata' Maikovsky), Condiogammarus n. gen. for Gammarus retz, and Jubeogammarus n. gen. for G. alsaticus; all 3 taxa from the lower Oligocene of the Alsace. All known fossil amphipods are reviewed).
- KARAMAN, G.S., 1984. Revision of Eriopisa - complex of genera (Gammaridea) (Contribution to the knowledge of the Amphipoda 139). ____ Poljoprivreda i Šumarstvo 30-34, 39-72. (The author divides this complex as follows: Eriopisa with elongata (type) and incisa, Confodiopisa n. gen. (type Psamogammarus caesiculus, also scopulorum and possibly garthi), Flagitopisa n. gen. (for Niphargus philippensis, Imperiopisa

- n. gen. (for *Eriopisa gracilis*), *Psammogammarus* (type *caesus*, also *longiramus*), *Roropisa* n. gen. (type *Victoriopisa atlantica*, also *epistomata*), *Tunisopisa* (type *E. seuratii*), *Victoriopisa* (type *Niphargus chilensis*, also *V. c. griffithsi* n. ssp. (S. Africa) and *australiensis*) and *Vicitopisa* n. gen. (type *E. inaequicaudata*). *V. chilensis* is redescribed from Sri Lanka material).
- KARAMAN, G.S., 1984. Contribution to the knowledge of the Amphipoda. 140. On some gammaridean amphipods from Sri Lanka and adjacent regions. ____ *Studia mar.*, Kotor 15/16, 109-130. (Deals with *Ceradomaera plumosa* (with which *Maera othonides* s. Chilton, K.H. Barnard and Nayar may be identical) and *Quadrivisio bengalensis*. The new genus *Animoceradocus* n. gen. (Melitidae) is erected for *Megamoera semiserrata* (type) and possibly *Ceradocus baffini*).
- KARAMAN, G.S., 1984. Contribution to the knowledge of the Amphipoda. 141. *Quadrus vagabundus*, new genus and species, and revision of genus *Eriopisella* Chevr. (Gammaridea). ____ *Studia mar.* Kotor 15/16, 131-148. (*Quadrus vagabundus* n. gen., n. sp. (Melitidae) is described from Jaffna, Sri Lanka. Karaman revises the *Eriopisella* - group of genera as follows: *Eriopisella* (type *pusilla*, further spp. *capensis*, *epimera sechellensis*, *upolu*), *Cephalopisella* n. gen. (type *E. propagatio*), *Madapisella* (type *E. madagascarensis*), *Nippopisella*, (type *E. nagatai*) and *Spiniferopisella* n. gen. (type *E. spinosa*). The author further notes that the genera *Indocratus* and *Incratella*, both established in 1983, are objective synonyms).
- KARAMAN, G.S., 1984. (?). Contribution to the knowledge of the Amphipoda. 148. *Niphargus krameri* Schell. and *N. spinulifemur* S. Kar. in southern Europe. ____ *Bull. Mus. Hist. nat. Beograd B* 39, 85-104. (Received 1987. *N. spinulifemur*, originally described as ssp. of *N. krameri*, is here raised to specific rank. *N. krameri* is found for the first time in Italy, in the Trieste region).
- KARAMAN, G.S., 1985. Contribution to the knowledge of the Amphipoda 147. *Niphargus tamaninii* Ruffo 1953 and subspecies *N. t. barbatus* n. ssp. (fam. Niphargidae) in Italy. ____ *Poljoprivreda i Šumarstvo* 31 -1, 63-78. (*N. tamaninii* was originally described as ssp. of *N. kochianus*).
- KARAMAN, G.S., 1985. Contribution to the knowledge of the Amphipoda: 146. *Niphargus bodoni*, new species and *Niphargus pescei* in Italy (fam. Niphargidae). ____ *Fragm. balcan. Mus. macedonici Sci. nat.* 12, 81-83. (From Liguria, Italy).
- KARAMAN, G.S., 1985. The taxonomy of *Niphargus transitivus* Sket, 1971, with remarks to *N. armatus* G. Kar., 1985 (fam. Niphargidae) in Italy. (Contribution to the knowledge of the Amphipoda 149). ____ *Poljoprivreda i Šumarstvo* 31, 21-35. (Deals with *Niphargus t. transitivus*, *N. t. dissonus* and *N. armatus*).
- KARAMAN, G.S., 1985. Contribution to the knowledge of the Amphipoda. 151., *Gammarus salemaai*, new species from Lake Ohrid (Macedonia, Yugoslavia) (Family Gammaridae). ____ *Fragm. balc. Mus. Macedonici Sci. nat.* 12, 155-168. (This new species is i.a. characterized by a different chromosome number).
- KARAMAN, G.S., 1985. *Foroniphargus pori*, new genus and species of family Niphargidae. (Contribution to the knowledge of the Amphipoda 152). ____ *Poljoprivreda i Šumarstvo* 31, 51-66. (*E. pori* n. gen., n. sp. (Niphargidae) from subterranean freshwaters of N. Dan., Israel).
- KARAMAN, G.S., 1986. Contribution to the knowledge of the Amphipoda. 142. Two new taxa of suborder Gammaridea from Asia, with remarks to some of Sri Lanka's species. ____ *Poljoprivreda i Šumarstvo* 31 -1, 15-40. (*Dodophotis* n. gen. (Isaeidae) has *Photis distinguenda* as type and *P. digitata* as further species; the latter is redescribed. Also *Grandidierella* (*G. bonnieroides* is redescribed, and a new subgenus *G. (Bigrandidierella)* erected for *Microdeutopus megnae*).
- KARAMAN, G.S., 1986. One new species of family Niphargidae (Gammaridea), *Niphargus farroi* sp. n., from Hungary. ____ *Acta zool. hung.* 32, 61-72.
- KARAMAN, G.S., 1986. New data on the genus *Niphargus* Schiödte (fam. Niphargidae) in Italy and adjacent regions (Contribution to the knowledge of the Amphipoda 138). ____ *Boll. Mus. Civ. Stor. Nat. Verona* 12 (1985), 209-218. (Deals with *Niphargus armatus* n. sp. (Friuli), *N. ictus* n. sp. (Grotta del Fiuma), *N. stefanellii*, *N. hebereri*, *N. pescei* and *N. microcerberus*).
- KARAMAN, G.S., 1986. Contribution to the knowledge of the Amphipoda. 150. One new species of genus *Niphargus* (Gammaridea, Niphargidae) from France, *Niphargus renei* n. sp. ____ *Annis Limnol.* 22, 17-25. (From subterranean waters of the Rhone near Lyon. *N. renei* belongs to the *orcinus-group* of species).
- KARAMAN, G.S. & S. RUFFO, 1986. Amphipoda: *Niphargus* - group (Niphargidae sensu Bousfield). ____ Pp. 514-534 in L. BOTOSANEAU (ed.). *Stygofauna mundi*. E. J. Brill, Leiden.
- KAWAGUCHI, K., O. MATSUDA, S. ISHIKAWA & NAITO, 1986. A light trap to collect krill and other microcrustaceans and planktonic animals under the antarctic coastal fast ice. ____ *Polar Biol.* 6, 37-42.
- KENNEDY, V.S., 1985. A summer benthic survey in Conception Bay, Newfoundland, emphasizing zoogeography of annelids and amphipods. ____ *Can. J. Zool.* 63, 1863-1869. (Eighteen amph. species listed on p. 1868).
- KIM, C.B., 1986. (A taxonomic study of the marine Gammaridae (Crustacea Amphipoda) from Korea). ____ M. Sc. Thesis, Dept Zool. Grad. School, Seoul natn Univ., Seoul. 57 pp. (In Korean. Deals with *Orchestia platensis*, *O. pachypus*, *Talorchestia sinensis*, *T. trinitatis*, *Corophium uenoi*, *Ampithoe lacertosa*, *A. valida*, *Anisogammarus pugettensis* and *Maera pacifica*. Reference provided by Geoff Moore).
- KINGSFORD, M.J. & J. H. CHOAT, 1985. The fauna associated with drift algae captured with a plankton-mesh purse seine net. ____ *Limnol. Oceanogr.* 30, 618-630. (Not seen).

- KLEEF, H.L., 1984. A simple and time-saving method for quantitative collection of Corophium volutator (Crustacea: Amphipoda) from plankton samples rich in detritus. ____ *Hydrobiol. Bull.* **18**, 47-50. (Not seen).
- KOLDING, S., 1985. Genetic adaptation to local habitats and speciation processes within the genus Gammarus (Amphipoda: Crustacea). ____ *Mar. Biol. (Berl.)* **89**, 249-256.
- KOLDING, S., 1986. Interspecific competition for mates and habitat selection in five species of Gammarus (Amphipoda: Crustacea). ____ *Mar. Biol.* **91**, 491-496.
- KONOPACKA, A. & K. JAŹDŹEWSKI, 1985. Stream ecosystems in mountain grassland (West Carpathians). 13. Gammarid species ____ *Acta hydrobiol.* **27**, 371-380. (Not seen).
- KOTRSCHAL, K. & D.A. THOMSON, 1986. Feeding patterns in eastern tropical Pacific blennioid fishes (Teleostei: Tripterygiidae, Labrisomidae, Chaenopsidae, Blenniidae). ____ *Oikos* **70**, 367-378.
- KRAFT, C. & J. F. KITSCHALL, 1986. Partitioning of food resources by sculpins in Lake Michigan (USA). ____ *Environm. Biol. Fishes* **16**, 309-316. (Pontoporeia important prey).
- KROON, H. de, H. de JONG & J.T.A. VERHOEVEN, 1985. The macrofauna distribution in brackish inland waters in relation to chlorinity and other factors. ____ *Hydrobiologia* **127**, 265-275. (Not seen. A Dutch study).
- KUDRYASHOV, V.A. & A. Yu. ZVYAGINTSEV, 1981. (Amphipoda (Crustacea) fouling ships in the Far Eastern marine basin). ____ Pp. 56-72 in V.A. KUDRYASHOV, A.A. KUBANIN & I.R. MIKHAILDV (ed.). (Fouling organisms of the Far Eastern Seas). Akad. Nauk SSSR, Vladivostok. (In Russian. Not seen, unfortunately).
- KUKERT, H., 1984. Die Crustaceen der Brackwassertümpel im Aussendeichsland zwischen Spieka- Neufeld und Arensch-Berersch/Cuxhaven und ihre Verteilung in Beziehung zum Salzgehalt (Crustacea: Cladocera, Copepoda, Amphipoda, Decapoda). ____ *Abh. naturwiss. Ver. Bremen* **40**, 115-130. (Not seen).
- KUVIKOVA, A., 1985. (On the food of the European water shrew, Neomys fodiens, in Slovakia). ____ *Biologia (Bratislava)* **40**, 563-572. (In German, not seen. An important predator of amphipods).
- LaFRANCE, K. & E. RUBER, 1985. The life cycle and productivity of the amphipod Gammarus mucronatus on a northern Massachusetts salt marsh. ____ *Limnol. Oceanogr.* **30**, 1067-1077.
- LAGZDINS, G. & A. SOULE, 1984. (On the biology of Bathyporeia pilosa Lindstrom in the littoral of the Gulf of Riga). ____ Pp. 217-227 in G.P. ANDRUSHAITIS (ed) (Hydrobiology of the Gulf of Riga). Zinatne, Riga, 237 pp. (In Russian; not seen. Can anyone get me a copy of this paper?).
- LALANA-RUEDA, R. & F. GOSSELCK, 1986. Investigation of the benthos of mangrove coastal lagoons in southern Cuba. ____ *Int. Rev. ges. Hydrobiol.* **71**, 779-794.
- LANTOS, G., 1986. Data on the Amphipoda and Isopoda fauna of Toserdo and its environs in the Tisza valley (Hungary). 1. Amphipoda, Asellota (Crustacea, Peracarida). ____ *Tiscia (Szeged)* **26**, 81-87. (Nipharqus mediodanubialis and Synurella ambulans).
- LARSEN, P.F., 1985. The benthic macrofauna associated with the oyster reefs of the St. James River estuary, Virginia, USA. ____ *Int. Rev. ges. Hydrobiol.* **70**, 797-814. (Not seen).
- LARSON, H.K., 1983. Notes on the biology of the goby Kelloggella cardinalis (Jordan & Seale). ____ *Micronesica* **19**, 157-164. (A predator of amphipods).
- LAZO-WASEM, E.A., 1985. Notes on the amphipod genus Idunella with special reference to Idunella bowenae Karaman. ____ *Crustaceana* **50**, 111-112.
- LEDOYER, M., 1986. Une nouvelle espèce de Pleustidae (Crustacea, Amphipoda) de la Méditerranée. ____ *Boll. Mus. Civ. Stor. Nat. Verona* **12** (1985), 511-513. (Pleustoides mediterraneus n. sp. from off Marseille).
- LEDOYER, M., 1986. Crustacés Amphipodes Gammariens 2. Familles des Haustoriidae a Vitjazianidae. ____ *Faune de Madagascar* **59**, 595-1112. (With this second part Ledoyer's monumental monograph on the amphipods of Madagascar is completed. The following taxa have been described and fully illustrated: Indischnopus herdmani, Pseudurothoe benthedij n. gen., n. sp. (Urothoidae), Urothoe elegans, U. serrulidactylus, Urothopsis brevicaudata, Cerapus abditus, C. cf. tubularis, Erichthonius brasiliensis, E. latimanus, E. pugnax, Ischyrocerus oakui armatus, Jassa falcata (of the 'form' marmorata), Parajassa bidentata, P. chilkoa, P. spinipalma, Ventojassa crenulata, V. ventosa, Leucothoe crenatipalma, L. ctenochir, L. dentata, L. euryonyx, L. hyhelia, L. laticoxa, L. lihue, L. madrasana, L. micronesiae, L. predenticulata, L. richiardi, L. r. macrodonta n. ssp., L. squalidens, Leucothoella bannwarthi, ?Leucothoides torrida, Leucothopsis angusticoxa, Idunella brevicornis (originally described as Ronconoides b.), Liljeborgia akaroica, L. bousfieldi, L. enigmatica n. sp., L. gloriosae n. sp. (3700 m), L. heeia, L. mojada, L. mozambica n. sp. (3370 m), Listriella cf. excavata, Acontistoma prionoplax, Amaryllis macropthalma, A. (Pseudamaryllis) nonconstricta (Pseudamaryllis thus reduced to rank of subgenus), Ambasiopsis brevipes n. sp., Aristias madagascarensis, A. stenopodus n. sp., A. symbioticus, Azotostoma fusca, ?Bathycallisma armata n. sp., Cyphocaris cornuta, C. faurei, C. geyserensis n. sp. (2300-2500 m), Dounialella n. gen. (Lysianassidae s. l.), type D. longichelata n. sp. (1800 m), Ensayara angustipes, E. micropthalma n. sp., Euonyx biscayensis, Hippomedon benthedij n. sp. (2500 m), ?H. brevicaudatus n. sp. (3900 m), H. normalis, H. onconotus, Ichonopus nossibeensis n. sp., I. pseudoserrius n. sp., I. spinicornis, Izinkala fihla, Kerguelenia macropoda n. sp., K. micropthalma n. sp. (3700 m), Lepidepecreella pamanzi n. sp. (2500 m), Lepidepecreum

- madagascarensis* n. sp. (recte: *madagascarensis*), *Lysianassa ceratina*, *L. cinghalensis*, *L. c. latipes* n. ssp., *L. ewa*, *L. nasuta*, *L. variegata*, *Onesimoides cavimanus*, *O. chelatus*, *Orchomene plicata*, *Procyphocaris induratus*, *Pseudocyphocaris* n. gen. (*Lysianassidae* s. l.), type *P. coxalis* n. sp., *Schisturella parachelata* n. sp. (3450 m), *Socarnes obesa*, *Socarnoides indentata* n. sp., *Thrombasia incerta* n. sp., *Trischizostoma denticulata*, ?*Uristes latipes* n. sp., *Maxillipius rectitelson*, *Melphisana madagascarensis*, *Ochlesis carinatus* n. sp., *Kanaloa manoa*, *Monoculodopsis longimana*, ?*Oedicerina megalopoda* n. sp., ?*Oediceroides plumicornis*, *O. cf. weberi*, ?*O. aff. wolffii*, *Periocolodes acuticoxa*, *P. aequimanus mozambicus* n. ssp., *P. brevicarpus* n. sp., *P. longimanus*, *P. megapleon*, *P. serra*, *Amathillopsis comorensis* n. sp. (2500 m), *A. septemdentata*, *Epimeria bispinosa* n. sp. (3450 m), *Halice macronyx*, *Pardaliscella inermis* n. sp. (3700 m), *Pereionotus alaniphlias*, *P. natalensis* (*Palinnotus* is synonymized with *Pereionotus*), *Plioplateia nodiformis* n. sp. (the *Plioplateiidae* are synonymized with the *Phliantidae*), *Basuto stimpsoni*, *Diogodias longicarpus*, *D. platyrostris*, *Harpiniopsis bandelei* n. sp. (520-830 m), *Pseudoharpinia cf. brevirostris*, *Harpiniopsis cf. capensis* (described as *Harpinia laeva capensis*), *Harpinia cf. curtipes*, *Harpiniopsis pseudonadania* n. sp. (1300-1480 m), *Joubinella indentata* n. sp. (1100 m), *Metaphoxoides angustimanus*, *M. picardi*, *Metaphoxus (Vasco) brevidactylus* (*Vasco* is thus reduced to the rank of subgenus), *M. fultoni tulearensis* (described as *Vasco tulearensis*), *Proharpinia setifera* n. sp., *Parapleustes barnardi*, *P. honomu*, *Dulichlopsis brevidactylus* n. sp., *Laetmatophilus hala*, *L. intermedius*, *Neoxenodice caprellinoides*, *Podocerus gloriosae* n. sp., *P. hanapepe*, *P. madagascarensis* n. sp., *P. palinuroides* n. sp., *P. tulearensis* n. sp., *P. walkeri pedunculata* n. ssp., ?*P. zeylanicus*, *Seba ekepuu*, *S. gloriosae* n. sp., *S. typica*, *Anadaniexis australis*, *A. tridentata* n. sp. (3700 m), *Glorandaniotis* n. gen., (*Stegocephalidae*), type *G. fissicaudata* n. sp. (3700 m), *Parandaniexis inermis* n. sp. (3700 m), *Stegocephaloides australis*, *Proboloides anophthalma* n. sp., *P. armata* n. sp. (3700 m), *Stenothoe adhaerens*, *S. gallensis*, *S. inermis*, *S. valida*, *Wallametopa cabon*, *Bruzelia diodon*, *Illeustroie illegetes*, *Metatiron brevidactylus*, *M. caecus*, *Synopia ultramarina*, *S. variabilis*, *Hyale chevreuxi*, *H. honoluluensis*, *H. inermis*, *H. macrodactyla*, *H. nigra*, *Orchestia anomala*, *O. notabilis*, *Parhyale hawaiiensis*, *P. spec.*, *Talorchestia martensii*, *Tulearus thomassini*, and *Vemana geyserensis* n. sp. (2500 m). In an appendix the following additional species are treated. *Byblis inaequicornis* n. sp., *Byblisoides* sp., *Biancolina mauihina*, *Photis dolichommata*, *Unciola integrileura* n. sp. (1100-1150 m), ?*Oradarea scissicaudata* n. sp. and *Prolaphystiopsis latirostris* n. sp. (2300-2500 m). A general part deals with the biogeography of Indian Ocean amphipods; it contains a list of all species recorded from this area (pp. 1046-1064).
- LEDOYER, M. & M. MENIOUL, 1983. Considerations sur la répartition du gammarien (Crustacea Amphipoda) *Jassa falcata* (Montagu, 1808). ____ Bull. Inst. scient. Rabat 7, 93-114.
- LEINEWEHER, P., 1985. The life-cycles of four amphipod species in the Kattgat. ____ Holarct. Ecol. 8, 165-174. (*Chaetogammarus marinus*, *C. stoerensis*, *Gammarus oceanicus* and *G. salinus*).
- LEITE, F.P.P., Y. WAKABARA & A.S. TARARAM, 1986. On the morphological variations in oostegites of gammaridean species (Amphipoda). ____ Crustaceana 51 77-94. (A most interesting study on the marsupial plates of 14 Brazilian intertidal amphipods).
- LEONG, T.S., Y.K. LI, S. C. HO, K.H. KHOO, S. P. KAM, S. HANAPI, T.M. WONG, R.S. LEGORE, W. de LIGNY & G.T. TAN, 1987. Effects of a crude oil terminal on tropical benthic communities in Brunei. ____ Mar. Poll. Bull. 18, 31-35. (Many amphipods).
- LEWIS, F.G., 1987. Crustacean epifauna of seagrass and macroalgae in Apalachee Bay, Florida, USA. ____ Mar. Biol. 94, 219-229.
- LOMBAS, I. & N. ANADON, 1985. (A study on the crevice fauna of intertidal zone of Luanco (Asturias), North of Spain) ____ Rev. Biol. Univ. Oviedo 3, 107-120. (In Spanish, not seen. Amphipoda?).
- LOWRY, J.K., 1986. The callynophore, a eucaridan/peracaridan sensory organ prevalent among the Amphipoda (Crustacea). ____ Zool. Scripta 15, 33-349. (The callynophore is the newly coined name for the sensory organ found on the fused proximal antennular flagellar articles in many amphipods. The organ is probably of chemosensory nature).
- LOWRY, J.K. & S. RUFFO, 1986. The rediscovery of *Lysianassa costae* H. Milne Edwards from the Gulf of Naples (Crustacea Amphipoda Lysianassidae). ____ Boll. Mus. civ. St. nat. Verona 11 (1984), 205-216. (Redescription and establishment of a neotype for this, the type species of *Lysianassa*. *L. costae* is exclusively Mediterranean, while the Atlantic *L. ceratina*, in spite of many published records, probably does not occur in the Mediterranean).
- LOWRY, J.K. & H.E. STODDART, 1986. Protandrous hermaphrodites among the lysianassoid Amphipoda. ____ J. crust. Biol. 6, 742-748.
- LUPKES, G. & H.G. MÜLLER, 1985. Crustaceen als Lebensformtypen des Grundwassers. ____ Mikrokosmos 74 (3), 71-76.
- MACDONALD, J. S., & R.H. GREEN, 1986. Food resource utilization by five species of benthic feeding fish in Passamaquoddy Bay, New Brunswick. ____ Can. J. Fish. Aquat. Sci. 43, 1534-1546.
- MACQUART-MOULIN, C., 1985. Le controle des phases pelagiques nocturnes chez les crustacés peracarides benthiques. ____ Tethys 11, 275-287.
- MAGNHAGEN, C., 1985. Random prey capture or active choice? An experimental study on prey size selection in three marine fish species. ____ Oikos 45, 206-216. (Prey on *Corophium volutator*).
- MAGUIRE, G.B., P. J. GIBBS & L.C. COLLETT, 1984. The macrobenthic fauna of brackish water prawn farming ponds at Port Stephens, New South Wales. ____ Austr. Zool. 21, 445-458. (Not seen. Amph. ?).
- MANNING, R.B., 1986. A small trap for collecting crustaceans in shallow waters. ____ Proc. biol. Soc. Wash. 99, 266-268.
- MARCHANT, R., 1981. The ecology of *Gammarus* in running water. ____ Pp. 225-249 in M.A. LOCK & D.D. WILLIAMS (eds). Perspectives in running water ecology. Plenum Press.

- MARSDEN, I.D., 1985. Some factors affecting survival and oxygen uptake in a subtropical beach flea. ____ J. exp. mar. Biol. Ecol. **88**, 213-225. (*Chroestia lota*).
- MASON, C.F., 1986. Invertebrate populations and biomass over four years in a coastal, saline lagoon. ____ Hydrobiologia **133**, 21-30.
- MATHIEU, J. & D. MARTIN, 1986. (Growth by weight and mortality of the subterranean amphipod *Nipharqus rhenorhodanensis* reared under different experimental conditions). ____ Bull. mens. Soc. linn. Lyon. (In French, not seen).
- MATHIEU, J. & D. MARTIN & P. HUISSOUD, 1984. Influence des conditions hydrologiques sur l'évolution de la structure spatiale et de la démographie d'une population phréatique de l'amphipode *Nipharqus rhenorhodanensis*. Premiers résultats. ____ Mem. biospeol. **11**, 27-37.
- MAZE, R.A. & A.J. LABORDA, 1986. (Some aspects of the distribution of the intertidal amphipods of the Playa de Area Longa, Ria del Barquero, Lugo (NW Spain)). ____ Actas 8. Jornadas A & E, Sevilla, 156-166. (In Spanish. Data on *Haustorius arenarius*, *Bathyporeia pelagica*, *Urothoe brevicornis*, *U. poseidonis*, *Talitrus saltator*, *Talorchestia brito*, *T. deshayesii* and *Gammarus locusta*. Haustoriids identified by WV).
- MCCAIN, J.C., 1984. Marine ecology of Saudi Arabia. The intertidal infauna of the sand beaches in the northern area, Arabian Gulf, Saudi Arabia. ____ Fauna Saudi Arabia **6**, 53-78.
- MCCAIN, J.C., 1984. Marine ecology of Saudi Arabia. The nearshore, soft-bottom benthic communities of the northern area, Arabian Gulf, Saudi Arabia. ____ Fauna Saudi Arabia **6**, 79-101. (Amph., identified by Barnard & Thomas, listed on p. 98; Three caprellids are listed separately).
- MCDONALD, J.H. Size-related and geographic variation at two enzyme loci in *Megalorchestia californiana* (Amphipoda: Talitridae). ____ Heredity **54**, 359-366.
- McLACHLAN, A., 1985. The biomass of macro- and interstitial fauna on clean and wrack-covered beaches in western Australia. ____ Est. coast. Shelf Sci. **21**, 587-599. (*Allochrestes compressus* dominates macrofauna).
- McLACHLAN, A., 1985. The ecology of two sandy beaches near Walvis Bay. ____ Madoqua **14**, 155-163.
- MEARNS, A.J., R.C. SWARTZ, J.M. CUMMINS, P.A. DINNELL, P. PLESHA & P.M. CHAPMAN, 1986. Inter-laboratory comparison of a sediment toxicity test using the marine amphipod *Rhepoxynius abronius*. ____ Mar. environm. Res. **19**, 13-38.
- MEISSEL, W., U. MATTIAS & ZIMMERMANN, 1985. Ecophysiological studies on acid tolerance of *Gammarus fossarum*. ____ Arch. Hydrobiol. **104**, 287-302.
- METCALFE, N., 1985. Prey detection by intertidally feeding Lapwing. ____ Z. Tierpsychol. **67**, 45-57.
- MEURS, H-G, 1985. Untersuchungen zum Vorkommen und zum Lebenszyklus euryhaliner Gammariden (Amphipoda, Crustacea) in der mixo-mesohalinen Zonen von Elbe, Weser und Ems. ____ Diplomarbeit Univ. Oldenburg, 100 pp.
- MEYER-ROCHOW, V.B., 1985. A study of unusual intracellular organelles and ultrastructural organization of the eye of *Gammarus oceanicus* fixed in the midnight sun of the Spitsbergen (Svalbard) summer. ____ Biomed. Res. **6**, 353-366.
- MEYER-ROCHOW, B.B. & T. SUZUKI, 1986. A study of visual pigments in the two antarctic crustaceans *Orchomene plebs* (Amphipoda) and *Glyptonotus antarctica* (Isopoda). ____ Comp. Biochem. Physiol. **83 B**, 75-78.
- MYERING, M.P.D., 1985. Einflüsse einem Ortslage auf die Verbreitung von *Gammarus*-Arten in einen Bachsystem (Maarbach, Osthessen). ____ Mitt. Ergänzungsstudium ökol. Umweltsicherung **10**, 125-141.
- MYERING, M.P.D. & H-G. PIEPER, 1985. Zur Verbreitung von *Gammarus* (Crustacea: Amphipoda) im Fulda- Eder- Abflussgebiet, mit besonderer Berücksichtigung der Bachversauerung. ____ Mitt. Ergänzungsstudium ökol. Umweltsicherung **10**, 91-123.
- MICHAELIS, F.B., 1985. Rare or threatened species from inland waters of Tasmania, Australia. ____ Rec. Queen Victoria Mus. **87**, 1-14. (Not seen. Includes amphipods).
- MILLER, D.C., 1981. A new video-computer system to quantify swimming behavior for toxicological studies. ____ Can. techn. Rep. Fish. aq. Sci. **990**, 69-71. (Not seen).
- MILLER, D.C., 1984. Mechanical post-capture particle selection by suspension- and deposit- feeding *Corophium*. ____ J. exp. mar. Biol. Ecol. **82**, 59-76.
- MITCHELL, D.G., J.D. MORGAN, G.A. VIGERS & P.A. CHAPMAN, 1985. Acute toxicity of mine tailings to four marine species. ____ Mar. Poll. Bull. **16**, 450-455. (i.a. *Rhepoxynius abronius*).
- MITSUKOVA, L.D., 1986. (Chemical composition and power value of *Gammarus lacustris* Sars). ____ Gidrobiol. Zh. **22**(1), 18-23. (In Russian, not seen).
- MLADENOV, P.V. & I. POWELL, 1986. A simple underwater magnifying device for the diving biologist. ____ Bull. mar. Sci. **38** 558-561.
- MÜLLER, P., L. PIHL & R. ROSENBERG, 1985. Benthic faunal energy flow and biological interaction in some shallow marine soft bottom habitats. ____ Mar. Ecol. Progr. Ser. **27**, 109-121.

- MOORE, J., 1984. Altered behavioural responses in intermediate hosts - an acanthocephalan host strategy. ____ Am. Natur. 123, 572-577.
- MOORE, P.G., 1986. Seaweed-associated animal communities in the Firth of Clyde, with special reference to the population biology of the amphipod *Hyale nilssonii* (Rathke). ____ Proc. roy. Soc. Edinburgh 90 B, 271-286.
- MOORE, P.G., 1986. A new species in the genus *Grandidierella* Coutière (Crustacea: Amphipoda) from an Australian solar salt works. ____ J. nat. Hist. 20, 1393-1399. *G. propodentata* sp. nov. from a hypersaline lagoon in Casuarina, Queensland).
- MOORE, P.G., 1987. Taxonomic studies on Tasmanian phytal amphipods (Crustacea): the families Anamixidae, Leucothoidae and Sebidae. ____ J. nat. Hist. 21, 238-262. (Deals with *Anamixis yarrega* (both *Anamixis* and *Leucothoides* -forms), *Leucothoe boolpooli*, *L. commensalis*, *L. clenochasma* n. sp., *L. neptunea* n. sp. and *Seba chiltoni* n. sp. All new species are from Tinderbox in the d'Entrecasteaux Channel, SE Tasmania).
- MOORE, P.G. & C.H. FRANCIS, 1985. Some observations on food and feeding of the supralittoral beach-hopper *Orchestia gammarellus* (Pallas) (Crustacea: Amphipoda). ____ Ophelia 24, 183-197.
- MOORE, P.G. & C.H. FRANCIS, 1985. On the water relations and osmoregulation of the beach-hopper *Orchestia gammarellus* (Pallas) (Crustacea: Amphipoda). ____ J. exp. mar. Biol. Ecol. 94, 131-150.
- MOORE, P.G. & C.H. FRANCIS, 1986. Notes on breeding periodicity and sex ratio of *Orchestia gammarellus* (Pallas) (Crustacea: Amphipoda) at Millport, Scotland. ____ J. exp. mar. Biol. Ecol. 95, 203-209.
- MOORE, P.G. & C.H. FRANCIS, 1986. Environmental tolerances of the beach-hopper *Orchestia gammarellus* (Pallas) (Crustacea: Amphipoda). ____ Mar. environm. Res. 19, 115-129.
- MOORE, P.G. & J.I. SPICER, 1986. On the status of *Arcitalitrus dorrieni* (Crustacea: Amphipoda) on the island of Colonsay, Inner Hebrides. ____ J. nat. Hist. 20, 667-680.
- MORGAN, E., G.J. HARRIS & W.F. HOLMSTROM, 1985. Circa-tidal pacemakers in the nervous system of an amphipod. ____ Pp. 177-180 in P.H. REDFERN, I.C. CAMPBELL, J.A. DAVIS & K.F. MARTIN (eds.) Circadian rhythms in the central nervous system. MacMillan Press Ltd, Basingstok & London.
- MORINO, H., 1985. Revisional notes on *Jesogammarus* - *Annanogammarus* group (Amphipoda - Gammaroidea) with description of four new spp. from Japan. ____ Publ. Itako hydrobiol. Stn. 2, 9-55. (A regional monographic study. *Annanogammarus* is considered to be only subgenerically different from *Jesogammarus*. The following taxa are fully described: *J. (J.) jesoensis*, *J. (A.) annandalei*, *J. (J.) hokurikuensis* n. sp. (Fukui prefecture), *J. (J.) spinopalpus* n. sp. (Ibaraki pr.), *J. (A.) naritai* n. sp. (L. Biwa, Shiga pr.) and *J. (A.) fluviatilis* n. sp. (L. Biwa, Shiga pr.)).
- MURDOCH, M.H., F. BARLÖCHER & M.L. LALTOO, 1986. Population dynamics and nutrition of *Corophium volutator* (Pallas) in the Cumberland basin (Bay of Fundy). ____ J. exp. mar. Biol. Ecol. 103, 235-249.
- MUSHKO, I.B., 1984. (Effect of the insecticide K-othrine on the epithelium of the hepatic caeca in the amphipod *Gammarus roeselii*). ____ Dokl. Akad. Nauk SSR 276, 504-507. (In Russian, not seen).
- MYERS, A.A., ?1985. Studies on the genus *Lembos* Bate. 11. *Globosolembos* sub-gen. nov. *L. (G.) francanni* Reid, *L. (G.) indicus* Ledoyer, *L. (G.) ovatus* sp. nov., *L. (G.) tiafaui* sp. nov., *L. (G.) excavatus* Myers. ____ Boll. Mus. Civ. St. nat. Verona 10 (1983), 341-367. (In addition to the spp. mentioned in the title, also *L. leapakahi*, *L. ruffoi* and *L. smithi* (type) belong to the new subgenus, characterized by enlarged female gnathopod 1. the type locality of *L. ovatus* is Viti Levu, Fiji, that of *L. tiafaui* Samoa).
- MYERS, A.A., ?1985. Studies on the genus *Lembos* Bate. 12. Tropical Pacific islands. *L. dentischium* Myers ssp. *taparum* nov., *L. saloteae* sp. nov., *L. waipio* Barnard, *L. aequimanus* Schellenberg, *L. virgus* sp. nov., *L. regius* sp. nov., *L. tui* sp. nov. ____ Boll. Mus. civ. St. nat. Verona 10 (1983), 369-406. (*L. d. taparum* and *L. saloteae* come from Tonga, *L. virgus* from Viti Levu, Fiji, *L. regius* from Vanuatu, N. Hebrides, and *L. tui* from W. Samoa).
- MYERS, A.A., ?1985. A new species of *Cheiriphotis* Walker from the Mediterranean Sea (Amphipoda: Isaeidae). ____ Boll. Mus. civ. St. nat. Verona 10 (1983), 541-542. (*C. mediterranea* n. sp. from the coast of Israel).
- MYERS, A.A., 1985. Shallow-water, coral reef and mangrove Amphipoda (Gammaridea) of Fiji. ____ Rec. austr. Mus. Suppl. 5, 1-143. (Deals with *Ampelisca melaniensis* n. sp., *A. pygmaea*, *Amphilochus menehune*, *Gitana bilobata*, *G. gracilis* n. sp., *Gitanopsis tai* n. sp., *Moolaphaeonoides coocoo seraa* n. ssp., *Amphithoe hirsuta* (was ssp. of *A. pollex*), *A. kava* n. sp., *A. kuala* n. sp., *A. ramondi*, *Cymadusa brevidactyla*, *C. imbroglio*, *C. lunata* n. sp., *Paragrubia vorax*, *Peramphithoe orientalis*, *Pleonexes kanehe navosa* n. ssp., *P. kulafi*, *Paranamixis madagascarensis*, *Aorella multiplex*, *Globosolembos ovatus*, *Grandidierella bispinosa*, *G. perlata*, *G. teres*, *Lembos virgus*, *Biancolina algicola*, *Colomastix lunallo*, *Corophium* sp., *Paradexamine rewa* n. sp., *Podocerus crenulatus* n. sp., *P. hanapepe*, *P. talegus levuensis* n. ssp., *Bircenna dronga* n. sp., *Eusiroides diplonyx*, *Hyale chevreuxi*, *H. galateae distorta* n. ssp., *H. maroubrae*, *Lelehua malevua* n. sp., *Parhyale hawaiiensis*, *Aorchoides dilatata*, *Chevalia aciculata*, *Gammaropsis atlantica*, *G. digitata*, *G. pacifica*, *G. pokipoki*, *G. setifera*, *G. togoensis*, *Photis kapapa*, *P. pirloti* n. sp., *Cerapus pacificus*, *Ventojassa ventosa*, *Leucothoe diemenensis*, *L. gavalis* n. sp., *Leucothoe bannwarthi*, *Parambasia nui* n. sp., *Parawaldeckia lowryi* n. sp., *Ceradocus rubromaculatus*, *Elasmopus lapu* n. sp., *E. molokoi*, *E. pseudaffinis*, *E. seticarpus* n. sp., *E. spinidactylus*, *E. spinimanus*, *Maera octodens*, *M. pacifica*, *M. quadrimana*, *M. serrata*, *Mallacoota insignis*, *M. nananui* n. sp., *M. subcarinata*, *Melita simplex* n. sp., *M. zeylanicus*, *Parelmopus suensis*, ?*Quadrivisio bousfieldi*, *Paracalliope mapela* n. sp., *Pereionotus alaniphias*, *Wildus parathambaroo* n. sp., *Stenothoe kaia* n. sp. and *Platorchestia platensis*. Key to all species are provided).

- MYERS, A.A., 1986. Amphipoda from the South Pacific: Tonga. ____ *Rec. austr. Mus.* **38**, 271-290. (Deals with Leucothoe hyhelia, Parawaldeckia mua n. sp., Elasmopus alalo n. sp. (= E. pseudaffinis auct. non Schellenberg), E. gracilis, E. molokai, Parelasomopus suensis, Mallacoota nananni, Eriopisella seychellensis, Cymadusa pilipes (= Paradusa bilobata pilipes, while C. lunata is a synonym) and Globosolembos excavatus. A key to all male Globosolembos is provided).
- MYERS, A.A. & M.J. COSTELLO, 1986. The amphipod sibling pair Leucothoe lilljeborgi and L. incisa in British and Irish waters. ____ *J. mar. biol. Ass. U.K.* **66**, 75-82.
- NAYLOR, C. & J. ADAMS, 1987. Sexual dimorphism, drag constraints and male performance in Gammarus duebeni (Amphipoda). ____ *Oikos* **48**, 23-27.
- NEALSON, K.H., A.C. ARNESON & M.E. HUBER, 1986. Identification of marine organisms using kinetic and spectral properties of their bioluminescence. ____ *Mar. biol.* **91**, 77-84.
- NEBEKER, A.N., M.A. CAIRNS, J.H. GAKSTATTER, K.W. MALUEG, G.S. SCHUYTEMA & D.F. KRAWCZYK, 1984. Biological methods for determining toxicity of contaminated freshwater sediments for invertebrates. ____ *Environm. Toxicol. Chem.* **3**, 617-630. (i.a. Hyaella azteca - 'a recommended benthic test organism' - and Gammarus lacustris).
- NELSON, W.G., 1986. Predation and prey population variation in a high energy sandy beach macrofaunal community. ____ *Ophelia* **26**, 305-316. (A study from the Atlantic coast of Florida, USA).
- NICHOLS, F.H. & J.K. THOMPSON, 1985. Time scales of change in the San Francisco Bay benthos. ____ *Hydrobiologia* **129**, 121-138.
- NOTENBOOM, J., 1986. Sensonator valentiensis n.g., n. sp. (Amphipoda) from different biotopes in southern Valencia. ____ *Bijdr. Dierk.* **56**, 60-74. (This new Spanish genus is difficult to place. The author concludes his discussion as follows: Sensonator is most probably a relict of an early free-swimming Gammaroidea- and/or Pardaliscoidea- like ancestral group. For the time being, the author refrains from erecting a new family).
- NOTENBOOM, J., 1986. The species of the genus Pseudoniphargus Chevreux, 1901 (Amphipoda) from northern Spain. ____ *Bijdr. Dierk.* **56**, 75-122. (Deals with P. longicarpus n. sp., P. semielongatus n. sp., P. montanus n. sp., P. elongatus, P. burgensis n. sp., P. eborarius n. sp., P. jereanus n. sp., P. sp., P. gorbeanus n. sp., P. vasconiensis n. sp., P. unisexualis, P. incantatus n. sp., P. guernicae n. sp. and P. spiniferus n. sp., all from northern Spain).
- O'CONNOR, J.M., J.H. SAMUELSON, K.J. SALAMON & J.C. PIZZA, 1985. Measurement of the effect of aroclor-1254 on the respiration of Gammarus using potentiometric respirometry. ____ *Water Res.* **19**, 639-643. (Not seen).
- OGI, H., H. TANAKA & T. TSUJITA, 1985. The distribution and feeding ecology of murrelets in the northeastern Bering Sea. ____ *J. Yamashina Inst. Ornithol.* **17**, 44-56. (Main prey is Parathemisto libellula).
- OLAFSSON, E.B. & L-E. PERSSON, 1986. The interaction between Nereis diversicolor O.F. Muller and Corophium volutator Pallas as a structuring force in a shallow brackish sediment. ____ *J. exp. mar. Biol. Ecol.* **103**, 103-117.
- OLIVER, J.S. & R.G. KVITEK, 1984. Side-scan sonar records and diver observations of the Gray Whale (Eschrichtius robustus) feeding grounds. ____ *Biol. Bull. (Woods Hole)* **167**, 264-269.
- OLIVER, J.S. & R.G. KVITEK & P.N. SLATTERY, 1985. Walrus feeding disturbance: scavenging habits and recolonization of the Bering Sea benthos. ____ *J. exp. mar. Biol. Ecol.* **91**, 233-246.
- OLIVER, J.S. & P.N. SLATTERY, 1985. Destruction and opportunity on the sea floor: effects of gray whale feeding. ____ *Ecol.* **66**, 1965-1975.
- OLIVER, J.S. & P.N. SLATTERY, 1985. Effects of crustacean predators on species composition and population structure of soft-bodied infauna from McMurdo Sound, Antarctica. ____ *Ophelia* **24**, 155-175. (A study of the influence of i.a. Heterophoxus videns, a predator, on the species composition and population size structure of soft-bodied infauna).
- OLIVER, J.S., P.N. SLATTERY, M.A. SILBERSTEIN & E.F. O'CONNOR, 1983. A comparison of Gray Whale, Eschrichtius robustus, feeding in the Bering Sea and Baja California. ____ *Fish. Bull.* **81**, 501-512.
- OMORI, K., 1985. (Life history of a mud-flat amphipod Corophium volutator - the adaptive significance of semi-lunar and Japanese).
- ORTIZ TOUZET, M., 1984. (Discovery of an ectoparasitic amphipod from a whale in Cuban waters) ____ *Rev. Invest. mar.* **5**, 95-96. (In Spanish, not seen. Cyamus boopis).
- OVERSTREET, R.M., 1983. Metazoan symbionts of Crustacea. ____ Pp. 155-250 in A.J. PROVENZANO (ed.) *The biology of the Crustacea Vol. 6*. Pathobiology. Acad. Press., N. York.
- PARDI, L., A. ERCOLINI, F. FERRARA, F. SCAPINI & A. UGOLINI, 1985. (Zonal solar and magnetic orientation in intertidal amphipod crustaceans of equatorial regions). ____ *Atti Accad. naz. Lincei R. (8)* **76**, 312-320. (In Italian; not seen).
- PEDERZANI, H.A., 1985. Fundort Aquarium- ein Amphipode, aber welcher? ---- *Mikrokosmos* **74**, 50-54. (Not seen, so I don't know).
- PEER, D.L., L.E. LINKLETTER & P.W. HICKLIN, 1986. Life history and reproductive biology of Corophium volutator (Crustacea: Amphipoda) and the influence of shorebird predation on population structure in Chignecto Bay, Bay of Fundy, Canada. ____ *Neth. J. Sea Res.* **20**, 359-373.

- PENNAK, R.W. & J.V. WARD, 1986. Interstitial faunal communities of the hyporheic and adjacent groundwater biotopes of a Colorado mountain stream. ____ Arch. Hydrobiol. Suppl. **74**, 356-396. (Amph., 2 *Stygobromus* spp., on pp. 375-376).
- PEREIRA, V.F. & C. GOULART, 1985. (Redescription of *Hyaella pernix* (Amphipoda, Hyaellidae) with a discussion of its synonym *Hyaella curvispina*) ____ Rev. brasil. Zool. **3**, 209-218. (In Portuguese, not seen. There are now according to the authors, 29 spp. of *Hyaella*).
- PERSSON, L.-E. & E.B. OLOFSSON, 1986. Distribution and abundance of mobile epifauna and macrozoobenthos in south Swedish shallow marine areas. ____ Ophelia, Suppl. **4**, 201-210.
- PINKNEY, A.E., G.V. POJE, R.M. SANSUR, C.C. LEE & J.M. O'CONNOR, 1985. Uptake and retention of 14 C- Aroclor 1254 in the amphipod, *Gammarus tigrinus*, fed contaminated fungus, *Fusarium oxysporum*. ____ Arch. environm. Contam. Toxicol. **14**, 59-64.
- PINKSTER, S. & D. PLATVOET, 1986. Remarks on the identity of *Echinogammarus thoni* (Schäferna, 1922) with description of a new species, *Echinogammarus cyrtus*, from southern France (Crustacea, Amphipoda). ____ Bull. zool. Mus. A'dam **10**, 173-181. (*E. thoni* is redescribed, *E. cyrtus* n. sp. (Dept. Hérault, France) described, and *E. antalyae* Karaman, described as ssp. of *thoni*, raised to specific rank).
- PISCITELLI, G., 1985. (Characteristics of the reproductive cycle of *Echinogammarus olivii* (H. Milne Edwards, 1830) (Amphipoda Gammaridae) of the littoral zone of Puglia). ____ Olbalia (N.S.) **11**, 833-836. (In Italian; not seen).
- PISCITELLI, G. & L. SCALERA LIACI, 1985. (Characteristics of the reproductive cycle of *Echinogammarus olivii* (H. Milne Edwards, 1830) (gammarid amphipod) of the littoral zone of Puglia). ____ Olbalia **11**, 833-836. (In Italian; not seen).
- PLATVOET, D., 1985. Side line organs in gammarids (Crustacea, Amphipoda). ____ Beaufortia **35**, 129-133.
- PLATVOET, D. & S. PINKSTER, 1985. The present position of the alien amphipods *Gammarus tigrinus* and *Crangonyx pseudogracilis* in the Netherlands (Crustacea, Amphipoda). ____ Bull. zool. Mus. A'dam **10**, 125-128.
- POLIKARPOV, G.G., V.G. TSYTSUGINA, V.I. TIMOSICHUK, N.V. DEMINA & N.N. TERESHCHENKO, 1985. (Toxicity of Black Sea deep water for benthos amphipods *Gammarus olivii*). Dokl. Akad. Nauk Ukrain, SSR, Ser. B Geol. Khim. Biol. Nauk 1985 (**8**), 74-76. (In Russian, not seen).
- PONYI, J.E. & L. BANKÓ, 1985. The effect of different pesticides on the amphipod *Gammarus roeselii* Gervais. ____ Misc. zool. hung. **3**, 83-90.
- POOVACHIRANON, S., K. BOTO & N. DUKE, 1986. Food preferences and ingestion rate measurements of the mangrove amphipod *Parhyale hawaiiensis* (Dana). ____ J. exp. mar. Biol. Ecol. **98**, 129-140.
- PRESLER, P., 1986. Necrophagous invertebrates of the Admiralty Bay of King George Island (South Shetland Islands, Antarctica). ____ Pol. polar Res. **7**, 25-61. (An important study, with data on *Cheirimeton femoratus*, *Hippomedon kergueleni*, *Orchomene plebs*, *O. rotundifrons* and *Waldeckia obesa*).
- PRIDMORE, R.C. & D.S. ROPER, 1985. Comparison of the macroinvertebrate faunas of runs and riffles in three NZ streams. ____ N.Z. J. mar. Freshw. Res. **19**, 283-291.
- RAFFAELLI, D., 1987. An experimental investigation of the effects of shorebird and flatfish predation on estuarine invertebrates. ____ Est. coast. Shelf sci. **24**, 1-13.
- RAFFAELLI, D. & P.R. BOYLE, 1986. The intertidal macrofauna of Nigg Bay. ____ Proc. roy. Soc. Edinb. **91** B, 113-141.
- RAGA, J.A. & C. SANPERA, 1986. (Ectoparasites and epizotes of *Balaenoptera physalus* (L. 1758) in Atlantic Therian waters). ____ Invest. Pesq. **50**, 489-498. (In Spanish i.a. *Cyamus balaenopterae*, p. 495).
- RAINBOW, P.S. & P.G. MOORE, 1986. Comparative metal analyses in amphipod crustaceans. ____ Hydrobiologia **141**, 273-289.
- RAPPÉ, G., 1985. (*Isocyamus delphinii* (Guérin, 1836), first record of a whale louse (Amphipoda, Cyamidae) from our coast). ____ De Strandvl (Belgium) **5**, 63-65. (In Flemish. First Belgian record).
- RAUSCHERT, M., 1985. *Eurythenes gryllus* (Liechtenstein) (Crustacea, Amphipoda) in der marinen Fauna von King George (Süd-Shetlandinseln, Antarktis). ____ Milu **6**, 319-324.
- REGNAULT, M., 1987. Nitrogen excretion in marine and fresh-water Crustacea. ____ Biol. Rev. **62**, 1-24. (A review paper).
- REICHERT, W.L., B.-T. le EBERHART & U. VARANASI, 1985. Exposure of two species of deposit-feeding amphipods to sediment-associated (3 H) benzo(a)pyrene: uptake, metabolism and covalent binding to tissue macromolecules. ____ Aq. Toxicol. **6**, 45-56. (not seen).
- REINHARDT, S.B. & E.S. VAN VLEET, 1986. Lipid composition of twenty-two species of Antarctic midwater zooplankton and fish. ____ Mar. Biol. **91**, 149-159. (i.a. *Cyphocaris richardi*, *Eurythenes gryllus*, *Eusirus propeperdentatus*, *Parandania boeckii* and *Parathemisto gaudichaudii*).
- RICHARDSON, A.M.M. & H.P. MORTON, 1986. Terrestrial amphipods (Crustacea, Amphipoda, family Talitridae) and soil respiration. ____ Soil Biol. Biochem. **18**, 197-200.

- RIDLEY, M. & D.J. THOMPSON, 1986. Sexual selection of population dynamics in aquatic Crustacea. ____ Pp. 409-422 in R.M. SIBLY & R.H. SMITH (eds.) Behavioural ecology. Blackwell, Oxford. (Not seen).
- RIETSMA, C.S., I. VALIELA & A. SYLVESTER-SERIANNI, 1982. Food preferences of dominant salt marsh herbivores and detritivores. ____ P.S.Z.N. 1: Mar. Ecol. 3, 179-189. (i.a. *Orchestria grillus*).
- RIMET, M., 1983. A comparative study of the effects of white light and ultraviolet rays on *Gammarus pungenis*. Part 1. ____ Annls Zool., Agra 20, 87-92.
- ROBERTSON, C.H. & R.W.G. WHITE, 1986. Feeding patterns of *Nesogobius* sp., *Gymnapistes marmoratus*, *Neodax balteatus* and *Acanthaluteres spilomelanurus* from a Tasmanian seagrass meadow. ____ Austr. J. mar. Freshw. Res. 37, 481-490.
- ROBINSON, B.W. & R.W. DOYLE, 1985. Trade-off between male reproduction (amplexus) and growth in the amphipod *Gammarus lawrencianus*. ____ Biol. Bull. (Woods Hole) 168, 482-488.
- RODRIGUEZ CASTELO, E. & J. MORA, 1984. (Introduction to the study of the dynamics of benthic populations of infralittoral muds with organic enrichment (Ria de Pontevedra, NW Spain)). ____ Cuadern. Area Sci. mar. Seminario Est. Galegas 1, 291-302. (In Spanish).
- RODRIGUEZ CASTELO, E. & J. MORA, 1984. (Population dynamics of infralittoral sand bottoms of the Ria de Pontevedra (NW Spain)). ____ Actos 4. Symp. iberico Est. Benthos mar., Lisboa, 13-22. (In Spanish).
- ROSTILLON, D., 1985. Seasonal variations in the benthos of a chalk trout stream, the river Samson, Belgium. ____ Hydrobiologia 126, 253-262. (i.a. *Gammarus forssarum*).
- ROWE, G.T., M. SIBUET & A. VANGRIESHEIM, 1986. Domains of occupation of abyssal scavengers inferred from baited cameras and traps on the Demerara abyssal plain. ____ Deep-Sea Res. 33, 501-522.
- RUFFO, S., 1984. *Bogidiella nubica* n. sp. from interstitial waters of the Sudan (Crustacea: Amphipoda). ____ Hydrobiologia 110, 131-134.
- RUFFO, S., 1985. (New mesopsammic Amphipoda from Andaman Islands). ____ Boll. Mus. civ. St. nat. Verona 10 (1983), 485-509. (In Italian. Describes and illustrates *Seborgia schieckei* n. sp., *Eriopisella chierogoi* n. sp. and *Josephosella andamana* n. gen., n. sp. (Melitidae), all from S. Andaman. In an appendix, the rediscovery of *Bollegidia sootai* on the coast of Malaysia is recorded).
- RUFFO, S., 1985. A new member of the Amphipoda suborder Ingolfiellidea from Namibia: *Stygobarnardia caprellinoides*, new genus, new species). ____ Atti Soc. ital. Sci. nat. Mus. civ. St. nat. Milano 126, 43-53. (In Italian, not seen. The new taxon is based on the supposed males of *Ingolfiella opisthodoris* K.H. Barnard, 1966).
- RUFFO, S., 1985. (Studies on amphipod crustaceans. 102. The Mediterranean species of the genera *Tmetonyx* Stebbing and *Tryphosella* Bonnier (Crustacea, Amphipoda, Lysianassidae). ____ Ann. Mus. civ. St. nat. Genova 85. ____ (In Italian. Deals with *Tmetonyx nardonis* (Heller) revived (= *T. exiguus*), *T. similis*, *Tryphosella minima*, *T. similima* n. sp. and *T. longidactyla* n. sp. (both from the Golfo di Napoli), *T. nanoides*, *T. caecula* and *T. palpiserrata*, the last two here transferred from *Tmetonyx*).
- RUFFO, S., 1986. The systematic status of *Orchomenella dilatata* Chevreux, 1903 (Crustacea, Amphipoda, Lysianassidae). ____ Boll. Mus. Civ. Stor. Nat. Verona 12 (1985), 525-528. (This is a *Tryphosella*. *Tmetonyx palpiserrata* is a junior synonym).
- RUFFO, S., 1986. Amphipoda; groupe *Metacrangonyx*. ____ Pp. 550-552 in L. BOTOSANEANU (ed.). Stygofauna mundi. E.J. Brill, Leiden (*Metacrangonyx* and *Pygocrangonyx* are intermediary between the Hadziidae and the Crangonyctidae).
- RUFFO, S., 1986. Amphipoda: Salentinellidae. ____ Pp. 564-566 in L. BOTOSANEANU (ed.). Stygofauna mundi. E.J. Brill, Leiden.
- RUFFO, S., 1986. Amphipoda: Talitroidea. ____ Pp. 571-573 in L. BOTOSANEANU (ed.). Stygofauna mundi. E.J. Brill, Leiden.
- SAALEMAA, H., 1986. Karyology of the northern Baltic peracaridan Crustacea. ____ Sarsia 71, 17-25. (All *Gammarus* spp. in the area have n = 26 (except n = 27). *Calliopius* has n=9, *Corophium volutator* n = 14 and *Leptocheirus pilosus* 2n = 22. Also *Pontoporeia femorata* has n = 14, but *P. affinis* is presumably polyploid, with n = 26).
- SAINTE-MARIE, B., 1986. Effect of bait size and sampling time on the attraction of the lysianassid amphipods *Anonyx sarsi* Steele & Brunel and *Orchomenella pinquus* (Boeck). ____ J. exp. mar. Biol. Ecol. 99, 63-77.
- SAINTE-MARIE, B. & P. BRUNEL, 1985. Suprabenthic gradients of swimming activity by cold-water gammaridean amphipod Crustacea over a muddy shelf in the Gulf of St. Lawrence. ____ Mar. Ecol. (Progr. Ser.) 23, 57-69.
- SALMAN, S.D., 1985. *Stenothoe irakiensis*, a new species of stenothoid amphipod from the Arabian Gulf. ____ Crustaceana 49, 244-250.
- SALMAN, S.D., 1986. *Parhyale basrensis*, a new species of talitrid amphipod from the Shutt Al-Arab region, Iraq. ____ Crustaceana 50, 286-294.
- SAMARAS, W.F. & F.E. DURHAM, 1985. Feeding relationship of two species of epizoic amphipods and the gray whale, *Eschrichtius robustus*. ____ Bull. S. Calif. Acad. Sci. 84, 113-126 (*Cyamus scammoni* and *C. ceti* feed on the irritated epidermis surrounding the ectocommensal barnacle *Cryptolepas rhachianecti*. The food niche of *Cyamus kessleri* remains unknown).

- SANDLUND, O.T., G. KJELLBERG & G. NOHEIM, 1987. (Mercury in fish and invertebrates in Lake Mjøsa.) ____ Fauna, Oslo 40, 10-15. (In Norwegian. Deals with i.a. Pallasea quadrispinosa and Gammaracanthus lacustris).
- SARVALA, J., 1986. Interannual variation of growth and recruitment in Pontoporeia affinis (Crustacea: Amphipoda) in relation to abundance fluctuations. ____ J. exp. mar. Biol. Ecol. 101, 41-60.
- SCAPINI, F., 1986. Inheritance of solar direction finding in sandhoppers. 4. Variation in the accuracy of orientation with age. ____ Monit. Zool. ital. 20, 53-62.
- SCHNEPPENHEIM, R. & R. WEIGMANN-HASS, 1986. Morphological and electrophoretic studies of the genus Themisto (Amphipoda: Hyperideae) from the South and North Atlantic. ____ Polar Biol. 6, 215-226.
- SCHODEL, H., 1985. Epizoische Einzeller auf Flohkrebse. ____ Mikrokosmos 74, 225-230.
- SCHODEL, H., 1985. Epizoische Einzeller auf Flohkrebse. 2. Besiedler der Gammaridenbeine. ____ Mikrokosmos 74, 269-273.
- SCHOLTZ, C., 1984. Untersuchungen zur Bildung und Differenzierung des postnauplialen Keimstreifs von Neomysis integer Leach (Crustacea, Malacostraca, Peracarida). ____ Zool. Jahrb. (Anat.) 112, 295-349. (The discussion deals with many problems of general Peracaridan interest).
- SCONFIETTI, R., 1984. (Contribution to the ecology of amphipods, isopods and tanaids: observations on some coastal lagoons in Sardinia.) ____ Rendiconti Semin. Fac. Sci. Univ. Cagliari 54 suppl., 59-67. (In Italian; not seen).
- SCHRIMPF, E. & F. FOEHLER, 1985. Gammarids in streams of northeastern Bavaria, FRG. 1. Prediction of their general occurrence by selected hydrochemical variables. ____ Arch. Hydrobiol. 103, 479-495.
- SCHWEDHELM, E., 1984. Ein neues Sinnesorgan bei Gammarus. ____ Naturwiss. 71, 51-52. (Not seen).
- SHEADER, M., 1985. Feeding in hyperiid amphipods. ____ Porcupine Newsl. 3 (3), 59-62. (Not seen).
- SHEADER, M., 1986. Primno evansi n. sp. (Amphipoda: Hyperideae) from the eastern North Atlantic. ____ J. nat. Hist. 20, 975-980. (W. and N of Madeira).
- SIEGFRIED, C.A., 1985. Life history, population dynamics and production of Pontoporeia hoyi (Crustacea, Amphipoda) in relation to the trophic gradient of Lake George, New York. ____ Hydrobiologia 122, 175-180.
- SIEGISMUND, H.R., 1985. Genetic studies of Gammarus: 2. Geographical variation at polymorphic enzyme loci in Gammarus salinus and Gammarus oceanicus. ____ Hereditas 102, 15-23.
- SIEGISMUND, H.R., 1985. Genetic studies of Gammarus: 3. Inheritance of electrophoretic variants of the enzymes mannose phosphate isomerase and glucose phosphate isomerase in Gammarus oceanicus. ____ Hereditas 102, 25-31.
- SIEGISMUND, H.R., V. SIMONSEN & S. KOLDING, 1985. Genetic studies of Gammarus. 1. Genetic differentiation of local populations. ____ Hereditas 102, 1-13.
- SIMPSON, K.W., J.P. FAGNANI, D.M. DeNICOLA & R.W. BODE, 1985. Widespread distribution of some estuarine crustaceans in the limnetic zone of the Lower Hudson River, New York. ____ Estuaries 8, 373-380. (i.a. Gammarus tigrinus).
- SKADSHIM, A. & H.R. SIEGISMUND, 1986. Genetic relationships among north-western European Gammaridae (Amphipoda). ____ Crustaceana 51, 163-175.
- SKET, B., 1985. Bogidiella gammariformis, new species (Amphipoda) from Ecuador. ____ Biol. Vestn. 33, 81-88.
- SLATTERY, P.N., 1985. Life histories of infaunal amphipods from subtidal sands of Monterey Bay, California. ____ J. crust. Biol. 5, 635-649. (Rhepoxynius abronius, R. fatigans and Eohaustorius sencillus).
- SLATTERY, P.N. & J.S. OLIVER, 1986. Scavenging and other feeding habits of lysianassid amphipods (Orchomene spp.) from McMurdo Sound, Antarctica. ____ Polar Res. 6, 171-178.
- SOLDATOVA, I.N., 1986. Ecophysiological properties of Pontogammarus maeoticus (Amphipoda) in a salinity gradient. ____ Mar. Biol. 92, 115-123.
- SOURENIAN, B., 1985. Le benthos littoral d'El Dabaa (Méditerranée, Egypt). 4. Premières données sur la macrofaune des sables infralittoraux. ____ Rapp. Comm. int. Mer Médit. 29, 253-255. (Not seen).
- SPEHAR, R.L., H.P. NELSON, M.J. SWANSON & J.W. RENOOS, 1985. Pentachlorophenol toxicity to amphipods and fathead minnows at different test pH values. ____ Environm. Toxicol. Chem. 4, 389-397.
- SPICER, J.I. & A.C. TAYLOR, 1986. A comparative study of the gill area relationships in some talitrid amphipods. ____ J. nat. Hist. 20, 935-947.

- SPOEL, S. van der, 1984. The northern distribution border of Phronimidae (Crustacea: Amphipoda) in the north Atlantic. ____ Uttar Pradesh J. Zool. 4, 10-16. (Not seen).
- STANHOPE, M.J. & C.D. LEVINGS, 1985. Growth and production of *Eogammarus confervicolus* (Amphipoda: Anisogammaridae) at a log storage site and in areas of undisturbed habitat within the Squamish estuary, British Columbia. ____ Can. J. Fish. aq. Sci. 42, 1733-1740.
- STAROBOGATOV, Y.I., 1986. (The system of Crustacea). ____ Zool. Zh. 65, 1769-1781. (In Russian. A new 'simplified' classification, with very many new names.)
- STEELE, D.H., 1986. The genus *Anonyx* (Crustacea, Amphipoda) in the North Pacific and Arctic oceans: *Anonyx laticostae* group. ____ Can. J. Zool. 64, 2603-2623. (Consists of *A. schefferi* n. sp. (Unimak Isl., Alaska), *A. stegnegeri* n. sp. (Bering Isl. not *stegnegeri*?), *A. gurjanovai* n. sp. (St. Lawrence Isl., Alaska, recte: *gurjanovae*, *A. hurleyi* n. sp. (= *A. lilljeborgi* s. Hurley 1963, from S. Juan Isla., WA, USA), *A. multiarticulatus*, *A. petersoni* n. sp. (St. Paul Isl., Pribilofs), *A. affinis* (= *A. japonicus* s. Gurj. 62, non Gurj. 51), *A. lebedi* (originally described by Gurjanova as ssp. of *A. lilljeborgi*), *A. sculptifer*, *A. orientalis* (originally described as ssp. of *A. debruyni*) and *A. volkovi* (which does not really belong in this group).
- STEELE, D.H., R.G. COOPER & D. KEATS, 1986. Two corophioid amphipods commensal on spider crabs in Newfoundland. ____ J. crust. Biol. 6, 119-124. (*Gammaropsis inaequistylis* and *Ischyrocerus commensalis* on *Chionoecetes opilio* and *Hyas araneus*).
- STEELE, D.H. & V.J. STEELE, 1986. The cost of reproduction in the amphipod *Gammarus lawrencianus* Bousfield, 1956. ____ Crustaceana 51, 176-182.
- STEELE, V.J., 1986. Ultrastructure of paired coniform 9+0 sensory cilia: a new type in the organ of Bellonci of the marine amphipod *Gammarus setosus*. ____ Cell tissue Res. 245, 117-125.
- STEELE, V.J. & P.E. OSHEL, 1987. The ultrastructure of an integumental microtrich sensillum in *Gammarus setosus* (Amphipoda). ____ J. crust. Biol. 7, 45-59.
- STEELE, V.J. & D.H. STEELE, 1986. The influences of photoperiod on the timing of reproductive cycles in *Gammarus* species (Crustacea, amphipoda). ____ Am. Zool. 26, 459-467.
- STEPHENSON, M. & G.L. MACKIE, 1986. Lake acidification as a limiting factor in the distribution of the freshwater amphipod *Hyaella azteca*. ____ Can. J. Fish. aq. Sci. 43, 288-292. (*H. azteca* is a good indicator of lake acidification in Ontario).
- STOCK, J.H., 1985. Stygobiont Crustacea of the hadzioid group from Haiti. ____ Bijdr. Dierk. 55, 331-426. (Haiti has a very diverse amphipod stygofauna, with part of the new taxa upsetting the current attempts to establish a new classification of the Gammaridae s.l. This paper describes and illustrates 13 new taxa and revises the subgeneric taxonomy of *Metaniphargus*. It deals with *Crangoweckelia mixta* n. gen. n. sp., *C. spinicauda* n. sp., *Pintoweckelia grandis* n. gen. n. sp., *Bahadzia latipalpus* n. sp., *Zombiweckelia parvipalpus* n. gen. n. sp., *Radaweckelia brevicauda* n. gen. n. sp. and *Apoweckelia serrata* n. gen. n. sp. *Metaniphargus* is divided into a number of subgenera as follows *Metaniphargus* (*C. curasavicus*, *C. orientalis*), *Guadzia* n. subgen. (*M. (G.) bullipes* (type), *M. (G.) juberthiei*, *M. (G.) crenatus* n. sp.), *Hispadzia* n. subgen. (*M. (H.) longidactylus* n. sp. (type), *M. (H.) chaetodactylus* n. sp.), *Haidzia* n. subgen. (*M. (H.) plumicauda* n. sp.), *Jamadzia* n. subgen. (*M. (J.) jamaicae* (type), *M. (J.) craterensis*, *Caribdzia* n. subgen. (*M. (C.) nicholsoni* (type), *M. (C.) palpator*, *M. (C.) bousfieldi*, *M. (C.) 1. longipes*, *M. (C.) 1. christophorensis*, *M. (C.) venezolanus*, *M. (C.) haitianus* n. sp., *M. (C.) hyporheicus* and *M. (C.) anchilhalinus*), and *Croidzia* n. subgen. (*M. (C.) beattyi*). All new taxa are from Haiti).
- STOCK, J.H., 1986. Deep sea origin of cave faunas: an unlikely supposition. ____ Stygologia 2, 105-111.
- STOCK, J.H., 1986. Two new amphipod crustaceans of the genus *Bahadzia* from 'blue holes' in the Bahamas and some remarks on the origin of the insular stygofaunas of the Atlantic. ____ J. nat. Hist. 20, 921-933. (*B. setimana* n. sp. from S. Andros, *B. obliqua* n. sp. from Cat Island. With key to all 5 species).
- STOCK, J.H., 1986. Order Amphipoda: an introduction. ____ Pp. 494-496 in L. BOTOSANEANU (ed.) Stygofauna mundi. E.J. Brill, Leiden.
- STOCK, J.H., 1986. Amphipoda: gammarid grouping (Gammaridae s. (ed.) Stygofauna mundi. E.J. Brill, Leiden.
- STOCK, J.H., 1986. Amphipoda: melitid grouping (Melitidae sensu Bousfield 1973, emend.) ____ pp. 504-513 in L. BOTOSANEANU (ed.) Stygofauna mundi. E.J. Brill, Leiden.
- STOCK, J.H., 1986. Amphipoda: Pleustidae. ____ Pp. 560-561 in L. BOTOSANEANU (ed.) Stygofauna mundi. E.J. Brill, Leiden (Here Stock places *Awacaris* and *Relictomaera*).
- STOCK, J.H., 1986. Amphipoda: Phoxocephalidae. ____ P. 562 in L. BOTOSANEANU (ed.) Stygofauna mundi. E.J. Brill, Leiden.
- STOCK, J.H., 1986. Amphipoda: Liljeborgiidae. ____ P. 563 in L. BOTOSANEANU (ed.) Stygofauna mundi. E.J. Brill, Leiden.
- STOCK, J.H., 1986. Amphipoda: Pardaliscidae. ____ P. 567 in L. BOTOSANEANU (ed.) Stygofauna mundi. E.J. Brill, Leiden.
- STOCK, J.H., 1986. Amphipoda: Aoridae. ____ P. 570 in L. BOTOSANEANU (ed.) Stygofauna mundi. E.J. Brill, Leiden.
- STOCK, J.H., 1986. Amphipoda: Ingolfiellidae. ____ Pp. 581-584 in L. BOTOSANEANU (ed.) Stygofauna mundi. E.J. Brill, Leiden.

- STOCK, J.H., T.M. ILIFFE & D. WILLIAMS, 1986. The concept 'anchialine' revisited. ____ *Stygologia* **2**, 91-93.
- STOCK, J.H. & B.L.M. RONDE-BROEKHUIZEN, 1986. Stygo fauna of the Canary Islands. 1. A new species of *Pygocrangonyx*, an amphipod genus with African affinities, from Fuerteventura. ____ *Bijdr. Dierk.* **56**, 247-266. (The taxonomy of the *Metacrangonyx* group is discussed, with i.a. a cladistic analysis. *P. repens* n. sp. was discovered in wells on the island of Fuerteventura).
- STOCK, J.H. & J.J. VERMEULEN, 1985. Crustacés amphipodes stygobies de la Guadeloupe en relation avec l'histoire géologique de l'île. ____ *Mem. Biospeol.* **11** (1984), 265-273. (*Metaniphargus juberthiei* n. sp. was the only amphipod found).
- STRETCH, J.J., 1985. Quantitative sampling of demersal zooplankton: reentry and airlift dredge sample comparisons. ____ *J. exp. mar. Biol. Ecol.* **91**, 125-136. (A study using demersal amphipods from Catalina Isl., California).
- STUART, V., E.J.H. HEAD & K.H. MANN, 1985. Seasonal changes in the digestive enzyme levels of the amphipod *Corophium volutator* (Pallas) in relation to diet. ____ *J. exp. mar. Biol. Ecol.* **88**, 243-256.
- SUNDELIN, B., 1984. Single and combined effects of lead and cadmium on *Pontoporeia affinis* (Crustacea, Amphipoda) in laboratory soft-bottom microcosms. ____ Pp. 237-258 in G. PERSOONE, E. JASPERS & C. CLAUS (eds.). *Ecotoxicological testing for the marine environment*, Ghent 1984.
- SWANSON, S.A., 1984. Dissemination of amphipods by waterfowl. ____ *J. Wildl. Mgmt.* **48**, 988-991.
- SZANIANWSKA, A. & M. WOLOWICZ, 1986. Changes in the energy content of common species from Hornsund, southwest Spitsbergen (Arctic Ocean). ____ *Polar Res.* **4**, 85-90.
- TARARAM, A.S., Y. WAKABARA & F.P.P. LEITE, 1987. Vertical distribution of amphipods living on algae of a Brazilian intertidal rocky shore. ____ *Crustaceana* **51**, 183-187.
- TARARAM, A.S., Y. WAKABARA & H. de SOUSA LIMA MESQUITA, 1985. Feeding habits of *Hyale media* (Dana, 1853) (Crustacea-Amphipoda). ____ *Bolm Inst. oceanogr. S. Paulo* **33**, 193-199.
- TARR, J.G., C.L. GRIFFITHS & R. BALLY, 1985. The ecology of three sandy beaches on the Skeleton Coast of South West Africa. ____ *Madoqua* **14**, 293-304. (Not seen).
- TAYLOR, A.C. & J.I. SPICER, 1986. Oxygen-transporting properties of the blood of two semi-terrestrial amphipods, *Orchestia gammarellus* (Pallas) and *O. mediterranea* (Costa). ____ *J. exp. mar. Biol. Ecol.* **97**, 135-150.
- TAYLOR, P.M., 1985. Water balance in the estuarine crustacean *Corophium volutator* (Pallas) (Amphipoda). ____ *J. exp. mar. Biol. Ecol.* **88**, 21-19.
- TAYLOR, P.M. & R.R. HARRIS, 1986. Osmoregulation in *Corophium curvispinum* (Crustacea: Amphipoda), a recent colonizer of freshwater: 1. Sodium ion regulation. ____ *J. comp. Physiol. B. Biochem. Syst. environm. Physiol.* **156**, 323-330.
- TAYLOR, P.M. & R.R. HARRIS, 1986. Osmoregulation in *Corophium curvispinum* (Crustacea: Amphipoda), a recent colonizer of freshwater. II. Water balance and the functional anatomy of the antennary organ. ____ *J. comp. Physiol. B. Biochem. Syst. environm. Physiol.* **156**, 331-338.
- THOMAS, J.D. & J.L. BARNARD, 1986. New genera and species of the *Megaluropus* group (Amphipoda, Megalurotidae) from American seas. ____ *Bull. mar. Sci.* **38**. (The family Megalurotidae (n. fam?) contains the following American taxa: *Resupinus* n. gen., type *R. spinicaudatus* n. sp. (Belize), further spp. *Megaluropus visendus* and *R. coloni* n. sp. (Panama), and *Gibberulus* n. gen., type *Megaluropus longimerus* (from Nigeria, here redescribed) with american spp. *M. falciformis* (earlier described as ssp of *M. longimerus*), *M. myersi* and *G. devaneyi* n. sp. (La Jolla Calif.). Megalurotids feed like melphidippids, 'standing upside down on the substrate').
- THOMAS, J.D. & J.L. BARNARD, 1986. Two species of *Hornellia* (subgenus *Metaceradocus*) from the Florida keys and Belize (Amphipoda, Melphidippoidea). ____ *Bull. mar. Sci.* **38**, 477-487. (*H. (M.) tequestae* n. sp. from the Florida Keys and *H. (M.) atlanticus* (recte 'atlantica') n. sp. from Belize).
- THOMPSON, D.B.A., 1982. The abundance and distribution of intertidal invertebrates, and an estimation of their selection by Shelduck. ____ *Wildfowl* **33**, 151-158.
- THOMPSON, D.B.A., D.J. CURTIS & J.C. SMYTH, 1986. Patterns of association between birds and invertebrates in the Clyde estuary. ____ *Proc. roy. Soc. Edinburgh* **90** B, 185-201.
- THOMSON, D.H., C. M. MARTIN & W.E. CROSS, 1986. Identification and characterization of Arctic nearshore benthic habitats. ____ *Can. techn. Rep. Fish. Aq. Sci.* **1434**, 1-70.
- THURSTON, M.H., D.S.M. BILLETT & E. HASSACK, 1987. An association between *Exspina typica* Lang (Tanaidacea) and deep-sea holothurians. ____ *J. mar. biol. Ass. U.K.* **67**, 11-15. (One hundred sixty-four specimens of *Valetia* sp. were collected from the cloaca or intestine of deep-sea holothurians).
- TIGANUS, V., 1985. Sur la distribution bathymetrique des populations d'Amphipodes de substrats meubles des eaux roumaines de la Mer Noire. ____ *Rapp. Comm. int. Mer Méditerr.* **29**, 331-332.

- TSYTSUGINA, V.G., 1985. (Adaptation of the natural population of amphipods Gammarus olivii to the anthropogenic pollution of the environment). ____ dokl. Akad. Nauk SSSR 279, 1270-1271. (In Russian).
- TSYTSUGINA, V.G., 1985. (Comparative radiosensitivity of Gammarus olivii populations living in different ecological conditions). ____ Radiobiologiya 25, 812-815. (In Russian, not seen).
- TULLY, O. & O'CEIDIGH, 1986. The ecology of Idotea species (Isopoda and Gammarus locusta (Amphipoda) on surface driftweed in Galway Bay (West of Ireland). ____ J. mar. biol. Ass. U.K. 66, 931-942.
- TURQUIN, M.-J., 1983. Un amphipode de la faune du sol en France. ____ Pp 674-676 in Ph. LEBRUN et al. (eds). New trends in soil biology. (Niphargus rhenorhodanensis).
- TURQUIN, M.-J., 1984. Age et croissance de Niphargus virei (Amphipode pérennant) dans la système Karstique de Drom: methodes d'estimation. ____ Mém. Biospéol. 11, 37-49.
- TURQUIN, M.-J., 1986. Un cas de courbure pleonale teratologique chez Gammarus. ____ Bull. mens. Soc. linn. Lyon 55, 91-95.
- TURQUIN, M.-J., 1986. Mortalité et stock de l'amphipode Niphargus virei, dans le système karstique de Drom. ____ Bull. Soc. linn. Lyon 55, 293-304.
- TURQUIN, M.-J., 1986. La stratégie démographique de deux espèces d'amphipodes cavernicoles. ____ Colloque Biol. Popul. Lyon, not paginated.
- TURQUIN, M.-J., 1986. The relationship between the densities of a population of the troglobite Niphargus virei and the rainfall. Floods: malentities versus resources. ____ U.Z. 1. Rome.
- TZVETKOVA, N.L. & V.A. KUDRJASCHOV, 1985. (On the fauna and ecology of gammarids (Amphipoda, Gammaridea) in the biocoenoses of the upper part of the shelf of the south Sakhalin) ____ Akad. Nauk SSSR, Issled. Faun. Morei 30 (38), 292-345 (In Russian. Data on 74 species, of which 59 are new to the area, and Biancolina obtusata sachalinensis n. ssp. and Melita mikulitschae Gurjanova n. sp. are new taxa. Many qualitative and quantitative biological and synecological data are provided).
- UGOLINI, A., F. SCAPINI & L. PARDI, 1986. Interaction between solar orientation and landscape visibility in Talitrus saltator. ____ Mar. Biol. (Berl.) 90, 449-460.
- UZUNOV, Y.I. & S.G. KOVACHEV, 1985. Comparative estimation as determined by different methods of the production of Gammarus balcanicus Schaeff. ____ Hidrobiologiya 26, 75-80. (Not seen).
- VADER, W., 1985. Notes on Norwegian marine Amphipoda 9. Aristias megalops Sars, 1895 (Lysianassoidea) rediscovered. ____ Fauna norv. Serv. A 6, 1-2.
- VADER, W. & K. ROMPPAINEN, 1985. Notes on Norwegian marine Amphipoda. 10. Scavengers and fish associates. ____ Fauna norv. Ser. A. 6, 3-8.
- VADER, W., 1986. Menigratopsis svennilssoni (Amphipoda, Lysianassoidea) found in northern Norway. ____ Fauna norv. Ser. A 7, 47.
- VADON, C., 1984. La faune carcinologique associée à l'actinie Anemonia sulcata (Pennant) sur les côtes françaises de Méditerranée. ____ Oceanis 10, 551-555. (i.a. Phtisica marina, found on c. half of the sea anemones on one site, but absent in the other. Up to 100 amphipods were found on one host, mostly on the tentacles).
- VAN SENUS, P., 1985. The effects of temperature, size, season and activity on the metabolic rate of the amphipod, Talorchestia capensis (Crustacea, Talitridae). ____ Comp. Biochem. Physiol. A. Comp. Physiol. 81, 263-270.
- VAN SENUS, P. & A. McLACHLAN, 1985. Distribution and behavior of the amphipod Talorchestia capensis (Crustacea: Talitridae). ____ S. Afr. J. Zool. 20, 253-257.
- VAN SENUS, P. & A. McLACHLAN, 1986. Growth, production, and a partial energy budget for the amphipod, Talorchestia capensis (Crustacea: Talitridae) in the Eastern Cape, South Africa. ____ Mar. Ecol. 7, 165-180.
- VARANASI, K., W.L. REICHEL, J.E. STEIN, D.W. BROWN & H.R. SANBORN, 1985. Bioavailability and biotransformation of aromatic hydrocarbons in benthic organisms exposed to sediment from an urban estuary. ____ Environm. Sci. Technol. 19, 836-841 (i.a. Rhepoxynius abronius and Eohaustorius sencillus).
- VASSILENKO, S.V., 1985. (The role of oxygen consumption and size-weight characteristics of four caprellid species). ____ Biol. Morya, Vladivostok 1985-5, 40-45 (In Russian. Caprella cristibrachium, C. kroyeri, C. penantis and C. mutica).
- VILAS, F., 1986. Activity of amphipods in beach sediments and nearshore environments: Playa Ladeira, northwestern Spain. ____ J. Coast. Res. 2, 285-296.
- VINCENT, M., 1985. Etude comparée des températures lethales superieures et du preferendum thermique chez deux espèces d'amphipodes épigés du centre-ouest. ____ Annls scient. Limoges 1985-1, 43-47. (Not seen).
- VOGEL, F., 1985. Die abdominale Muskulatur von Orchestia cavimana Heller, 1865 (Amphipoda, Talitridae). ____ Crustacean 56, 11-26.

- WARD, P.I., 1986. A comparative field study of the breeding behavior of a stream and a pond population of Gammarus pulex (Amphipoda). ____ Oikos 46, 29-36.
- WARD, P.I., 1987. Sexual selection and body size in Gammarus pulex; a reply to Greenwood & Adams. ____ Oikos 48, 108-109.
- WILDISH, D.J. & M.J. DADSWELL, 1985. Sublittoral gammaridean amphipods of soft sediments in the Bay of Fundy. ____ Proc. N. Sc. Inst. Sci 35, 1-15 (Ninety-nine spp. of which two (Haploops fundiensis and a Melita) were previously undescribed, and 6 (Ampelisca abdita, Acanthohaustorius millsi, Gammaropsis maculatus, Menigrales obtusifrons, Melita nitida and Stenopleustes gracilis) constitute additions to the Canadian fauna).
- WILDISH, D.J., D.L. PEER & D.A. GREENBERG, 1986. Benthic macrofaunal production in the Bay of Fundy and the possible effects of a tidal power barrage at Economy Point - Cape Tenny. ____ Can. J. Fish. aq. Sci. 43, 2410-2417.
- WILLIAMS, L.G., P.M. CHAPMAN & T.C. GINN, 1986. A comparative evaluation of marine sediment toxicity using bacterial luminescence, oyster embryo and amphipod sediment bioassays. ____ Mar. environm. Res. 19, 225-249.
- WILLIAMS, W.D., 1984. Chemical and biological features of salt lakes of the Eyre peninsula, South Australia, and an explanation of regional differences in the fauna of Australian salt lakes. ____ Verh. Ver. theor. angew. Limnol. 22, 1208-1215. (Not seen).
- WILLIAMS, W.D., 1986. Amphipoda on land masses derived from Gondwana. ____ Pp. 553-559 in L. BOTOSANEANU (ed.). Stygofauna mundi. E.J. Brill, Leiden.
- WITEK, Z., W. KITTEL, H. CZYKIETA, M.I. ZMIJEWSKA & E. PRESLER, 1985. Macrozooplankton in the southern Drake Passage and in the Bransfield Strait (Antarctic) during BIOMASS SIBEX (Dec. 1983 - Jan. 1984). ____ Pol. polar Res. 6, 95-116.
- WOLDWICZ, M. & SZANIAWSKA, 1986. Calorific value, lipid content and radioactivity of common species from Hornsund, southwest Spitsbergen (Arctic Ocean). ____ Polar Res. 4, 79-84.
- WRIGHT, D.A., 1986. Trace metal uptake and sodium regulation in Gammarus marinus from metal polluted estuaries in England. ____ J. mar. biol. Ass. U.K. 66, 83-92.
- YAMASHITA, Y., D. KITAGAWA & T. AOYAMA, 1985. A field study of predation of the hyperiid amphipod Parathemisto japonica on larvae of the Japanese sand eel, Ammodytes personatus. ____ Bull. jpn Soc. scient. Fish. 51, 1599-1608.
- YAMATO, Sh., 1985. Discrimination of four intertidal melitid species (Amphipoda: Melitidae) in the Inland Sea of Japan, and evidence of their reproductive isolation. (Preliminary report). ____ Benthos Res. (Bull. jpn. Assoc. Benthol. 28)? (Not seen. Melita koreana s.l. is split into for taxa: M. koreana s. str., M. rylovae, and two undescribed spp. Breeding tests showed neither hybrids nor interspecific amplexing pairs).
- YOO, K-T, 1985. Zoogeography of genus Parathemisto (Amphipoda: Hyperiidea) in Korean waters. ____ Bull. mar. Sci. 37, 782 (Abstract only).
- ZEIDLER, W., 1986. Status of Australian crustacean type material described by K. Sheard 1936-39. ____ Crustaceana 51, 109-111 (Much of the material, esp. slides, is missing).
- ZHENG, Y., M.-t. LI & F.-q. TIAN, 1985. (An application of Corophium spp. (Crustacea, Amphipoda) in aquaculture of prawn). ____ Mar. Sci., Qingdao 9, 46. (In Chinese).

BIBLIOGRAPHY

July 1990

- AOKI, M., 1988. (Factors affecting population fluctuations of caprellid amphipods inhabiting Sargassum patens bed (Preliminary report). ____ Benthos Res. 32, 42-49. (In Japanese, English summary).
- AOKI, M. & T. KIKUCHI, 1990. Habitat adaptations of caprellid amphipods and the importance of epiphytic secondary habitats in a Sargassum patens bed in Amakusa, southern Japan. ____ Publ. Amakusa mar. biol. Lab. 10, 123-133.
- BARNARD, J.L. & A.Y. DAI, 1988. Four species of Gammarus (Amphipoda) from China. ____ Sinozoologia 6, 85-112. (Not seen. Deals with Gammarus lacustris, G. shanxiensis n. sp., G. hongyuanensis n. sp. and G. lasaensis n. sp. all in the pulex - group. Apparently contains much data on and references to freshwater amphipods of China).
- BARNARD, J.L. & J.D. THOMAS, 1990. Ensayara jumane, a new species from Belize, Caribbean Sea (Amphipoda, Lysianassidae). ____ Proc. biol. Soc. Wash. 103, 120-126. (With key to Ensayara spp).
- BLINN, D.W., B. DEHDASHTI, C. RUNCH & R.W. DAVIES, 1990. The importance of prey size and density in an endemic predator-prey couple (leech Ergobdella montezuma - amphipod Hyalella montezuma). ____ J. Anim. Ecol. 59, 187-192.
- BOTOSANEANU, L. & J. NOTENBOOM, 1988. Un amphipode stygobie de la Thaïlande (Crustacea, Amphipoda). ____ Annls. limnol. 24, 51-59. (Bogidiella (B.) thai n. sp. from a cave in S. Thailand).

- BOUSFIELD, E.L. & K.E. CONLAN. Malacostracans. ____ Pp. 854-859 in The new Encyclopaedia Britannica. Vol. 16, Macropaedia. Encyclopaedia Britannica, Inc., Chicago.
- BOUTIN, C. & M. MESSOULI, 1988. *Metacrangonyx gineti* n. sp. d' une source du Mont Atlas Marocain et la famille des Metacrangonyctidae n. fam. (Crustacés Amphipodes stygobies). ____ Vie Milieu **38**, 67-84. (*Pygocrangonyx* and *Afrocrangonyx* are merged with *Metacrangonyx*).
- BUSCHMANN, A.H., 1990. Intertidal macroalgae as refuse and food for amphipods in Central Chile. ____ Aquat. Bot. **36**, 237-245.
- CARLS, M.G. & S. KORN, 1985. Sensitivity of arctic marine amphipods and fish to petroleum hydrocarbons. ____ Can. techn. Rep. Fish. aq. Sci. 1368, 11-26. (*Anonyx nugax*, *Boeckosimus nanseni* and *Gammaracanthus loricatus*)
- CHAUDRI, M.A., A.A. GHURI & M.S. MAHOON, 1978. Aquatic fauna of Swat Valley, Pakistan. Part 3. Crustacea. ____ Biologie, Lahore **24**, 177-198. (*Anisogammarus madgensis* n.sp. (Madyen, NWFP, Pakistan) is described on pp. 181-183 and illustrated on pp. 193-194).
- CONLAN, K.E., 1990. Revision of the crustacean amphipod genus *Jassa* Leach (Corophioidea: Ischyroceridae). Can. J. Zool. **68**: 2031-2075. (The genus *Jassa* is revised to encompass the type species *J. falcata*, the previously recognized species *J. ingens*, *J. herdmani*, *J. pusilla*, and *J. marmorata*, and the new species *J. alonsoae*, *J. borowskyae*, *J. carltoni*, *J. fenwicki*, *J. gruneri*, *J. hartmannae*, *J. justii*, *J. morinai*, *J. myersi*, *J. oclairi*, *J. shawi*, *J. slatteryi*, *J. staudei*, and *J. thurstoni*. *Jassa odontonyx* is synonymized with *J. pusilla*, and *J. pulchella* is confirmed as synonymous with *J. falcata*. *Jassa wandeli*, *J. multidentata*, *J. goniamera*, *J. barnardi*, *J. lillipuna*, and *J. ocia* will be assigned to other genera in a later publication).
- DAUVIN, J.-C. & F. SENTIL, 1990. Conditions of the peracarid populations of subtidal communities in northern Brittany ten years after the Amoco Cadiz oil spill. ____ Mar. Poll. Bull. **21**, 123-130. (Most populations have recovered, but some formerly common species are still absent).
- DE GROVE, S. & C. d' UDEKEM d' ACOZ, 1988. (Observations concerning the marine fauna of northern France in 1987). ____ De Strandlo **8**, 86-99. In Dutch. Amph. pp 89-90, i.a. *Melita gladiosa*, *Gammaropsis nitida*, (in hermit-crab inhabited *Buccinum* shells), and *Corophium sextonae*.
- DEWITT, T. H., R.C. SWARTZ & J.O. LAMBERSON, 1989. Measuring the acute toxicity of estuarine sediments. ____ Environm. Toxicol. Chem. **8**, 1035-1048. (Tests with *Eohaustorius estuarius*).
- DICK, J.T.A., D.E. IRVINE & R.W. ELLWOOD, 1990. Differential predation by males on molted females may explain the competitive displacement of *Gammarus duebeni* by *Gammarus pulex* (Amphipoda). ____ Behav. Ecol. Sociobiol. **26**, 41-46. (This is *G. d. celticus*).
- FRICKE, H. & J. OEHLenschläger, 1988. Fatty acid and sterol composition of the Antarctic amphipod *Themisto gaudichaudii* Guérin 1828. ____ Comp. Biochem. Physiol. **89B**, 39-42.
- GARCIA, J.J. & N.B. CAMINO, 1987. (Preliminary studies on parasites on amphipods (Crustacea - Malacostraca) from Argentina). ____ Neotropica **33**, 57-64 (In Spanish. Diverse parasites in *Hyalella*).
- GARTNER, J.V. & J.A. MUSICK, 1989. Feeding habits of the deep-sea fish, *Scopelogadus beanii* (Pisces: Melamphaidae), in the Western North Atlantic. ____ Deep-Sea Res. **36A**, 1457-1470. *Vibilia*, *Parathemisto* and *Phronima* important prey).
- GLATZEL, T., 1989. (An inventory of the groundwater fauna in northwestern Lower Saxony, with special reference to Crustacea). ____ Drosera **89**, 11-22. (In German, not seen).
- GLEASON, L.N., 1989. Movements of *Pomphorhynchus bulbocollis* larvae from the hemocoel to the peripheral circulation of *Gammarus pseudolimnaeus*. ____ J. Parasitol. **75**, 982-985.
- GODDARD, J., 1989. Presumptive Batesian mimicry of a aeolid nudibranch by an amphipod crustacean. ____ Shells & Sea Life **16**, 220-222. (*Podocerus* sp. mimics *Flabellina trilineata*).
- HARGEBY, A., 1990. Macrophyte associated invertebrates and the effect of habitat permanence. ____ Oikos **57**, 338-346. (*Gammarus* and *Asellus* in stands of *Choraceae*).
- HAY, M.E., J.E. DUFFY & W. FENICAL, 1990. Host-plant specialization decreases predation on a marine amphipod: a herbivore in plant's clothing. ____ Ecology **71**, 733-743. (A study of *Pseudamphithoides incurvaria*).
- HOLSINGER, J., 1989. Preliminary zoogeographic analysis of five groups of crustaceans from anchialine caves in the West Indian region. ____ Proc. 10. Int. Congr. Speleol. **1**, 25-26. (Abstract only).
- HONG, J.-S. & D.J. REISH, 1987. Acute toxicity of cadmium to eight species of marine amphipod and isopod crustaceans from Southern California. ____ Bull. environm. Contom. Toxicol. **39**, 884-888.
- HUDSON, A.V. & J.D. REYNOLDS, 1977. Distribution of Irish intertidal Talitridae. ____ Bull. Ir. biogeogr. Soc. **2**, 63-76.
- JAŻDŹEWSKI, K., 1987. *Corophium multisetosum* Stock in the southern Baltic Sea - some notes on its morphology, ecology and biology. ____ Proc. 4 Symp. Baltic mar. Biol., Gdynia 1975, 245-247.
- JAŻDŹEWSKI, K., 1990. A redescription of *Tiron antarcticus* K.H. Barnard, 1932 (Crustacea: Amphipoda: Synopiidae) with an updated key to the species of *Tiron* Liljeborg, 1865. ____ Proc. biol. Soc. Wash. **103**, 110-119.

- KLAGES, M. & J. GUTT, 1990. Observations on the feeding behavior of the antarctic gammarid Eusirus perdentatus Chevreux, 1912 (Crustacea: Amphipoda) in aquaria. ____ Polar Biol. **10**, 359-364. ('A carnivorous predator'(!)).
- KÖHN, J. & F. GOSSELCK, 1989. The recent distribution of glacial relict Malacostraca in the western and southern Baltic. ____ Zool. Anz. **222**, 57-74.
- KÖHN, J. & A. WATERSTRAAT, 1990. The amphipod fauna of Lake Kummerow (Mecklenburg, German Democratic Republic), with reference to Echinogammarus ischnus Stebbing, 1899. ____ Crustaceana **58**, 74-82.
- KUSAND, H., 1989. Life history variation of a freshwater amphipod, Jesogammarus paucisetulosus. ____ Jpn. J. Limnol. **50**, 189-198.
- LALITHA, M., K. SHYAMASUNDARI & K.H. RAO, 1989. Effect of salinity and temperature on the littoral sand hopper Talorchestia marknssi (Weber) (Crustacea: Amphipoda). ____ Riv. Gdrobiol. **27**, (1988), 121-130.
- LALITHA, M., K. SHYAMASUNDARI & K.H. RAO, 1989. Studies on the embryonic development of Talorchestia martensii (Weber) (Crustacea: Amphipoda). ____ Arch. Ital. Anot. Embriol. **94**, 185-196.
- LAZO-WASEM, E.A., A.J. BALDINGER & M.F. GABLE, 1989. Pariphinotus Kunkel, 1910, the senior synonym of Heterophlia Shoemaker, 1933. (Crustacea: Amphipoda: Phliantidae). ____ Postilla **205**, 1-5.
- LAZO-WASEM, E.A. & M.F. GABLE, 1989. First report of a freshwater amphipod (Gammaridea: Hyalellidae), Hyalella azteca, new record (Saussure), from nonanchialine waters of Bermuda. Postilla **204**, 1-4. (No doubt introduced).
- LEDOYER, M., 1986. Faune mobile des herbiers de phanérogames marines (Halodule et Thalassia) de la Laguna de Termino' (Mexique, Campeche). 2. Les Gammariens. ____ An. Inst. cienc. Mar. Limnol. Univ. Natl. auton. Mexico **13**, 171-200. (Deals with Ampelisca vadorum, Gitanopsis laguna (with which G. petulans is tentatively synonymized), Cymadusa compta, Corophium cf. insidiosum, C. louisianum, Gammaropsis togoensis, Grandidierella bonnieroides, Lembos unicornis, Atylus minikoi, Nasageneia yucatanensis n.sp., Dulichella appendiculata, Elasmopus levis, Gammarus (Mucrogammarus) sp., Maera quadrimana, Melita planaterga, Cerapus benthophilus and Paraphoxus spinosus).
- LEIN, T.E., R. KÜFNER & J.R. HANSEN, 1989. (Plants and animals on the rocky shores of Finnmark. Consequences of oil pollution). ____ Ökofersk Repp. 1989 **15**, 1-56 (In Norwegian).
- LOP, A.E., 1989. On the morphological variation of two circum-mediterranean brackish - water (sic) gammarids Rhipidogammarus chipidophorus (Catta) and Echinogammarus foxi (Schelleberg) (sic), from the Spanish inland waters. ____ Spixiana **12**, 115-124.
- LOWRY, J.K. & H.E. STODDART, 1989. Stephonyx, a new, widespread genus of lysianassoid Amphipoda. ____ Zool. Scripta **18**, 519-525. (Stephonyx n. gen. has Euonyx biscayensis, as type, and five further spp., all transferred from Euonyx. E. chelatus and S. biscayensis are completely redescribed and reillustrated).
- MARGOLIS, L. & Z. KABATA, 1989. Guide to the parasites of fishes of Canada. Part 3. Acanthocephala and Cnideria. ____ Can. spec. Publ. Fish. aq. Sci. **107**, 1-90.
- MARRIOTT, D.R., M. L. COLLINS, R.M. PARIS, D.R. GUDGIN, C.J. BARNARD, P.K. MCGREGOR, F.S. GILBERT, J.C. HARTLEY & J.M. BEHNKE, 1989. Behavioural modifications and increased predation risk of Gammarus pulex infected with Polymorphus minutus. ____ J. biol. Educat. **23**, 136-141.
- MATEUS, A. & E. MATEUS, 1986. Campagne de la 'Calypso' dans le Golfe de Guinée et aux Iles Principe, São Tomé et Annobon (1976). Amphipodes récoltés à bord de la Calypso. ____ An. Fac. Cienc. Porto **66**, 125-133. (Deals with Waldeckia scrupulosa n. sp. (Guinea Bissau), Ampelisca acutidentata n. sp. (Principe), Leucothoe campi n. nom. (= L. denticulata auct., non Costa), Maera excavata n. sp. (?), M. trisinuata n. sp. (Principe), M. leopoldinae n. sp. (Portugal), M. atlantica n. sp. (Portugal), Elasmopus sachoni, E. spinipes n. sp. (?), Eurystheus aculeata n. sp. (Rio d'Óro), Amphithoe nobrei n. sp. (Annobon), A. dentimana n. sp. (Annobon), and Caprella acutifrons annobonensis n. ssp. (Annobon). Also contains a list of amphipods found in W. Africa).
- MCDONALD, J.H. 1987. Repeated geographic variation at three enzyme loci in the amphipod Platorchestia platensis. ____ Evolution **41**, 438-441.
- McLACHLAN, A., 1990 Dissipative beaches and macrofauna communities on exposed intertidal sands. ____ J. coast. Res. **6**, 57-71. (Not seen).
- MEADOWS, P.S. & J. TAIT, 1989. Modification of sediment permeability and shear strength by two burrowing invertebrates. ____ Mar. Biol. **101**, 75-82.
- MENIOU, M., M. DAKKI & P. AGUESSE, 1990. Une biotypologie des peuplements infralittoraux superficiels de crustacés des côtes rocheuses marocaines. ____ Vie Milieu **40**, 57-66. (Seventy-one spp of amphipods listed on p. 63).
- MESSANA, G. & L. CHELAZZI, 1986. The fauna of the subterranean waters of East Afric, and particularly of Somalie. ____ Stygologia **2**, 339-351 (Amph. 347-348).
- MEYER, E., 1990. A subsampling device for macroinvertebrates with general remarks on the processing of stream benthos samples. ____ Arch. Hydrobiol. **117**, 309-318.
- MUNRO, M.A., P.J. WHITFIELD & R. OFFLEY, 1989. Pomphorhynchus laevis (Müller) in the flounder, Platichthys flesus L., in the tidal river Thames: population structure, microhabitat utilization and reproductive status in the field and under conditions of controlled salinity. ____ J. Fish Biol. **35**, 719-736. (Viable larval stages in Gammarus zaddachi).

- NELL, J.M., M.H. BOTHNER, N.J. MACIOLEK & J.F. SPASSLE, 1989. Impact of exploratory drilling for oil and gas on the benthic environment of Georges Bank. ____ Mar. Environm. Res. 27, 77-114.
- NEWMAN, R.M., W.C. KERFOOT & Z. HANSCOM, 1990. Watercress and amphipods. Potential chemical defence in a spring stream macrophyte. ____ J. chem. Ecol. 16, 245-259.
- NIPPER, M.G., D.J. GREENSTEIN & S.M. BAY, 1989. Short-term and long-term sediment toxicity test methods with the amphipod Grandidierella japonica. ____ Environm. Toxicol. Chem. 8, 1191-1200.
- PECK, S.B. & K. CHRISTIANSEN, 1990. Evolution and zoogeography of the invertebrate cave fauna of the Driftless Area of the Upper Mississippi River Valley of Iowa, Minnesota, Wisconsin, and Illinois, USA. ____ Can. J. Zool. 60, 74-88.
- PETRESCU, I., 1989. Contribution to the study of the amphipods of Romania. Comparative study of the populations of Melita palmata (Montagu, 1804) (Crustacea, Amphipoda). ____ Trav. Mus. Hist. nat. Grigore Antipa 30, 187-191. (Material from Black Sea and English Channel).
- PIKE, D.G. & H.E. WELCH, 1990. Spatial and temporal distribution of sub-ice macrofauna in the Barrow Strait area, Northwest Territories, Canada. ____ Can. J. Fish. aq. Sci. 47, 81-91.
- PLATVOET, D., 1987. The genus Salentinella Ruffo, 1947 (Crustacea, Amphipoda), in Spain. ____ Stygologia 3, 217-240. (Deals with S. angelieri and its new variety S. a. longispina, S. carracensis n.sp. (prov. Guadalajara), S. seriliensis (prov. Sevilla), S. longicaudata n.sp. (prov. Valencia), S. cazemierae n.sp. (prov. Huseca), S. meijersae n.sp. (prov. Burges), and S. petiti. A key to Salentonella spp. is also provided).
- PLIGIN, Y.V. & L.V. YEMEL'YANOVA, 1989. Acclimatization of Caspian invertebrates in Dnieper reservoirs. ____ Hidrobiol. Zh. 25 (1), 1-9. (In Russian).
- PRETUS, J. L., 1988. A new stygobiont amphipod Pseudoniphargus mereadali n.sp., from the island of Minorca (Balearic archipelago). ____ Stygologia 4, 229-241.
- QUEIROSA, H., 1990. Corophium multisetosum (Amphipoda: Corophiidae) in Canal de Mira, Portugal: some factors that affect its distribution. ____ Mar. Biol. 104, 397-402.
- RAUSCHERT, M., 1988. Gammaridea (Crustacea, Amphipoda) from the coastal area of King George Island (South Shetland Islands): Podoceridae. ____ Mitt. Zool. Mus. Berlin 64, 299-310. (Deals with Podocerus capillimanus and Dulichia antarctica n.sp.).
- RAUSCHERT, M., 1990. Neue Stenothoidae (Crustacea, Amphipoda, Gammaridea) aus dem Sublitoral von King George (Süd-Shetland - Inseln). ____ Mitt. zool. Mus. Berlin 66, 3-39. (Deals with Metopoides antarcticus n. comb. (transferred from Proboloidea), M. macromanus n.sp., M. foliodactylus n.sp., M. andresi n.sp., M. lanceolatus n.sp., M. latus n.sp., M. serratus n.sp., M. angustus n.sp., M. leptomanus n.sp., and Prometopa edentata n.sp.).
- READ, A. T. & D.D. WILLIAMS, 1990. The role of the calceoli in precopulatory behaviour and mate recognition of Gammarus pseudolimnaeus Bousfield (Crustacea, Amphipoda). ____ J. nat. Hist. 24, 351-359.
- ROBINSON, A.M., J.O. LAMBERSON, F.A. COLE & R.C. SWARTZ, 1988. Effects of culture conditions on the sensitivity of a phoxocephalid amphipod, Rhepoxynius abronius, to cadmium in sediment. ____ Environm. Toxicol. Chem. 7, 953-959.
- RUSSO, A.R., 1990. The role of seaweed complexity in structuring Hawaiian epiphytal amphipod communities. ____ Hydrobiologia 194, 1-12.
- SA DE SA REGO, E., 1987. Allorchestes chelonitis Oliveira, 1953, a synonym of Parhyale hawaiiensis (Dana, 1853) (Crustacea, Amphipoda). ____ Gheringia, Zool. 66, 141-147.
- SALMAN, S.D. & N. JABBAR, 1990. A new species of the genus Cheiriphotis Walker, from the north-west Arabian Gulf, with a redescription of S. megacheles (Giles) (Amphipoda, Isaeidae). ____ Crustaceana 58, 214-226. (Cheiriphotis williamsoni n.sp. (= Ch. megacheles s. Walker, 1904) from Iraq).
- SCHEEPMACKER, M. & J. van DALFSEN, 1989. Genetic differentiation in Gammarus fossarum and Gammarus caparti (Crustacea, Amphipoda) with reference to Gammarus pulex pulex in northwestern Europe. ____ Bijdr. Dierk. 59, 127-140. (G. fossarum is genetically very heterogeneous while G. caparti is similar to some fossarum populations).
- SCONFIETTI, R., 1988. Research on spatial distribution of amphipods, isopods and tanaids (Peracarida) in a Mediterranean estuary (River Dese, Lagoon of Venice). ____ Crustaceana 55, 193-201.
- SHEA, J.R. & J.A. PERCY, 1990. Salinity tolerance and osmoregulation of the arctic marine amphipods, Onisimus littoralis (Kröyer) and Anonyx nugax (Phipps). ____ Polar Biol. 10, 275-281.
- SKADSHEIM, A., 1990. A cohort life table for Gammarus salinus (Amphipoda). ____ Oikos 57, 207-214.
- SMITH, S.J. & R. SWAIN, 1983. Observations on the taxonomy of Austrochiltonia (Hurley) (Amphipoda, Ceinidae). ____ Bull. austr. Soc. Limnol. 8, 39-43. (Afrochiltonia and Austrochiltonia are again considered valid genera).
- SOARES, C.M.A., ?? . (Preliminary note on the amphipods of the estuary of Itamaraca, Pe.). ____ Cienc. Cult. Suppl. 26, 356. (In Portuguese. Abstract only).

- SPICER, J.I., A.C. TAYLOR & B.R. McMAHON, 1990. O₂-binding properties of haemocyanin from the sandhopper Talitrus saltator (Montagu, 1808) (Crustacea: Amphipoda). ____ J. exp. mar. Biol. Ecol. 135, 213-228.
- STAPLETON, J.L., W.D. WILLIAMS & J.L. BARNARD, 1988. The morphology of the calceolus of an Australian crangonyctoid freshwater amphipod. ____ Crustaceana 55, 157-162.
- STOCK, J.H., 1988. Two new stygobiont Amphipoda (Crustacea) from Polynesia. ____ Stygologia 4, 79-100. (Fiha schminkei n. gen. n. sp. ('hadzioids') from Viti Levu, Fiji. The name Sriha n. nom. is introduced as replacement for preoccupied genus name Quadrus from Sri Lanka. Josephosella hamata n.sp. was collected from Tongatapu, Tonga).
- STOCK, J.H. & L. BOTOSANEANU, 1989. Discovery of the first bogidiellid (cavernicolous Amphipoda) east of Wallace's Line in Indonesia. ____ Stygologia 4 (1988), 371-377. (B. dehorrengi n.sp. from a cave on Halmohera).
- STOCK, J.H. & T. M. ILIFFE, 1990. Amphipod crustaceans from anchialine cave waters of the Galapagos Islands. ____ Zool. J. Linn. Sec. 98, 141-160. (Deals with Valettietta cavernicola n.sp., Galapsielus leleuporum and Antronicippe serrata n. gen., n. sp. (Pardaliscidae). Both new species are collected from Isle St. Cruz).
- STRELNIKOVA, V.M., 1989. (Body weight-length ratio and calory value in Antarctic hyperiid Parathemisto gaudichaudii) ____ Gidrobiol. Zh. 25(1), 107-105. (In Russian).
- VONK, R., 1989? Nuuanu curvata n.sp. and Melita leioteison n.sp. (Crustacea, Amphipoda) from beach interstitia on Curaçao. ____ Pp. 185-198 in Stud. in honour of Dr. Pieter Wagenaar Hummellinck 123. (Not seen. Reference incomplete?).
- WAKABARA, Y., A.S. TARARAM, M.T. VALERIO, B.E. RARDO & P.F. LEITE, 1988. Liljeborgiidae (Amphipoda - Gammaridea) from the southeastern coast of Brazil. ____ Relat. int. Inst. oceanogr., Univ. S. Paulo 23, 1-10. (Deals with Liljeborgia dubia, L. quinquidentata, and Listriella titinga n.sp. (Ilha Anchieta, SE Brazil)).
- WILLIAMS, J.A., 1990. The respiratory quotient of the high shore amphipod, Talorchestia deshayesi. ____ Comp. Biochem. Physiol. A 95, 177-180.
- WILSON, W.H., 1989. Predation and the mediation of intraspecies competition in an infaunal community in the Bay of Fundy. ____ J. exp. mar. Biol. Ecol. 132, 221-145. (Corophium volutator preyed upon by Semipalmated Sandpipers).

